Physical development of children is a marker of adequacy of nutrition, and affects adaptation, severity of pathological processes, rate of recovery, effectiveness of therapy, terms of hospitalization. Not enough teaching time is devoted to the study and training of nutrition status assessment skills in various ways.

The skill of evaluation of physical development (FR) and nutritional status (NS) was worked out on the questionnaire data of specific patients. The results of the programme WHO Antro assessment, the centyl tables and the impedance data were compared in 101 patients with eating disorders (54 boys), from 2 m.o to 17 y.o (average 8.5 y.o) Malnutrition 30%, severe malnutrition – 6% cases. Obesity revealed 39% kids. The results of the WHO Antro assessment, the percentile tables and the impedance data were compared. Direct correlation links between the child's percentile weight corridor, the WHO Antro (z-score), and the percentage of deficiency or excess of fat and active cell mass are obtained ($p=0.05$). The child's percentile growth corridor and body mass index also correlates with the growth z- scor, fat, and active cell mass percent according to body impedance ($p=0.05$).

The skill of assessing physical development and nutritional status by students can be formed using simulation technologies. Physical development, as well as the nutritional status of the child, can be investigated by the WHO Antro program using a computer application, or by using technologically complex equipment – impedance, or manually – with the help of centyl tables.

Deviations of nutritive status from physiological norm are determined by all proposed methods.

469

PREVENTIVE HEALTH CARE OF CHILDREN IN CROATIA DURING COVID-19 EPIDEMIC

Željka Draušnik*, Đurđa Španović, Ureljia Rodin, Alojz Radman. *Croatian Institute of Public Health*

10.1136/archdischild-2021-europaediatrics.469

To determine the impact of the COVID-19 epidemic on preventive health care visits and vaccinations in pediatrician practices in primary health care in Croatia.

Data about preventive health care visits and vaccinations of infants and preschool children in pediatrician practices in primary health care in Croatia were extracted from the Central Health Information System of the Republic of Croatia and comparison of 2019 and 2020 data was made, respectively month by month. We analyzed data only for infants and preschool children, from 0 to 6 years old.

In total, during 2019, 268,556 preventive health care visits of infants and preschool children were recorded compared to 252,690 during the 2020, which is decline of 5.9%. The biggest decline, 42.6%, was recorded during the March 2020, but with notable increase in the following months which almost overturned the impact of the initial decline. New decline in the visits was seen during the November, but far smaller than during the beginning of epidemic. Regarding vaccinations, during the 2019, 238,799 vaccinations were recorded to be administered, compared to 262,157 during the 2020, which is increase of 9.8%. The biggest decline was also seen during the March, 26.1%.

Beginning of COVID-19 epidemic in Croatia caused a major decline in the number of preventive health care visits and

vaccinations already during March 2020. It was probably direct consequence of 'lockdown' and parental fear of going to a health facility. Negligible is also the earthquake that hit Zagreb and surroundings in mid-March. Joint and rapid response of together primary pediatricians and Croatian Institute of Public Health with the thrust of the parents, manage to stop the decline of the numbers and even to increase the numbers in the following months. A second round of the decline in preventive health care visits, during the late autumn and the beginning of winter, was probably result of increase in the number of SARS-CoV- 2 positive people, including children, self-isolations and possible delay of vaccination after child is recovered from COVID-19.

However, this favorable situation with frequency of preventive health care visits and vaccinations in Croatia during the COVID-19 epidemic gives us hope that in the future Croatia will not face vaccine-preventable communicable diseases outbreak, as some countries expect and once more emphasizes the importance of the pediatricians in the primary health care for the well-being of infants and preschool children.

470

PARENTAL ASSESSMENT ON NEURODEVELOPMENTAL OUTCOMES IN CHILDREN BORN LATE PRETERM AT 2, 3, 5 AND 8 YEARS OLD

Diana Simão Raimundo*, Joana Mendão Carreira, João Vasco, Eulália Viveiros. *Hospital do Divino Espírito Santo, Ponta Delgada, Portugal*

10.1136/archdischild-2021-europaediatrics.470

In recent years, there has been growing evidence that children born late preterm (LPT) (34 0/7 – 36 6/7 weeks' gestation) are at higher risk for multiple morbidities during childhood and even adulthood. We aimed to evaluate late neurodevelopment in children born LPT at ages of 2, 3, 5 and 8 years old (y.o).

All LPT born in 2012, 2015, 2017 and 2018 in a Portuguese island region were selected from clinical records and a questionnaire was applied to their parents. Descriptive statistics analysis was performed.

Response rate was 67,0% (total= 207; 2 y.o. n=63; 3 y.o. n=62; 5 y.o. n=53; 8 y.o. n=29). Within LPT group, none of the evaluated outcomes recorded statistically significant differences between gestational ages. Visual and hearing impairment increased with age, until 55,2% and 26,0%, respectively, at 8y.o; this suggests that these potential problems may go unnoticed in the first years of life, becoming evident at school age. About 1/5 of the LPT presented difficulties in language/speech by 2 and 3 y.o., and it remained high at 5 years (11,3%), but with a lower rate at 8 years (3,4%). Warning signs for emotional/behavioral disturbance were reported in about 10% of 5 and 8 y.o., using the parameters recommended by the National Program for Child and Youth Health. Diagnosed attention deficit hyperactivity disorder (ADHD) was found in 6,9% of children aged 8, coincident with the prevalence estimated worldwide. However, 17,2% of all 8 y.o. LPT showed warning signs on Conners' Parent Rating Scale, suggesting this clinical entity may be underdiagnosed or that parental reports may indicate behavior dysfunctionality despite criteria for attention deficit hyperactivity disorder (ADHD) are not met. School learning difficulties with grade retention were reported in 17,2% of the 8 y.o. LPT children, much higher than the national mean retention rate in the 2nd grade (5,8%). These