

Supplemental table 3 Univariable and multivariable Cox proportional hazards model of all patients

	Death or transplant			Reintervention			Aortic valve replacement		
	Uni		Multivariable	Uni		Multivariable	Uni		Multivariable
	p	p	HR (CI 95%)	p	p	HR (CI 95%)	p	p	HR (CI 95%)
Critical stenosis, yes/no	0.10	0.58	2.7 (0.1-42.6)	0.007	0.31	1.6 (0.6-4.1)	0.09	0.56	0.7 (0.2-2.3)
Gender, male/female	0.41			0.9			0.44		
Gestational age, weeks	0.41			0.024	0.55	0.6 (0.5-0.9)	0.024	0.67	1.0 (0.8-1.2)
Birth weight	0.86			0.54			0.10		
Aortic annulus, z-score	0.1			<0.001	0.007	0.6 (0.5-0.9)	0.024	0.002	0.5 (0.3-0.8)
FFE	0.025	0.38	2.7 (0.3-24.9)	0.013	0.13	0.4 (0.1-1.3)	0.057	0.26	0.4 (0.1-1.8)
LVEDd, z-score	0.19			0.56			0.69		
LVPWd, z-score	0.25			0.30			0.15		
Mean aortic gradient	0.034	0.4	1.0 (0.9-1.9)	0.55			0.91		
Residual mean gradient	0.87			<0.001	1.1	1.1 (1.0-1.1)	0.053	0.03	1.1 (1.0-1.1)
Valve morphology	0.16			0.43			0.48		
Tricuspid (ref)									
Bicuspid	0.058			0.58			0.82		
Unicuspid	NA			0.19			0.24		
AR post first intervention							0.19		
No AR (ref)									
AR grade 1	0.84			0.76			0.56		
AR grade 2 or 3*	0.87			0.63			0.075		

Bold indicates p-values <0.1. Pre- and post-treatment aortic valve gradients were measured with echo Doppler.

*No patient had AR grade 4. (AS, aortic stenosis; HR, hazard ratio; CI, confidence interval; LVEDd, left ventricle end diastolic diameter; LVPWd, left ventricle posterior wall diameter; AR, Aortic regurgitation.)