

THE EARLY DIAGNOSIS OF RICKETS

BY

J. D. GRAY, M.D., and F. STEPHEN CARTER, M.D., B.Chir.D.C.H.
(From the Belgrave Hospital for Children, London)

With the widespread application of the knowledge of the relationship of vitamin D to rickets, the occurrence of the clinical picture familiar from Glisson's time onwards has nearly disappeared. In its place we are faced with changes brought about by a suboptimal intake of the necessary vitamin. Unfortunately the physical signs on which a diagnosis depends mimic the wide variation of normal to be found in infants. Cecil (1940) thus sums up the position: 'Clinical appraisal of the departure from normal in the rib-cartilage junction is difficult. A common clinical error is to diagnose rickets where no rickets is present.' The reverse is true. The same statement is applicable to the closing of the fontanelle, the eruption of teeth, and the width of the lower end of the radius. 'The clinical diagnosis of rickets based upon enlarged epiphyses and bow-legs is often unreliable.' (Price, 1946).

The radiological appearance of the lower end of the radius and ulna is often accepted as the arbiter. This is open to question, for the British Paediatric Association (1944) in an enquiry into the incidence of rickets in war-time, using a panel of experienced paediatricians and radiologists, found a clinical incidence of 12.5 per cent. to a radiological 0.5 per cent. Rackow (1949) says:

'The wrist joint in the infant is generally accepted by radiologists as being the most satisfactory for x-ray study in the diagnosis of rickets. It offers no technical difficulties in the making of the radiograph and when changes in the bone have occurred they are invariably present at the lower end of the radius and ulna. Being a small joint, all parts remain comparatively close to the film, so that as long as movement is excluded, the definition of bone structure is sharp.

'The florid and active form of rickets is readily diagnosed radiologically, and the diagnostic problem now mainly concerns itself with slighter degrees of rachitic change.

'The early radiological changes in the bone are mainly changes of degree and not of kind. Thus, slight expansion of the diaphyses of the radius and ulna is already present at the lower end of the normal bone. The ulna also normally shows slight cupping; osteoporosis may be simulated by slight over-penetration of the radiograph. For these reasons, and because radiological change of any kind in bone, which is due to systemic disorder, may take several weeks to establish itself, the early radiological diagnosis of rickets may be

uncertain and many cases can only be classed as doubtful.'

The plasma-alkaline-phosphatase has been accepted by many observers as a reliable means of detecting active and latent rickets (Cantarow and Trumper, 1945). In this respect, most investigators have used Bodansky's method for determining the phosphatase content of the blood in rickets. It has two disadvantages for routine use in paediatrics in that it requires well over an hour for a determination and it needs 2 ml. of serum. On the other hand, the King-Armstrong method can be completed in twenty-five minutes and requires only 0.4 ml. plasma. An alternative argument in favour of Bodansky's method is that it combines an estimation of the serum inorganic phosphate as well as the phosphatase. However, it has been our experience that the inorganic phosphate in the majority of cases of early rickets is within normal limits although the phosphatase is increased. Barnes et al. (1946) noted that in 40 per cent. of their cases of rickets the calcium and phosphorus levels were normal. Josefsson (1941) has shown that active rickets may be present in spite of normal calcium and phosphorus levels, but in the healing stage the phosphorus level may be low. Thus, there seems to be considerable evidence that the blood calcium phosphorus product is unreliable as a means of detecting early rickets. Because of the arguments presented here, the plasma alkaline phosphatase has been estimated by the King-Armstrong method in all cases of clinically suspected rickets seen at the Belgrave Hospital for Children during the last two and a half years. The normal range is 0-10 units in the adult. Little information was available as to the normal and pathological ranges to be found in young children, for only two references were found which had a bearing on rickets. May and Wygant (1939) considered that rickets could not be diagnosed under 55 King-Armstrong units of phosphatase activity, and on the other hand Corner (1944) put the normal range at 3-10 units with no relation to age, and suggested that readings of 15 units or over were diagnostic of rickets. In our experience, the first was too high and the latter too low. An attempt, therefore, has been made to assess the normal range, to relate the clinical condition to King-Armstrong units of phosphatase activity, and to use the phosphatase content as a guide to treatment.

TABLE 1
ANALYSIS OF CASES OF CLINICAL RICKETS SHOWING PLASMA PHOSPHATASE LEVELS AND RESPONSE TO VITAMIN D

No.	Sex	Vitamin D	Age	Signs	Phosphatase Units		Treatment	Remarks
					Date	No.		
1	F.	Given†	1 yr. 7 months	S+S+*	16 Jan., 1947 19 Feb., 1947 19 Mar., 1947	25 51 14	Calciferol 200,000 units	Radiograph showed epiphyses rachitic
2	F.	?	11 months	B+F1	21 Nov., 1946 28 Feb., 1947	23 18	Radiostoleum 5 ml. daily	
3	M.	None	5 months	B+C+E+	4 Dec., 1946 31 Mar., 1947	24 20	Calciferol 200,000 units	
4	M.	?	1 yr. 3 months	B+S+	4 June, 1947 23 Aug., 1947 5 Jan., 1948 5 April, 1948	30 23 20 17	Adexolin 30 m. daily	
5	M.	?	1 yr. 7 months	B+Bs	7 May, 1947 7 July, 1947	76 17	Calciferol 200,000 by 3	Radiograph showed epiphyses rachitic
6	M.	Given	1 yr. 11 months	B+F+C+	26 July, 1948 1 Sept., 1948	64 29	Calciferol 200,000 units	
7	M.	?	1 yr. 6 months	B+Bs	26 Jan., 1948 1 Mar., 1948	36 21	Calciferol 200,000 units	Radiograph normal
8	M.	?	1 yr. 5 months	B+F+	28 June, 1948 29 July, 1948	38 25	Calciferol 200,000 units	Radiograph normal
9	F.	?	2 yrs. 6 months	B+	16 Jan., 1947 13 Feb., 1947	25 19	Calciferol 200,000 units	
10	M.	?	8 months	B-F+C+	30 Sept., 1946 25 Oct., 1946 24 Nov., 1946	29 42 46	Ostelin 30 m. daily	Radiograph showed recalcifying rickets
11	M.	?	3 yrs. 6 months	Bs	12 July, 1948 4 Oct., 1948	30 20	Calciferol 200,000 units	
12	M.	?	1 yr. 3 months	B-S+F1	29 Apr., 1948 16 June, 1948 13 July, 1948 18 Oct., 1948 20 Dec., 1948	30 42 58 37 18	Calciferol 200,000 by 3	
13	M.	?	1 yr. 11 months	F-Bs	11 Aug., 1948 6 Oct., 1948	44 34	Cod-liver oil dr. 2 daily	
14	M.	Given	1 yr. 8 months	B-B-Bs	6 Aug., 1948 5 Oct., 1948 16 Nov., 1948 28 Dec., 1948	51 30 38 39	Calciferol 200,000 by 3	
15	F.	?	1 yr. 7 months	B-B-BsS+	6 Oct., 1948 15 Nov., 1948 16 Dec., 1948	42 43 22	Calciferol tab. co. t.d.s. (500 units, Vit. D per tablet)	
16	F.	?	1 yr. 1 month	B+E+BsF1	11 Nov., 1948 15 Dec., 1948	38 16	Calciferol 200,000 by 2	
17	M.	?	1 yr. 5 months	B+E+F1	4 Jan., 1948 8 Feb., 1948	46 17	Calciferol 200,000 units	
18	M.	Given	1 yr. 3 months	B+B+	25 Aug., 1948 6 Dec., 1948	58 22	Calciferol 200,000 by 2	
19	M.	Given	1 yr. 9 months	B+E-F-	7 May, 1948 9 June, 1948	32 22	Calciferol 200,000 units	Radiograph showed ? rickets
20	F.	Given	1 yr. 2 months	E+F+	12 Mar., 1947 7 May, 1947 16 June, 1947	36 23 17	Calciferol 200,000 units	Radiograph showed healing rickets
21	F.	Given	1 yr. 4 months	B+Bs		45		Radiograph showed epiphyses rachitic
22	F.	None	1 yr. 4 months	F+F1		33		
23	F.	Given	1 yr.	B+F+F1		49		
24	F.	?	1 yr. 8 months	B+		49		
25	M.	Given	7 months	F+C+F1		36		
26	F.	?	2 yrs. 3 months	B+BsF1		37		

TABLE 1—*cont.*

No.	Sex	Vitamin D	Age	Signs	Phosphatase Units		Treatment	Remarks
					Date	No.		
27	F.	None	11 months	B-E-C--		32		
28	M.	None	1 yr. 4 months	B+BsC--		32		
29	F.	None	7 months	B+		30		
30	F.	?	1 yr.	B-FI		27		Radiograph showed epiphyses rachitic
31	M.	None	2 yrs. 3 months	E-FI		30		
32	M.	Given	1 yr. 6 months	B+		26		Radiograph showed epiphyses rachitic
33	M.	Given	4 months	B+C-		26		
34	M.	None	1 yr. 3 months	E+B _s		30		
35	M.	?	2 yrs.	E+B _s FI		32		
36	F.	?	2 yrs.	B+E+B _s		26		Radiograph normal
37	M.	None	2 yrs. 1 month	B-FI		32		
38	M.	?	1 yr. 3 months	B+E+		29		
39	M.	Given	1 yr. 3 months	B+F+		29		
40	F.	None	2 yrs. 6 months	B-BsF-		31		
41	M.	?	1 yr. 9 months	B-E+		33		
42	F.	?	1 yr. 6 months	B+B _s		32		
43	M.	?	9 months	B-F+FI		27		
44	M.	None	4 months	FI		37		Radiograph showed epiphyses rachitic
45	M.	Given	1 yr. 8 months	B-F+FI		54		Radiograph showed rachitic changes
46	F.	?	6 months	B+E-FI		30		Radiograph showed epiphyses rachitic

* In the tables:

B+ = beading at the costochondral junction. This was assessed as far as possible on the recommendations of Dalyell and Mackay (1922), and beading has not been indicated unless it could be graded 1½-2 by their standards.

B_s = bossing of the skull.

C+ = craniotabes.

E+ = enlarged epiphyses at the wrists.

S+ = Harrison's sulcus.

F+ = a fontanelle larger than age permits.

FI = muscle tone poor.

† Where the mother was insistent that cod-liver oil had been given it is stated as 'given'; doubtful evidence has been indicated by a question mark.

Method

The plasma phosphatase was determined in forty-six apparently healthy children from one month to three years old with no clinical signs of rickets. The average obtained for this group was 17 King-Armstrong units with a spread of 11-20. Ten children of three to five years averaged 12 units with a spread of 8-18 units. In thirteen of the controls picked at random, radiographs of radial and ulnar epiphyses appeared normal.

We have not included three cases of florid rickets in which the phosphatase levels fell between 80 and 220 units as diagnostic confirmation was not required.

One of us (F.S.C.) is responsible for the clinical diagnoses in all the cases reported.

Discussion

The contention that rickets cannot be diagnosed under 55 King-Armstrong units of phosphatase activity is not borne out by the findings in the series of cases presented. In most the diagnostic titre lay between 30 and 40 units, and with the institution of specific therapy the response was prompt and marked. On the other hand, it has not been possible to substantiate clearly Corner's claim that rickets could be related to findings in the 15-20 units range. Setting aside the control values given in this paper and considering the problem from different aspects, the following points are put forward to strengthen the view that her figures for normality are too low for the first two years of life. Barnes and Munks (1940) in establishing values for infants

TABLE 2
ANALYSIS OF CASES WITH SYMPTOMS BUT NO SIGNS

No.	Sex	Vitamin D	Age	Signs	Phosphatase Units		Treatment	Remarks
					Date	No.		
1	M.	Given	1 yr. 2 months	Irritable	20 Oct., 1948	35	Tab. Calciferol Co. t.d.s. (5 units, Vit. D. per tab.)	
					19 Nov., 1948	27		
2	F.	None	1 yr. 9 months	Irritable	14 Oct., 1948	29	Tab. Calciferol Co. t.d.s. (5 units, Vit. D. per tab.)	
					30 Nov., 1948	23		
					25 Jan., 1949	23		
3	F.	None	1 yr. 8 months	Irritable Sweats	27 May, 1948	38	Calciferol 200,000 by 3	Radiograph normal
					25 June, 1948	24		
					28 July, 1948	45		
					29 Oct., 1948	35		
					27 Jan., 1949	17		
4	M.	?	1 yr. 4 months	Head sweats	26 May, 1948	33	Calciferol 200,000 by 5	Radiograph showed ? rickets
					30 June, 1948	36		
					26 July, 1948	35		
					23 Aug., 1948	42		
					20 Sept., 1948	34		
					20 Jan., 1949	26		
5	M.	None	9 months	No symptoms	18 Nov., 1946	33	Radiostoleum 5 ml. daily	Radiograph showed epiphyses rachitic Radiograph showed epiphyses normal
					1 Jan., 1947	15		
					23 June, 1947	17		
6	F.	None	6 months	Irritable	24 Aug., 1948	31	Calciferol 200,000 units	
					23 Sept., 1948	16		
7	M.	?	11 months	Irritable	9 July, 1947	27	Adexolin m. 10 daily	
					7 Apr., 1948	20		
8	F.	None	4 months	Irritable Head sweats	25 Mar., 1948	28	Adexolin m. 10 daily	
					11 May, 1948	20		
9	M.	None	6 months	Irritable	20 May, 1948	36	Calciferol 200,000 by 4	Radiograph normal
					23 June, 1948	58		
					13 Oct., 1948	24		

TABLE 3
ANALYSIS OF CASES WITH SYMPTOMS AND A NORMAL PHOSPHATASE

No.	Sex	Vitamin D	Age	Signs	Phosphatase Units		Treatment	Remarks
					Date	No.		
1	M.	Given	7 months	Irritable Now showed B-F1	10 Aug., 1948	24	No treatment given Calciferol 200,000 by 3	Radiograph normal
					15 Nov., 1948	54		
2	F.	Given	1 yr. 3 months	Delayed dentition F-?	12 Dec., 1947	20	No treatment given Adexolin ml. 5 daily	Radiograph normal
					16 June, 1948	32		
					1 Sept., 1948	60		
					27 Oct., 1948	24		
					24 Nov., 1948	22		

in Bodansky units have shown that phosphatase activity rises from 7.1 units on the third day to 13.0 units by the fourth or fifth month, and then gradually declines to 11.5 units by the end of the third year, the normal adult range being 1.5-4 units.

Klasmer (1944) in his controls found a variation of 9-15 Bodansky units in children ranging in age from six months to two and a half years. It is not possible to convert Bodansky to King-Armstrong units, but as they both measure phosphatase activity, it is to be expected that the variation in one should be reflected by the other. If Corner's figure of up to 10 ± 5 units normal range for the infant is accepted, then it shows very little difference from the adult normal and does not mirror the rise which other observers have noted for Bodansky's method.

Further, the greatest rate of growth in length in a child, about eighteen inches, occurs during the first three years of life, dropping to less than half that figure for the next three years. It seems reasonable to expect the tremendous epiphyseal activity to be paralleled by a phosphatase activity greater than that found in the static adult. It is thought therefore that the figure of 20 units as the upper limit of normal for the first three years of life given by the control series in this paper is approximately correct.

A phenomenon discussed by Yieh and Wisler (1938) as well as by Josefsson is seen in some of the cases in this series, namely in table 1 nos. 1, 10, 12; in table 2 nos. 3, 4, 9; and in table 3, no. 2. With the institution of vitamin treatment, particularly of the

TABLE 4
ANALYSIS OF CASES CLINICALLY DIAGNOSED AS RICKETS WITH NO BIOCHEMICAL CONFIRMATION

No.	Sex	Vitamin D	Age	Signs	Phosphatase Units		Treatment	Remarks
					Date	No.		
1	M.	Given	7 months	B-C \pm	7 June, 1947	20	Calciferol 50,000 by 3	Radiograph showed epiphyses rachitic Radiograph showed improvement
					4 Feb., 1948	16		
2	M.	Given	1 yr.	Bs	17 Mar., 1948	31	Calciferol 100,000 by 3 Calciferol 200,000 units	Radiograph showed healing Radiograph normal
					9 Apr., 1948	20		
					21 July, 1948	13		
3	M.	Given	8 months	E-S	28 Oct., 1947	18	Calciferol 200,000 by 3	
					31 Dec., 1947	20		
					11 Feb., 1948	20		
4	F.	None	1 yr. 6 months	B-E \pm	14 Nov., 1947	20		
5	F.	None	2 yrs.	B-BsE \pm	25 Feb., 1947	21		
6	M.	Given	1 yr. 2 months	B-S	9 May, 1947	22		Radiograph showed ? rickets
7	M.	Given	9 months	B-Bs	25 Mar., 1947	21		
8	F.	Given	1 yr. 5 months	B-BsE \pm	28 Apr., 1947	22		Radiograph showed slight rachitic change
9	M.	Given	8 months	E-		16		Radiograph showed rachitic changes present
10	M.	None	1 yr. 1 month	B-E \pm		13		Radiograph showed healing rickets

shock variety, there is first a fall in phosphatase activity, then a secondary rise, finally a slow fall to normal. This is assumed to be associated with the process of healing in more advanced cases. The usual reaction in milder types is a prompt drop to a point either near or within the normal range.

Dikshit and Patwardhan (1946) working on experimental rickets in puppies noted that the phosphatase started to rise slowly from the very beginning of vitamin D deficiency. This is illustrated in table 2 by cases 1-9 when symptoms which might have been due to early rickets were present. None of the usual stigmata were found on physical examination. The phosphatase activity averaged 35 units. It is interesting to note that the phosphatase findings in these very early cases were in the main in keeping with those which had definite signs. It is suggested that although a rise in phosphatase activity is diagnostic of early rickets, it cannot be used in this type of case as an index of the severity of vitamin D deficiency, confirming a similar finding of Morris et al. (1937).

Cases 1 and 2 (table 3) demonstrate the necessity of checking the phosphatase at frequent intervals when the diagnosis is entertained but the biochemical findings are equivocal. Clinical judgment had been swayed by the reported phosphatase level. Consequently no treatment had been given, with the result that three and six months later both biochemical and clinical rickets were present in each case.

Cases 1-10 (table 4) require separate consideration. Case 3 is probably an example of healed rickets in which the physical signs remain but the phosphatase level has returned to normal. In this case, large

doses of 'calciferol' had no effect on the enzymatic activity. This has been suggested as diagnostic of healed rickets by Klasmer (1944). Cases 4-6 are difficult to classify because of a lack of fuller information. They could be cases of active rickets with a borderline phosphatase level, or of healed rickets with persistent physical signs. A dose of 200,000 units of calciferol and a phosphatase estimation three weeks later would probably have settled the point. This reaction is illustrated by cases 1 and 2. Case 8 is probably borderline, and for nos. 9 and 10 no explanation is offered.

No attempt has been made to correlate radiological and biochemical findings, a difficulty experienced by Bodansky and Jaffe (1934) and appearing in this series as well. They found cases with a normal phosphatase content and radiological evidence of rickets, and a review of the cases presented here shows the same inconsistency. Radiological diagnosis of early rickets is renowned for its difficulty, and is one in which error is great even among the most highly skilled observers. The limitations of the radiologist in this respect are noted by Barnes and Carpenter (1937), and it is felt that too much reliance should not be based on x-ray examinations for diagnostic purposes in those cases where, at best, a radiograph can only be expected to show minimal changes.

Summary

An estimation of the plasma-alkaline-phosphatase by the King-Armstrong method in 56 controls gives a range of 11-20 units with an average of 17 for

children between one month and three years of age. Over that age, the average approximates to the figure for adults.

In sixty-four cases of clinical rickets, blood phosphatase levels were evaluated, many of them serially before and after treatment. The most frequent diagnostic titre lay between 25-40 units.

The prompt response to specific therapy is shown. The raised blood phosphatase level due to shock therapy has been noted and discussed. The phosphatase level in very early cases showing symptoms but no signs averages 35 units.

The difficulties of diagnosis in borderline cases are mentioned, and it is suggested that a dose of calciferol, 200,000 units, followed by a further blood phosphatase estimation three weeks later may be of assistance in solving the problem. The difficulties of early radiological diagnoses are stressed, and it is suggested that the discrepancies which occur occasionally in cases of clinical rickets and a normal phosphatase may be due to the persistence of physical signs in healed rickets.

We are indebted to Dr. A. M. Rackow for his assistance in discussing the radiology of rickets.

REFERENCES

- Barnes, D. J., and Carpenter, M. D. (1937). *J. Pediat.*, **10**, 596.
- , Kaucher, M., and Munks, B. (1946). *Amer. J. Dis. Child.*, **71**, 622.
- , and Munks, B. (1940). *Proc. Soc. exp. Biol.*, **44**, 327.
- Bodansky, A., and Jaffe, H. L. (1934). *Arch. intern. Med.*, **54**, 88.
- Brit. Paed. Assoc. (1944). *Arch. Dis. Child.*, **19**, 54.
- Cantarow, A., and Trumper, M. (1945). *Clin. Biochem.* 3rd ed., 198. Philadelphia.
- Cecil, R. L. (1940). *Text Book of Med.*, 5th ed., 645. Philadelphia.
- Corner, B. D. (1944). *Arch. Dis. Childh.*, **19**, 68.
- Dalyell, E. J., and Mackay, H. M. M., M.R.C. (1922). *Spec. Rep. Ser.*, **77**, 137.
- Dikshit, P. K., and Patwardhan, V. N. (1946). *Indian J. med. Res.*, **34**, 263.
- Josefsson, E. (1941). *Ann. Paediatr.*, **157**, 169.
- Klasmer, R. (1944). *Amer. J. Dis. Child.*, **67**, 348.
- May, E. W., and Wygant, T. M. (1939). *Arch. Pediat.*, **56**, 274.
- Morris, N., and Peden, O. D. (1937). *Quart. J. Med.*, n.s. **6**, 211.
- , Stevenson, M. M. and Small, J. M. D. (1937). *Arch. Dis. Childh.*, **12**, 45.
- Price, F. (1946). *Text Book of Prac. Med.*, 7th ed., 486 et seq.
- Rackow, A. M. (1949). *Personal communication.*
- Yieh, V., and Wissler, O. (1938). *Ann. Paediatr.*, **152**, 348.