

# MARMITE

yeast extract

## nutrition in infancy

contains  
RIBOFLAVIN (vitamin B<sub>2</sub>) 1.5 mg. per oz.  
NIACIN (nicotinic acid) 16.5 mg. per oz.

Malnutrition in infancy presents a serious problem in many tropical countries. Nutritional deficiencies during this period of rapid growth not only cause morbidity among young children, but also undermine the health of the individual in later life. Anaemia, marasmus and kwashiorkor are some of the common nutritional disorders of early childhood. Although lack of protein is considered responsible for much ill-health among infants, vitamin deficiencies are also prevalent.

Marmite yeast extract supplies every known member of the vitamin B complex and, within the limits of the amount consumed, it is a useful source of predigested protein. It can be given to babies from birth and has been particularly recommended for infants that are not thriving.

*Literature on Request*

MARMITE LIMITED, WALSINGHAM HOUSE, SEETHING LANE, LONDON, E.C.3  
CA 5609

### ~~~~~ J. & A. CHURCHILL LTD ~~~~~

#### DISEASES OF INFANCY AND CHILDHOOD

By WILFRID SHELDON, C.V.O., M.D., F.R.C.P.  
Seventh Edition. 213 Text-figures and including 18 Plates  
(5 in colour). 50s.

#### INFANT FEEDING AND FEEDING DIFFICULTIES

By P. R. EVANS, M.D., F.R.C.P., and RONALD MacKEITH, D.M., F.R.C.P. Second Edition. 66 Illustrations. 12s. 6d.

#### PROGRESS IN CLINICAL OBSTETRICS AND GYNAECOLOGY

By T. L. T. LEWIS, M.D., F.R.C.S., M.R.C.O.G. 90 Illustrations. 55s.

#### RECENT ADVANCES IN PAEDIATRICS

By various authors under the Editorship of DOUGLAS GAIRDNER, D.M., F.R.C.P. 117 Illustrations. 42s.

#### ANTENATAL AND POSTNATAL CARE

By F. J. BROWNE, M.D., D.Sc., F.R.C.S.Ed., F.R.C.O.G., and J. C. McCLURE BROWNE, M.B., B.S., F.R.C.S.Ed., F.R.C.O.G. Eighth Edition. 94 Illustrations. 37s. 6d.

#### PHYSIOTHERAPY IN PARAPLEGIA

By E. P. G. HOBSON, F.C.S.P. 30 Illustrations. 12s. 6d.

#### CHILD HEALTH AND DEVELOPMENT

By Various Authors. Edited by RICHARD W. B. ELLIS, O.B.E., M.A., M.D., F.R.C.P. 81 Illustrations. 42s.

Second (Enlarged) Edition.

#### THE QUEEN CHARLOTTE'S TEXTBOOK OF OBSTETRICS

By MEMBERS OF THE CLINICAL STAFF OF THE HOSPITAL. New (Ninth) Edition. 4 Coloured Plates and 230 Text-figures. 45s.

#### TUBERCULOSIS IN OBSTETRICS AND GYNAECOLOGY

By GEORGE SCHAEFER, M.D., F.A.C.S., F.I.C.S., Cornell University Medical College. 58 illustrations. 63s.

#### ESSENTIALS OF ORTHOPAEDICS

By PHILIP WILES, M.S., F.R.C.S., F.A.C.S. Second Edition. 7 Coloured Plates and 393 Text-figures. 55s.

#### COMMON DISEASES OF THE EAR, NOSE AND THROAT

By PHILIP READING, M.S., F.R.C.S. Second Edition. 2 Coloured Plates and 38 Text-figures. 22s. 6d.

#### THE NORMAL CHILD

Some Problems of the First Three Years and Their Treatment. By R. S. ILLINGWORTH, M.D., F.R.C.P. 64 Illustrations. 30s.

#### BABIES AND YOUNG CHILDREN

Feeding, Management and Care. A Book for Parents. By R. S. ILLINGWORTH, M.D., F.R.C.P., D.C.H., and CYNTHIA M. ILLINGWORTH, M.B., B.S., M.R.C.P. 24 Plates and 50 Text-figures. 18s.

~~~~~ 104 GLOUCESTER PLACE, LONDON, W. ~~~~~

# For the medical treatment of hypertrophic pyloric stenosis

## SKOPYL

BRAND OF METHYL SCOPOLAMINE NITRATE

### a powerful antispasmodic without central depressant action

**SKOPYL\*** is one of the most powerful spasmolytics known, the effect in vitro being five times greater than that of atropine, and ten times greater than that of scopolamine. The toxicity is three times that of scopolamine. Because of the greater activity the dose and the relative toxicity are correspondingly less.

**SKOPYL** has been in clinical use for about ten years. Comprehensive experience has shown the preparation to be especially valuable in all conditions where spasmolysis and inhibition of secretion is desirable. Side-effects are rare and usually very slight, even when large doses are used.

**IT IS** in paediatrics that **SKOPYL** is most useful, and the value of the preparation is best seen in the treatment of hypertrophic pyloric stenosis, where it has been proved able to replace surgery in many cases. Good results have also been obtained in infantile dyspepsia, and during the spasmodic period in pertussis.

For clinical references and directions for dosage, see a special booklet that will be sent on request.

**SKOPYL** is available in tablets of 0.5 mg. and 0.1 mg. (**SKOPYL MITE**), and in 0.25% w/v solution for sublingual use.



PHARMACIA Ltd.

UPPSALA, SWEDEN

*represented by***SAVORY & MOORE, LTD.**

61 Welbeck Street, London, W.1

\* Trade Mark