

Gluten-free

Diets

Coeliac disease in children and certain cases of idiopathic steatorrhoea in adults have been found amenable to treatment with a gluten-free diet. As patients suffering from the coeliac syndrome remain intolerant to gluten for long periods, wheat and rye flour which contain gluten may have to be excluded from the diet for an indefinite time. The withdrawal of all sources of wheat and rye flour results in a severely restricted diet and special care must be taken to ensure an adequate intake of essential nutrients.

Marmite is recommended for inclusion in the diet of coeliac patients. It supplies all known factors of the vitamin B complex which, in the absence of flour, may be in short supply. Moreover, the appetite-stimulating properties of Marmite are of importance to patients on a restricted dietary regime.

MARMITE

yeast extract contains RIBOFLAVIN (vitamin B₂) 1.5 mg. per oz. NIACIN (nicotinic acid) 16.5 mg. per oz.

MARMITE LIMITED, 35 SEETHING LANE, LONDON, E.C.3

ADC 581c

~~~~~ J. & A. CHURCHILL LTD. ~~~~~

New (Second) Edition

RECENT ADVANCES IN PAEDIATRICS

Just Ready

Edited by DOUGLAS GAIRDNER, D.M., F.R.C.P.
82 illustrations. 48s.

INFANT FEEDING AND FEEDING DIFFICULTIES

By P. R. EVANS, M.D., F.R.C.P., M.Sc. and RONALD MacKEITH, D.M., F.R.C.P., D.C.H.
New (Third) Edition. 66 illustrations. 16s.

CHILD HEALTH AND DEVELOPMENT

By various authors. Edited by R. W. B. ELLIS, O.B.E., M.A., M.D., F.R.C.P.
Second Edition. 81 illustrations. 42s.

THE PREMATURE BABY

By V. MARY CROSSE, O.B.E., M.D., D.Obst.R.C.O.G., D.P.H.
Fourth Edition. 39 illustrations. 20s.

DISEASES OF INFANCY AND CHILDHOOD

By WILFRID SHELDON, C.V.O., M.D., F.R.C.P.
Seventh Edition. 213 text-figures and including 18 plates (5 in colour). 50s.

A New Book

RECENT ADVANCES IN CEREBRAL PALSY

Just Published

Edited by RONALD S. ILLINGWORTH, M.D., F.R.C.P., D.P.H., D.C.H.
136 illustrations. 50s.

THE NORMAL CHILD: Some Problems of the First Five Years and Their Treatment

By R. S. ILLINGWORTH, M.D., F.R.C.P., D.C.H.
Second Edition. 69 illustrations. 33s.

BABIES AND YOUNG CHILDREN: Feeding, Management and Care

By R. S. ILLINGWORTH, M.D., F.R.C.P., D.C.H., and CYNTHIA M. ILLINGWORTH, M.B., M.R.C.P.
24 plates and 50 text-figures. A book for Parents. 18s.

ESSENTIALS OF ORTHOPAEDICS

By PHILIP WILES, M.S., F.R.C.S., F.A.C.S.
Second Edition. 7 coloured plates and 393 text-figures. 55s.

PRINCIPLES OF EPIDEMIOLOGY

By IAN TAYLOR, M.D., M.R.C.P., D.P.H., and JOHN KNOWELDEN, M.D., D.P.H.
25 illustrations. 30s.

A SYNOPSIS OF HYGIENE (Jameson & Parkinson)

New (Eleventh) Edition.
By LLYWELYN ROBERTS, M.D., M.R.C.P., D.P.H., assisted by KATHLEEN M. SHAW, M.B.E.
22 illustrations. 60s.

RECENT ADVANCES IN OBSTETRICS AND GYNAECOLOGY

By A. W. BOURNE, M.A., M.B., F.R.C.S., F.R.C.O.G., and L. H. WILLIAMS, M.D., M.S., F.R.C.S., F.R.C.O.G.
Ninth Edition. 78 illustrations. 38s.

~~~~~ 104 GLOUCESTER PLACE, LONDON, W.1 ~~~~~

# For the medical treatment of hypertrophic pyloric stenosis

## SKOPYL

BRAND OF METHYL SCOPOLAMINE NITRATE

a powerful antispasmodic  
without central depressant action

**SKOPYL\*** is one of the most powerful spasmolytics known, the effect in vitro being five times greater than that of atropine, and ten times greater than that of scopolamine. The toxicity is three times that of scopolamine. Because of the greater activity the dose and the relative toxicity are correspondingly less.

**SKOPYL** has been in clinical use for about ten years. Comprehensive experience has shown the preparation to be especially valuable in all conditions where spasmolysis and inhibition of secretion is desirable. Side-effects are rare and usually very slight, even when large doses are used.

**IT IS** in paediatrics that **SKOPYL** is most useful, and the value of the preparation is best seen in the treatment of hypertrophic pyloric stenosis, where it has been proved able to replace surgery in many cases. Good results have also been obtained in infantile dyspepsia, and during the spasmodic period in pertussis.

For clinical references and directions for dosage, see a special booklet that will be sent on request.

**SKOPYL** is available in tablets of 0.5 mg. and 0.1 mg. (**SKOPYL MITE**), and in 0.25% w/v solution for sublingual use.



PHARMACIA Ltd.

UPPSALA, SWEDEN

*represented by*

**SAVORY & MOORE, LTD.**

61 Welbeck Street, London, W.1

\*Trade Mark