JUVENILE POLYPS*

BY

IAN P. TODD

From 149 Harley Street, London W.1

Everyone at this meeting has seen and dealt with innumerable rectal polyps in children. It seems strange, therefore, that it is not commonly appreciated that there are several histological types of polyp in childhood. Whether the surgeon, the pathologist or perhaps semantics is to blame for this one does not know, but it does seem that 'rectal' or 'colonic polyp' is not a correct final diagnosis.

A classification of benign polyps of the large intestine has been suggested by Morson (1962b) (Table). Three are of importance in childhood: (a) the congenital cystic hamartoma or juvenile polyp, (b) Peutz-Jegher's polyps and (c) familial polyposis or adenomatosis coli. The solitary adenoma is exceedingly rare in childhood though a well-known surgical text calls the juvenile polyp an adenoma.

Familial polyposis is transmitted by either sex as a dominant gene according to Mendelian principles. The polyps, which are seldom present at birth, appear throughout childhood, and the condition should be treated before adult life is reached, probably as soon as symptoms arise, as malignant change is inevitable. Ileostomy and proctocolectomy, or colectomy with ileo-rectal anastomosis and constant supervision of the rectal stump is the correct treatment. But it is necessary to be sure that the multiple polyps are adenomata. The familial history may not be obtained. Gene mutation probably initiates the disease.

In the Peutz-Jegher's syndrome the polyps are not adenomata, but hamartomata: the malformation is of the muscularis mucosae which forms a thick tree-like structure covered with normal mucosa which includes both Paneth and Kultschitsky cells.

However, the common juvenile polyp is not an adenoma; and it is not premalignant. An adenoma is usually deep plum coloured with lobulation; on cross-section it has a tree-like central core but the muscularis mucosae is normal (Fig. 1); it is the mucosa that is abnormal, hyperchromatic and shows

some degree of epithelial hyperplasia; there are no special epithelial cells in the mucosa; and it may be sessile or pedunculated.

A juvenile polyp is cherry red, smooth, round and not lobulated (Todd, 1963); often it is surrounded by an aura of glistening yellow mucus; it may be elusive but the sight of the mucus should make one suspect its presence.

Most recent series suggest that they are slightly more common in the male (3 : 2 in my own small series of 25). Most occur in childhood or adolescence, though three cases in this series are over 30 years. Many can be felt with the finger though three cases were more than 20 cm. from the anus. The size varied from 4-20 mm.

The majority of cases exhibit a single polyp. Very rarely juvenile polyposis or hamartomatosis may occur.

Histologically these juvenile polyps are neither adenomata nor like Peutz-Jegher's polyps. The typical gross appearance has been mentioned. The cut surface, too, is characteristic (Fig. 2); there is no arborization of the muscularis mucosae, but large

TABLE
CLASSIFICATION OF BENIGN POLYPOID TUMOURS
OF THE LARGE INTESTINE

Classification	Single Polyp	Multiple Polyp
Neoplasms	Adenoma Papillary adenoma } Villous papilloma }	Familial intestinal polyposis
? Neoplasms or ? Hamartomas	Haemangioma Leiomyoma Lipoma Neurofibroma Juvenile polyp Peutz-Jegher's polyp	Lipomatosis Neurofibromatosis Juvenile polyposis Peutz-Jegher's syndrome
Hamartoma	Benign lymphoma	Lymphomatous polyposis
Inflammatory	Inflammatory (= pseudo) polyps	Following any diffuse inflammatory process, especially ulcerative colitis
Unclassified	'Hyperplastic' or 'metaplastic' polyp	Multiple lesions often seen Cystic pneumatosis

* A paper read at a meeting of the British Association of Paediatric Surgeons in Sheffield, July 1963.

FIG. 1.—An adenoma.

FIG. 2.—A juvenile polyp.

cystic spaces, filled with mucus, are easily seen. Microscopically almost the whole polyp consists of a characteristic connective tissue resembling the lamina propria of normal mucous membrane. Smooth muscle fibres are not seen and the whole polyp is covered by a layer of normal columnar mucus-secreting epithelium, one cell thick (Morson, 1962a).

Probably diathermy snare removal is the method of choice in the treatment of juvenile polyps. Incomplete removal, which is more likely to occur

with avulsion, may allow recurrence. A follow-up is unnecessary but if symptoms return a further polyp may have developed. Many are never found as auto-amputation is common, and it is the final more severe bleeding that brings mother and child to the surgery.

REFERENCES

- Morson, B. C. (1962a). Precancerous lesions of the colon and rectum. *J. Amer. med. Ass.*, **179**, 316.
 — (1962b). Some peculiarities in the histology of intestinal polyps. *Dis. Colon Rect.*, **5**, 337.
 Todd, I. P. (1963). Juvenile polyps. *Proc. roy. Soc. Med.*, **56**, 969.