


**NEW**

# STESOLID

## Diazepam-without the needle



Stesolid from Weddel  
the concept of rectal administration  
of diazepam

Stesolid is a unique system that  
obviates the need for needle in a  
variety of conditions and procedures.

Status Epilepticus - no needle

Febrile convulsions - no needle

Minor procedures of all types - no needle

There are two strengths Stesolid 5 mg and Stesolid 10 mg



### Prescribing Guidelines

**Presentation:** White, rectal tubes containing a solution of 2 mg/ml or 4 mg/ml diazepam. Approximately 2.5 ml can be squeezed from each tube giving an immediate dose of 5 mg or 10 mg diazepam. **Uses:** Diazepam is a muscle relaxant, anticonvulsant, sedative and anxiolytic. It is particularly useful in minor surgical and dental procedures or other circumstances in which a rapid effect is required but where intravenous injection is impracticable. Stesolid rectal tubes may be particularly useful for the treatment of convulsions in children and children with epilepsy. **Stesolid rectal tubes vary with age. Children:** 1-2 years of age - One 5 mg tube. Over 3 years of age - One 10 mg tube. **Adults:** One 10 mg tube. **Elderly patients -** One 5 mg tube. High doses may be required in some patients. **Contraindications:** Known sensitivity to diazepam. Acute pulmonary insufficiency. Pregnancy. **Precautions:** Stesolid rectal tubes should be stored in a cool place. **Legal Category:** POM. **Package Contents:** Packs of 5 tubes. **Batch No:** 10 mg tubes £2.25 5 x 10 mg tubes £4.25. **Product Licence Numbers:** 5 mg rectal tube 0495/0029; 10 mg rectal tube 0495/0030.

Further information is available from:




Weddel  
pharmaceuticals  
limited

Red Willow Road, Wrexham Industrial Estate, Wrexham, Cheshire LL13 9PX.

# milupa<sup>®</sup>

## means more to babies, mothers and their advisers because...



You can trust Milupa Infant Foods from the first tiny taste through to family meals.

All Milupa Foods are formulated to meet the nutritional needs of growing infants in order to promote healthy growth and development. They mix quickly

and easily in any amount to a delicious meal.

That's why Milupa means so much to babies, mothers and their advisers.

### milupa<sup>®</sup>

### The new generation of Baby Foods

Further information about the range of Milupa Infant Foods is available from Milupa Limited, Milupa House, Hercies Road, Hillingdon, Uxbridge, Middlesex UB10 9NA. Telephone: Uxbridge 59851.

# Atrovent<sup>®</sup>

ipratropium bromide inhaler

The  
difference  
speaks  
volumes

"... there is now a great body of evidence that we have a new, safe and extremely efficient anti-cholinergic drug."

Postgrad. Med. J., 1975,  
(Suppl. 7), 51, 161

"... as effective as salbutamol in increasing FEV<sub>1</sub>..."

Thorax, 1979, 34, 51

"The combination of [Atrovent and salbutamol] together more than doubled the FEV<sub>1</sub> change..."


Br. J. Dis. Chest, 1978, 72, 181

"... highly specific and virtually free of side effects even in doses well in excess of those recommended."

Thorax, 1980, 35, 137

"... more bronchodilatation than the more widely used  $\beta$ -adrenergic agonists..."

Br. Med. J., 1978, 7, 1123



**Presentation:** Metered dose inhaler containing 200 doses, each delivering ipratropium bromide 0.02mg. **Action and indications:** Anticholinergic bronchodilator for chronic reversible airways obstruction, particularly in chronic bronchitis. **Contra-indications, precautions and warnings:** Contra-indication: sensitivity to atropine. Caution in glaucoma, prostatic hypertrophy and first trimester of pregnancy. Do not spray into the eye. If a reduced response becomes apparent, the patient should seek medical advice. Anticholinergic side-effects are unlikely at therapeutic doses. **Dosage:** Adults usually 1 or 2 puffs, 3 or 4 times daily, although some patients need up to 4 puffs at a time to obtain maximum benefit during early treatment. Children 6-12 years, usually 1 or 2 puffs 3 times daily. Under 6 years usually 1 puff 3 times daily. **Pack size and basic NHS price (UK only):** 10ml vial complete with mouthpiece £4.00. Price correct at time of printing. PL 0015/0043 ▼ For full prescribing information please see data sheet.




**Boehringer  
Ingelheim**

Boehringer Ingelheim Limited,  
Bracknell, Berkshire, RG12 4YS


# A Planned Approach to Babycare

Years of experience in developing and clinically evaluating medical equipment in close co-operation with internationally known Paediatricians and Neonatologists now enables Vickers Medical to offer a complete range of equipment to assist the recovery process of the sick neonate.


Each unit fulfils individual clinical needs by providing infants at risk with better chances of survival and can be integrated to achieve even higher standards of infant critical care. The design philosophy of Vickers Medical embraces adaptability and flexibility to enable specific requirements to be fulfilled now and at the same time plan for future therapeutic needs—in short "A planned Approach to Babycare".



**The Resuscitaire 165**  
A comprehensive resuscitation system to provide all the facilities to help babies born with breathing difficulties survive the immediate post-natal period.



**The Transport Incubator System** For the safe movement of pre-term and critically ill neonates between hospital departments and for longer journeys by ambulance.



**The Neovent** A continuous-flow, pressure-limited, time-cycled ventilator for the respiratory support of critically ill neonates and infants up to 3 years old.

**The Neocare**  
A safe and accurately controlled environment for nursing critically ill babies. Now available to maintain either the baby's skin temperature or the air temperature at a constant pre-selected level.

...the treatment of jaundice.

Further information from....  
Vickers Medical, Priestley Road,  
Basingstoke, Hants, England, RG24 9NP.  
Telephone: (0256) 29141 Telex: 858697



---

## MILUPA STUDENT ELECTIVE GRANT FUND

### *Student elective grants for paediatric nutrition projects*

Milupa Limited are offering, during 1983, a limited number of Grants, of up to £500 each, to enable Medical Students to undertake a specific research project related to paediatric nutrition during their Elective period.

Applications should be supported with full details of the proposed project, a recommendation from the Head of the Department at the University Medical School, and documented agreement that the applicant will be received in the host department, plus a detailed C.V. and summary of expenses.

Grants will be made on a competitive basis and will be awarded at the end of June 1983. The closing date for applications will be 16 May 1983.

Applications for elective periods to be undertaken before 30 June 1983 cannot be considered. The selection of applicants for awards will be made by an independent scientific panel. Applications should be sent to Mr. T. L. Bell, Managing Director, Milupa Limited, Milupa House, Hercies Road, Hillingdon, Uxbridge, Middlesex UB10 9NA.

## MILUPA TRAVEL GRANT FUND

### *Travel grants for paediatric research*


Milupa Limited are prepared to award Travel Grants to young Paediatricians and Scientists working in a Paediatric Department for visits of short duration to Departments or Institutes in order to learn new techniques or to have discussions about specific research projects. Preference will be given to candidates wishing to study in the U.K., but Grants may be made for travel abroad if the particular technique is not available in the U.K.

Applications giving full details of the proposed programme and the names of two referees may be submitted at any time.

All applications should be sent to Mr. T. L. Bell, Managing Director, Milupa Limited, Milupa House, Hercies Road, Hillingdon, Uxbridge, Middlesex UB10 9NA, from whom further particulars may also be obtained. The selection of applicants for awards will be made by an independent scientific panel.

---

# This Publication is available in Microform.



## University Microfilms International

Please send additional information

for \_\_\_\_\_  
(name of publication)

Name \_\_\_\_\_

Institution \_\_\_\_\_


Street \_\_\_\_\_

City \_\_\_\_\_

State \_\_\_\_\_ Zip \_\_\_\_\_

300 North Zeeb Road  
Dept. P.R.  
Ann Arbor, Mi. 48106





When  
efficacy  
is the need  
and safety  
the concern

No need now to sacrifice efficacy for safety. Eumovate fulfils the need for a corticosteroid preparation with a high topical activity than hydrocortisone, yet with a wide margin of safety.



# Eumovate

(clobetasone butyrate)

a unique balance of efficacy and safety

#### Prescribing information

##### Uses

Eumovate is suitable for treating the milder forms of eczema, seborrhoeic dermatitis and other steroid responsive skin conditions.

##### Dosage and administration

Apply up to four times a day until improvement occurs, when the frequency of application may be reduced.

##### Contra-indications

Bacterial, fungal or viral diseases of the skin.  
Hypersensitivity to the preparation.

##### Precautions

In infants and children, long-term continuous topical corticosteroid therapy should be avoided where possible, as adrenal suppression can occur. In infants, the napkin may act as an occlusive dressing, and increase absorption.

Appropriate antimicrobial therapy should be used concurrently whenever treating inflammatory lesions which may become infected.

Any spread of infection requires withdrawal of topical corticosteroid therapy and, possibly, commencement of systemic chemotherapy.

Avoidance of occlusives, prolonged application to the face or sensitive sites.

In general, topical steroids should not be used extensively in pregnancy, i.e., in large amounts or for prolonged periods.

##### Side effects

In the unlikely event of signs of hypersensitivity appearing, application should stop immediately. If large areas were to be treated, some patients could absorb sufficient steroid to cause transient adrenal depression despite the low degree of systemic activity associated with clobetasone butyrate. Local atrophic changes could occur where moisture increases absorption of steroid, but only after prolonged use.

Basic NHS cost (exclusive of VAT)  
25 gram tube £1.63 (also available in 100 gram tubes)

##### Product licence numbers

Cream: 4/0233  
Ointment: 4/0254

**Glaxo**

leaders in topical steroid therapy

Further information on Eumovate (trade mark) is available from:

Glaxo Laboratories Limited  
Greenford, Middlesex UB6 2HE