officer with day to day responsibility for the service to have complete autonomy and consequent power to protect/develop the service, working from the present health authority base. A senior clinical medical officer in child health feels a more natural affinity for the paediatrician with whom he works as a clinical colleague. The BPA must really take the initiative and act swiftly to promote an integrated child health service. Community medicine should be a planning resource only and the faculty bid to stay in control must be firmly resisted.

A SENIOR CLINICAL MEDICAL OFFICER

Juvenile discitis

Sir,

Hensley *et al*¹ seemed unconvinced of the value of antibiotic treatment in their six cases of juvenile discitis. We have seen and treated four children (three under the age of 5 years) in the past four years with clinical, radiological, and laboratory features consistent with the condition as described by the authors (Table). None of our patients were systemically ill, although one, our first, did present with mild glomerulonephritis and in his case a throat swab yielded beta haemolytic streptococci Lancefield group A. All four patients responded to antibiotic treatment. Penicillin was stopped after 10 days in case 1 with recurrence of backache and inability to walk within a week. His symptoms resolved again within a few days of restarting penicillin and flucloxacillin and he remained well after being given treatment for a further 6 weeks. The other three children received antibiotic treatment for 6 weeks by which time the erythrocyte sedimentation rate had returned to normal in each case.

After two years follow up one of our patients had a recurrence of symptoms, but she again responded to antibiotic treatment with ampicillin and flucloxacillin and has since remained asymptomatic for over two years. We feel that although it is possible that some cases of juvenile discitis may be related to trauma,² infection is the common cause and broad spectrum antibiotic treatment should be offered to all cases. Immobilisation of the spine for a prolonged period has been recommended by some authors^{3,4} but we did not find this necessary in any of our patients.

M P SHAH AND C G MILLER
District General Hospital,
Lovely Lane,
Warrington WA5 1QG

Dr Hensley and co-workers comment:

Drs Shah and Miller confirm our opinion that the treatment and origin of juvenile discitis are controversial.¹ In the patients we described we failed to find any haematological or bacteriological support for an infective

Table Clinical and laboratory findings in four patients with juvenile discitis

Case no	Sex	Age at presentation (years)	Date of presentation	Presenting symptoms	Disc space affected	Bacteriology	ESR	Treatment	Outcome
1	M	4	SEPT 1979	Backache for 1 month, reluctance to sit, stand, or walk for 2 days	L4/L5	T/S Group A beta haemolytic streptococci	39	Oral penicillin for 10 days, then oral penicillin and flucloxacillin for 6 weeks	Initial relief within a week. Relapsed after stopping treatment. Prompt relief on restarting antibiotics.
2	F	13/12	DEC 1979	Backache for 3 weeks, stiff gait	L1/L2	Streptococcus viridans 1/6 blood culture bottles	50	Amoxicillin and flucloxacillin for 6 weeks on two occasions	Gradual recovery. Asymptomatic within a month. Relapsed 2 years later but again responded.
3	M	11/12	OCT 1980	Irritability, reluctance to sit, refusal to stand	L5/S1	Negative	41	Penicillin and flucloxacillin for 6 weeks	Asymptomatic after 2 weeks treatment. Well since.
4	M	2/12	AUG 1981	Limp for 8 days	L2/L3	Negative	56	Amoxicillin and flucloxacillin for 6 weeks	Resolution of symptoms within a week. Well since.