

## Phototherapy: an ocular hazard revisited

Phototherapy is an established and effective treatment for neonatal jaundice.<sup>1</sup> There remains concern, however, that this necessary light exposure may pose a potentially noxious stimulus to the eyes of neonates.<sup>2–4</sup> Recent studies have suggested that, irrespective of phototherapy, prolonged exposure to 'bright' ambient light may damage cone photoreceptors<sup>3</sup> and increase the incidence of retinopathy of prematurity.<sup>5</sup> In view of these findings it is perhaps timely to reconsider the potential ocular hazard posed by phototherapy and the practices required to minimise the likelihood of damage occurring.

### The efficacy of 'patching'

The eyes of phototreated infants are routinely 'patched', because it has been shown that the intensity of light in phototherapy units can cause retinal damage in animals.<sup>2</sup> In a study of 12 different eyeshields obtained from neonatal units within the United Kingdom, Chin *et al*<sup>6</sup> found that all shields tested were extremely effective in reducing the intensity of incident light. They and others,<sup>7</sup> however, noted that patches are prone to slip. Such occurrences are explicable on two counts: it is often difficult to secure eyeshields effectively and, in addition, nursing staff often think of 'patching' as of secondary importance to more immediate and potentially life preserving interventions.

To what extent does inadequate patching—or indeed no patching at all—increase the ocular hazard posed by phototherapy? In fact there is little evidence that phototherapy has caused retinal damage in human infants. This may be due partly to limited follow up testing, and partly to the fact that subtle damage may go undetected. Most studies, however, have reported carefully 'patched' infants who may not be expected to suffer damage.<sup>8</sup>

### Animal experiments

The case for routine patching of infants' eyes during phototherapy is mostly based on extrapolations from animal studies. Here an important distinction must be made between the substantial number of reports about ocular phototoxicity<sup>9–10</sup> and the relatively few studies that have, in an attempt to mimic clinical practice, used an appropriate (that is, neonatal) animal with light exposure obtained from

a phototherapy unit. Of the latter more relevant studies, Sisson *et al*<sup>11</sup> exposed newborn piglets (36 hours old) to phototherapy for 72 hours. In contrast to human neonates, the piglets' pupils were dilated to compensate for their orientation at 90° to the light source. At necropsy pronounced histological damage to rod and cone photoreceptors was found, and this was attributed to the light exposure. In a similar study, Messner *et al*<sup>12</sup> found retinal damage in the eyes of newborn monkeys that were exposed to phototherapy for up to seven days; in this experiment, however, the animals were restrained so that they directly faced the light source.

Neither of these studies accurately represents the conditions prevailing in clinical practice. Firstly, human neonates undergoing phototherapy are never purposely restrained to face the light source, nor will their pupils be kept dilated during treatment. Secondly, the amount of time the experimental animals spent with their eyes open was not addressed; studies in humans have shown that premature neonates may spend more than 80% of the time with their eyes fully closed.<sup>13–14</sup> If the transmission of light through the closed eyelids is similar to that of the adult, there will be a wavelength dependent reduction in the intensity of light reaching the cornea of between 10 to 100 times.<sup>15</sup>

### The developing human eye

Consideration of the developing human eye emphasises that further caution is warranted when extrapolating from animal studies. For example, corneal haze<sup>16</sup> and the transient tunica vasculosa lentis (present in the eye of the premature neonate until about 34 weeks' gestation) may reduce the intensity of light reaching the retina. Though photoreceptors in the peripheral retina of the human neonate are relatively well developed at full term, the macular region remains poorly differentiated.<sup>17–18</sup> Similarly, retinal vascular development is not complete until just after term.<sup>19</sup> The relevance of these observations to ocular phototoxicity in human neonates has yet to be addressed but they highlight the difficulties of extrapolating from animal experiments.

### Conclusions

We conclude that further research is required to evaluate the ocular hazard posed by phototherapy.

Firstly, future animal studies should be designed in such a manner as to ensure that the conditions prevalent in clinical practice are appropriately simulated. Secondly, estimates of typical exposures of the retina received by infants during phototherapy are currently unavailable and would be of great benefit in evaluating the potential hazard. To this end we are attempting to measure both the environmental (for example, the intensity and duration of light exposure) and biological (for example, eyelid opening, pupil size, transmittance of the eyelids, and optic media) factors that influence retinal exposure during phototherapy. It is hoped that such data can be related to known photobiological dose response associations obtained from in vivo and in vitro laboratory studies.

Should phototherapy be found to pose an important ocular hazard then the use of eyeshields needs to be afforded the appropriate clinical priority. If it is not, then the economics of 'patching' and its unknown effects on visual development may not justify the continuation of the practice.

Dr Moseley is supported by the Wellcome Trust.

#### References

- <sup>1</sup> Cashore WJ, Stern L. The management of hyperbilirubinemia. In: Zipursky A, ed. *Clinics in perinatology*. Philadelphia: WB Saunders, 1984.
- <sup>2</sup> Dobson V. Phototherapy and retinal damage. *Invest Ophthalmol* 1976;**15**:595-8.
- <sup>3</sup> Abramov I, Hainline L, Lemerise E, Brown AK. Changes in visual functions of children exposed as infants to prolonged illumination. *J Am Optom Assoc* 1985;**56**:614-9.
- <sup>4</sup> Wolke D. Environmental neonatology. *Arch Dis Child* 1987;**62**:987-8.
- <sup>5</sup> Glass P, Avery GB, Siva Subramanian KN, Keys MP, Sostek AM, Friendly DS. Effect of bright light in the hospital nursery on the incidence of retinopathy of prematurity. *N Engl J Med* 1985;**313**:401-4.
- <sup>6</sup> Chin KC, Moseley MJ, Bayliss SC. Light transmission of phototherapy eyeshields. *Arch Dis Child* 1987;**62**:970-1.
- <sup>7</sup> Patz A, Souri EN. Phototherapy and other ocular risks to the newborn. *The Sight-Saving Review* 1972;**42**:29-33.
- <sup>8</sup> Moseley MJ, Fielder AR. Light toxicity and the neonatal eye. *Clinical Vision Sciences* 1988 (in press).
- <sup>9</sup> Lanum J. The damaging effects of light on the retina. empirical findings, theoretical and practical implications. *Surv Ophthalmol* 1978;**22**:221-49.
- <sup>10</sup> Lerman S. Ocular phototoxicity. In: Davidson SI, Fraunfelder FT, eds. *Recent advances in ophthalmology*. Edinburgh: Churchill Livingstone, 1985.
- <sup>11</sup> Sisson TRC, Glauser SC, Glauser EM, Tasman W, Kuwabara T. Retinal changes produced by phototherapy. *J Pediatr* 1970;**77**:221-7.
- <sup>12</sup> Messner KH, Maisels MJ, Leure-DuPree AE. Phototoxicity to the newborn primate retina. *Invest Ophthalmol Vis Sci* 1978;**17**:178-82.
- <sup>13</sup> Berman J, Hutchins R, Gold S, Peli E, Lindsey P. Measurement of light exposure in infants at high risk for developing retinopathy of prematurity. *Invest Ophthalmol Vis Sci* 1987;**28** (suppl):119.
- <sup>14</sup> Moseley MJ, Thompson JR, Levene MI, Fielder AR. Effects of nursery illumination on frequency of eyelid opening and state in preterm neonates. *Early Hum Dev* 1988 (in press).
- <sup>15</sup> Moseley MJ, Fielder AR. Light transmission through the human eyelid: in vivo measurement. *Ophthalmic Physiol Opt* 1988 (in press).
- <sup>16</sup> Hittner HM, Hirsch NJ, Rudolph AJ. Assessment of gestational age by examination of the anterior vascular capsule of the lens. *J Pediatr* 1977;**91**:455-8.
- <sup>17</sup> Abramov I, Gordon J, Hendrickson A, Hainline L, Dobson V, LaBossiere E. The retina of the newborn human infant. *Science* 1982;**217**:265-7.
- <sup>18</sup> Yuodelis C, Hendrickson A. A qualitative and quantitative analysis of the human fovea during development. *Vision Res* 1986;**26**:847-55.
- <sup>19</sup> Ashton N. Retinal angiogenesis in the human embryo. *Br Med Bull* 1970;**26**:103-6.

M J MOSELEY and A R FIELDER  
 Department of Ophthalmology,  
 University of Leicester,  
 Clinical Sciences Building,  
 PO Box 65, Leicester Royal Infirmary,  
 Leicester LE2 7LX