be a problem only in that group of patients, of whom patient 1 is a typical example, who experience recurrent episodes of rejection.

Heart-lung transplantation is acceptable treatment for children with terminal respiratory disease. Although information on long term survival is not yet available, the improved quality of life in the short term in the children we have treated is encouraging.

Dr Smyth is supported by the Cystic Fibrosis Research Trust. We thank the staff of donor hospitals for their helpful cooperation and Mr Ben Milstein for his editorial comments and criticisms.

References

- ¹ Reitz B A, Wallwork J, Hunt S A, *et al*. Heart-lung transplantation. Successful therapy for patients with pulmonary vascular disease. *N Engl J Med* 1982;**306**:557-64.
- ² Penketh A, Higenbottam T, Hakim M, Wallwork J. Heart and lung transplantation in patients with end-stage lung disease. *Br Med J* 1987;**295**:311-4.
- ³ Scott J P, Higenbottam T W, Penketh A R L, Hodson M, Stewart S, Wallwork J. Heart-lung transplantation for cystic fibrosis. *Lancet* 1988;ii:192-4.
- ⁴ Hutter J A, Despins P, Higenbottam T W, Stewart S, Wallwork J. Heart-lung transplantation: better use of resources. *Am J Med* 1988;**85**:4-11.
- ⁵ McGavin C R, Artinintini M, Nave H, McHardy G J R. Dyspnoea disability and distance walked: comparison of estimates of exercise performance in respiratory disease. *Br Med J* 1978;ii:241-4.
- ⁶ Hakim M, Higenbottam T W, Bethune D, *et al*. Selection and procurement of combined heart and lung grafts for transplantation. *J Thorac Cardiovasc Surg* 1988;**95**:474-9.
- ⁷ Jamieson S W, Baldwin J, Stinson E B, *et al*. Clinical heart-lung transplantation. *Transplantation* 1984;**37**:81-4.
- ⁸ Jamieson S W, Stinson E B, Oyer P E, Baldwin J C, Shumway N C. Operative technique for heart-lung transplantation. *J Thorac Cardiovasc Surg* 1984;**87**:930-5.
- ⁹ Wallwork J, Jones K, Cavarocchi N, Hakim M, Higenbottam T W. Distant procurement of organs for clinical heart-lung transplantation using a single flush technique. *Transplantation* 1987;**44**:654-8.
- ¹⁰ Higenbottam T W, Stewart S, Penketh A, Wallwork J. The diagnosis of lung rejection and opportunistic infection by transbronchial lung biopsy. *Transplant Proc* 1987;**19**:3777-8.
- ¹¹ Stewart S, Higenbottam T W, Hutter J A, Penketh A R L, Zebro T J, Wallwork J. Histopathology of transbronchial biopsies in heart-lung transplantation. *Transplant Proc* 1988;**20**:764-6.
- ¹² Higenbottam T W, Stewart S, Penketh A R L, Wallwork J. Transbronchial lung biopsy for the diagnosis of rejection in heart-lung transplant patients. *Transplantation* 1988;**46**:532-40.
- ¹³ Higenbottam T W, Hutter J A, Stewart S, Wallwork J. Transbronchial biopsy has eliminated the need for endomyocardial biopsy in heart-lung recipients. *Journal of Heart Transplantation* 1988;**7**:435-9.
- ¹⁴ Cotes J E. Lung function throughout life; determinants and reference values. *Lung function*. 4th ed. London: Blackwell, 1979:340.
- ¹⁵ Fricker F J, Griffith B P, Hardesty R L, *et al*. Experience with heart transplantation in children. *Pediatrics* 1987;**79**:138-46.
- ¹⁶ Addonizio L J, Rose E A. Cardiac transplantation in children and adolescents. *J Pediatr* 1987;**111**:1034-8.
- ¹⁷ Starnes V A, Stinson E B, Oyer P E, *et al*. Cardiac transplantation in children and adolescents. *Circulation* 1987;**76**:(suppl V) V.43-7.
- ¹⁸ Fuster V, Steele P M, Edwards W D, Gersh B J, McGoon M D, Frye R L. Primary pulmonary hypertension: a natural history and the importance of thrombosis. *Circulation* 1984;**70**:580-7.
- ¹⁹ Siassi B, Moss A J, Dooley R R. Clinical recognition of cor pulmonale in cystic fibrosis. *J Pediatr* 1971;**78**:794-805.
- ²⁰ Bishop J M, Cross K W. Physiological variables and mortality in patients with various categories of chronic respiratory disease. *Bull Eur Physiopath Respir* 1984;**20**:495-500.
- ²¹ Wallwork J, Williams R, Calne R Y. Transplantation of liver heart and lungs for primary biliary cirrhosis and primary pulmonary hypertension. *Lancet* 1987;ii:182-5.
- ²² Otulana B A, Higenbottam T W, Hutter J, Wallwork J. Close monitoring of lung function in heart-lung transplants allows detection of pulmonary rejection and infection. *Am Rev Respir Dis* 1988;**137**:245.
- ²³ Gray J J, Alvey B, Smith D J, Wreghitt T G. Evaluation of a commercial latex agglutination test for detecting antibodies to cytomegalovirus in organ donors and transplant recipients. *J Virol Methods* 1987;**16**:13-9.
- ²⁴ Hutter J, Scott J P, Wreghitt T G, Higenbottam T, Wallwork J. The importance of cytomegalovirus in heart-lung transplant recipients. *Chest* 1989;**95**:627-31.
- ²⁵ Brooks R G, Hofflin J M, Jamieson S W, Stinson E B, Remington J S. Infectious complications in heart-lung transplant recipients. *Am J Med* 1985;**79**:412-22.
- ²⁶ Scott J P, Higenbottam T W, Clelland C, Hutter J, Wallwork J. The natural history of obliterative bronchiolitis in patients with heart-lung transplants. *Transplant Proc* 1989;**21**:2592-3.
- ²⁷ Scott J P, Higenbottam T W. Adverse reactions and interactions of cyclosporin. *Medical Toxicology* 1988;**3**:107-27.

Correspondence to Dr TW Higenbottam, Papworth Hospital, Papworth Everard, Cambridge CB3 8RE.

Accepted 14 March 1989

Commentary

J O WARNER

National Heart and Lung Institute, Brompton Hospital, London

The development of heart-lung transplantation as a new treatment for severe irreversible respiratory failure has given new hope to a large number of patients and their families facing the prospect of premature death. The extension of this procedure to the paediatric age range was inevitable and it is reassuring that the results that have been obtained from the first few cases are at least as good as those in adults. Any paediatrician who has the experience of seeing a child in terminal respiratory failure, miraculously given new life after a transplant, can only be immensely optimistic and positive about this procedure. However, it is most important that it should be put into perspective with a clear understanding of what is involved.

A practical treatment it may be for children in terminal respiratory failure. However, only a few individuals in this predicament will benefit from transplantation. From the Papworth experience so far, only 15% of referred patients have benefited.

The question that must be asked is whether the anguish of the remaining 85% is sufficiently offset by the successful outcome in the minority.

The Brompton Hospital paediatric respiratory department has assessed 18 children for transplant over the last two and half years. Only two have, so far, been transplanted. One remains well two years later and the other, after 15 months, has had a number of complications, including cytomegalovirus pneumonitis, which is currently controlled on ganciclovir and several rejection episodes culminating in the development of severe obstructive airways disease. Five children were accepted on the high priority waiting list for transplantation. Two of these patients have died and the other three are still alive. Eight referred patients were considered unsuitable for transplantation: five for medical reasons; two who were not considered sufficiently ill to warrant transplantation; and one with major psychosocial problems who was an extremely poor complier with treatment. Amazingly the parents of three children referred for transplant, on further reflection, did not wish their children to be included in the programme. This arose because of inadequate discussion with the family before referral. Thus my experience on smaller numbers is very similar to that at Papworth Unit, but there are additional factors to consider.

The emotional stress created by referral is enormous. Frequently it has involved travel over long distances, splitting families and creating a significant financial burden. The anguish of patients and families while waiting for transplants, in a situation when health is deteriorating rapidly, is considerable. Attitudes to impending death are funda-

mentally changed. Children in terminal respiratory failure must always be given appropriate treatment so that they die peacefully, with dignity, and without pain. The prospect of a transplant has sadly led to delays in institution of such treatment. This is wholly unacceptable, but the patient and parents' grim determination to struggle on as long as possible is perfectly understandable once the new hope has been generated. One of my patients who died on the waiting list, in fact indicated, a few hours before death, that she had not wanted to have a transplant. She had given her parents extra hope by acceding to their wishes. In the two weeks before death she had deliberately stopped all her treatment and hidden it in a paper bag. A psychiatric assessment of the patient and family has, therefore, now become an essential part of the selection procedure.

Thus, in the short term, paediatricians must be aware that at best only 20% of referrals will receive a transplant, three quarters of whom can be expected to be alive after one year. At least 20% will be considered unsuitable for medical reasons, and of the remainder, half will die on the waiting list and the other half will still be waiting, perhaps a year after original referral.

At a time of medical rationing, with restriction of resources, can this procedure be justified? I believe the answer is yes, but extreme care is required before even considering referral to a centre for assessment. Only the most psychologically robust individuals, fulfilling appropriate criteria for severe respiratory failure, and without any other complicating medical problems, can be accepted on transplant programmes.