

- their characteristics in digestion of dietary fat. *J Pediatr Gastroenterol Nutr* 1982;1:243-55.
- 4 Spendlove RS, Schaffer FL. Enzymatic enhancement of infectivity of reovirus. *J Bacteriol* 1965;89:597-602.
- 5 Barnett BB, Spendlove RS, Clark ML. Effect of enzymes on rotavirus infectivity. *J Clin Microbiol* 1979;10:111-3.

Dr Isaacs comments:

I certainly agree with Professor Dodge and Dr Sagher that milk lipids may provide important protective effects. Further, I find their observation that alterations in the medium chain fatty acid composition of mouse milk may reduce the severity of infection from rotavirus, a non-enveloped virus, extremely interesting. We, of course, did not determine the effect of the stomach contents of infants fed the various formulas on non-enveloped viruses. Rather, our statement that milk fatty acids and monoglycerides have antiviral activity against enveloped viruses but not non-enveloped viruses was based upon the results of a number of studies by us and others showing that only enveloped viruses are inactivated by milk lipids or purified lipids.

In the early studies of Welsh *et al* enveloped Semliki Forest virus and herpes simplex virus type 1 (HSV-1) were inactivated by milk lipids but enterovirus coxsackie B4, a non-enveloped virus, was not.¹ In addition, we found that human milk inactivated the enveloped measles virus, vesicular stomatitis virus and HSV-1 but not the non-enveloped vaccinia virus and poliovirus.² Milk lipids also have been shown to inactivate dengue virus³ and mouse mammary tumour virus⁴; these are both enveloped viruses.

In *in vitro* studies it has been found that purified free fatty acids and their derivatives inactivated the enveloped Sendai virus, Newcastle disease virus, influenza A virus, Sindbis virus, West Nile virus, HSV-1, and a number of enveloped bacteriophages but not the non-enveloped SV40, polio or encephalomyocarditis viruses (ECMV).^{5,6} This antiviral effect appeared to be due to the destruction of viral envelopes.⁷

These multiple findings suggest that the apparent protective effect of milk lipids against rotavirus infection observed by Dodge and Sagher is unlikely to be due to direct inactivation of the non-enveloped rotavirus. However, milk fatty acids may prevent the binding of rotavirus and other non-enveloped viruses to receptors or interfere with viral uncoating. These suggestions are supported by the observations of JFE Newman (Institute for Virology, Sandringham, South Africa), reported at the recent VIIIth International Congress of Virology (Berlin, 1990), that fatty acids with chain lengths of 12-15 prevent uncoating of some non-enveloped viruses, for example, bovine enterovirus and ECMV, but not others, for example, poliovirus type 1, coxsackie B5, and human rhinoviruses 1B and 14. It would, therefore, be interesting to see the results of an *in vitro* study examining the effects of medium chain fatty acids found in mouse milk on rotavirus infectivity.

- 1 Welsh JK, Arsenakis M, Coelen RJ, May JT. Effect of antiviral lipids, heat, and freezing on the activity of viruses in human milk. *J Infect Dis* 1979;140:322-8.
- 2 Isaacs CE, Thormar H, Pessolano T. Membrane-disruptive effect of human milk: inactivation of enveloped viruses. *J Infect Dis* 1986;154:966-71.

- 3 Falkler WA Jr, Diwan AR, Halstead SB. A lipid inhibitor of dengue virus in human colostrum and milk; with a note on the absence of anti-Dengue secretory antibody. *Arch Virol* 1975;47:3-10.
- 4 Sarkar NH, Charney J, Dion AS, Moore DH. Effect of human milk on the mouse mammary tumor virus. *Cancer Res* 1973;33:626-9.
- 5 Kohn A, Gitelman J, Inbar M. Unsaturated free fatty acids inactivate animal enveloped viruses. *Arch Virol* 1980;66:301-7.
- 6 Sands JA, Auperin DD, Landin PD, Reinhardt A, Cadden SP. Antiviral effects of fatty acids and derivatives: lipid-containing bacteriophages as a model system. In: Kabara JJ, ed. *Symposium on the pharmacological effect of lipids*. Champaign, Illinois: The American Oil Chemists Society, 1978:75-95.
- 7 Thormar H, Isaacs CE, Brown HR, Barshatzky MR, Pessolano T. Inactivation of enveloped viruses and killing of cells by fatty acids and monoglycerides. *Antimicrob Agents Chemother* 1987;31:27-31.

Fluorescein dilaurate test of exocrine pancreatic function in cystic fibrosis

SIR,—I read with interest the paper by Drs Dalzell and Heaf,¹ particularly as we had a very similar study published in the *Archives* four years previously (to which no reference was made!).² In both studies the index cases and controls were similar in age and number, but we prescribed double the dose of fluorescein dilaurate than did Dalzell and Heaf. Our study also demonstrated significantly different fluorescein dilaurate excretion ratios between patients with cystic fibrosis and normal subjects, with the ratios being significantly reduced in patients with cystic fibrosis ($p < 0.01$). An additional component to our study was to compare the fluorescein dilaurate test with faecal chymotrypsin estimation. We found a positive correlation between the two tests ($R = 0.69$, $p < 0.02$).

Although the fluorescein dilaurate test appears to detect exocrine pancreatic insufficiency, in practice it is of limited value as it is in capsule form and not suitable for the age group in which the presentation of cystic fibrosis is most prevalent. We did explore the possibility of the test being used to titrate pancreatic supplement administration. If cholesterol ester hydrolase, which is responsible for liberating the fluorescein from fluorescein dilaurate, was incorporated into a pancreatic enzyme supplement, it may be possible to use the fluorescein dilaurate test to determine the most effective dosage of pancreatic enzyme for individual patients. Unfortunately this enzyme does not appear to be present in any of the commercially available preparations.

J S FORSYTH
Dundee General Hospitals Unit,
Ninewells Hospital and Medical School,
Ninewells,
Dundee DD1 9SY

- 1 Dalzell AM, Heaf DP. Fluorescein dilaurate test of exocrine pancreatic function in cystic fibrosis. *Arch Dis Child* 1990;65:788-9.
- 2 Cumming JGR, Forsyth JS, Boyd EJS, *et al*. Diagnosis of exocrine pancreatic insufficiency in cystic fibrosis by use of fluorescein dilaurate test. *Arch Dis Child* 1986;61:573-5.

Hospital admission—a missed opportunity to immunise

SIR,—We agree with the conclusion of Ferson's study that there is poor documentation of a child's immunisation status on admission to hospital.¹ In 66% of the children admitted to our hospital and verified to have incomplete immunisation the 'medical record of immun-

isation' was absent or in so shortened a form as to be meaningless.

Using information from the district computer we determined the immunisation status of children from Salford district who were admitted to a general paediatric ward in a teaching hospital during November 1989. We also obtained details about the children verified to have incomplete immunisation, including the recorded immunisation history, from the hospital notes.

There were 139 admissions of 133 Salford children during the month. The children's ages ranged from 7 days to 15 years with 113 under 5 years of age. Using the immunisation schedules recommended at that time, information from the district computer showed that 70 children were fully immunised and 47 incompletely immunised; 16 were not traced by the computer.

Information from the hospital notes suggested that in five children there was a contra-indication to immunisation and three children had received the necessary immunisation within the two weeks before admission. The hospital notes for six children were not available. The remaining 33 children could have been offered: triple antigen ($n = 14$), oral polio ($n = 14$), combined diphtheria-tetanus ($n = 1$), measles-mumps-rubella ($n = 22$), tetanus ($n = 3$), and pertussis immunisations ($n = 12$).

Using Ferson's classification of the 'medical record of immunisation' in the hospital notes,¹ in 27 of the 41 notes (66%) documentation was absent or unclear. In two (5%) it was clear but incorrect, and documentation was correct and clear in only 12 (29%) of notes.

At our children's hospital, sited in a district with poor immunisation uptake, the opportunities for immunisation in a single month on one ward are considerable. To immunise in hospital we need accurate information on previous immunisations. Parental recall and medical recording of immunisation provide inadequate data. Two possible solutions are improving access to immunisation information held on district computers and using parent held child health records.

I A MCKINLAY
W J RANKIN
Agnew Unit,
Royal Manchester Children's Hospital,
Pendlebury, Manchester M27 1HA

- 1 Ferson MJ. Immunisation state and its documentation in hospital patients. *Arch Dis Child* 1990;65:763-7.

Effects of overweight on lung function

SIR,—We were very interested in the paper by Dr Fung and colleagues on overweight and lung function,¹ but we have some queries about their approach and conclusion.

To investigate the association between body mass index and lung function in children, the authors included subjects up to 20 years of age. In table 1 they show that distribution of height for the sexes was unequal: 45% of all girls were taller than 150 cm but only 12% were taller than 159 cm. This suggests that most of these females had reached their adult height: at age 13, when 75% of Hong Kong girls reach menarche, the median height is 151 cm.² On the other hand, 31% of all boys were taller than 159 cm with no skewed distribution, suggesting that most boys had not reached adult height yet.