

Nationwide study of haemolytic uraemic syndrome: clinical, microbiological, and epidemiological features

Department of Paediatrics and Child Health, University of Sydney and The Children's Hospital at Westmead, Sydney, Australia
E J Elliott

Department of Gastroenterology, The Children's Hospital at Westmead
E V O'Loughlin

Centre for Kidney Research, The Children's Hospital at Westmead
J Knight

Microbiological Research Unit, Department of Microbiology and Infectious Diseases, Royal Children's Hospital, Melbourne
R M Robins-Browne
V Bennett-Wood

Department of Nephrology, Royal Children's Hospital, Melbourne
H Powell

Royal Children's Hospital, Mater Hospital and Princess Alexandra Hospital, Brisbane, Queensland, Australia
J Bourke

Women's and Children's Hospital, Adelaide
P Henning

Microbiological Diagnostic Unit, Department of Microbiology and Immunology, University of Melbourne
G G Hogg

Australian Paediatric Surveillance Unit
D Redmond

Correspondence to:
A/Prof E Elliott, Dept of Paediatrics and Child Health University of Sydney, c/o The Children's Hospital at Westmead, Locked Bag 4001, Westmead, NSW, 2145, Australia
elizabe2@chw.edu.au

Accepted 28 March 2001

E J Elliott, R M Robins-Browne, E V O'Loughlin, V Bennett-Wood, J Bourke, P Henning, G G Hogg, J Knight, H Powell, D Redmond, Contributors to the Australian Paediatric Surveillance Unit

Abstract

Aims—To establish the incidence and aetiology of haemolytic uraemic syndrome (HUS) in Australia and compare clinical and microbial characteristics of sporadic and outbreak cases.

Methods—National active surveillance through the Australian Paediatric Surveillance Unit with monthly case notification from paediatricians, July 1994 to June 1998. Children under 15 years presenting with microangiopathic haemolytic anaemia, thrombocytopenia, and acute renal impairment were identified.

Results—Ninety eight cases were identified (incidence 0.64 per 10⁵ children <15 years/annum and 1.35 per 10⁵ children <5 years/annum). Eighty four were associated with diarrhoea (64 sporadic, 20 constituting an outbreak) and 14 were atypical. Shiga toxin producing *Escherichia coli* (STEC) O111:H- was the most common isolate in sporadic HUS and caused the outbreak. However O111:H- isolates from outbreak and sporadic cases differed in phage type and subtyping by DNA electrophoresis. STEC isolates from sporadic cases included O26:H-, O113:H21, O130:H11, OR:H9, O157:H-, ONT:H7, and ONT:H-. STEC O157:H7 was not isolated from any case. Only O111:H- isolates produced both Shiga toxins 1 and 2 and possessed genes encoding *E coli* attaching and effacing gene (intimin) and enterohemolysin. Outbreak cases had worse gastrointestinal and renal disease at presentation and more extrarenal complications.

Conclusions—Linking national surveillance with a specialised laboratory service allowed estimation of HUS incidence and provided information on its aetiology. In contrast to North America, Japan, and the British Isles, STEC O157:H7 is rare in Australia; however, non-O157:H7 STEC cause severe disease including outbreaks. Disease severity in outbreak cases may relate to yet unidentified virulence factors of the O111:H- strain isolated.

(Arch Dis Child 2001;85:125-131)

Keywords: Shiga toxin; E coli; haemolytic uraemic syndrome; surveillance

Haemolytic uraemic syndrome (HUS) is characterised by microangiopathic haemolytic anaemia, thrombocytopenia, and acute renal

impairment. It is usually preceded by diarrhoea, often bloody. Karmali *et al* first identified the association between HUS and infection with Shiga toxin producing *Escherichia coli* (STEC) O157:H7 in children,^{1 2} which is now well established.³⁻⁷ STEC were previously known as verotoxin producing *E coli*. In North America and Europe most HUS is caused by STEC of serotype O157:H7 and major outbreaks have occurred following ingestion of contaminated food or drink, swimming in contaminated water, contact with farm animals, through environmental contamination, or person to person spread.⁸⁻¹⁷ While the major cause of infection in humans is STEC O157:H7, other serotypes also may cause diarrhoea and HUS.¹⁸⁻²¹ Differences in the occurrence of STEC serotypes appear to be geographic; data suggest non-O157:H7 STEC strains predominate in Australia²²⁻²⁴ and contribute to disease in several other countries.^{18 20 21 25} Although typically associated with intestinal infection, HUS may also follow urinary tract infection with STEC²⁶ or systemic infection with neuraminidase producing bacteria.²⁷ HUS may be familial²⁸ or associated with immunodeficiency, immunosuppression, and pregnancy.²⁹

The epidemiology of HUS in the southern hemisphere is poorly described. Although diarrhoea associated HUS occurs in Australia, STEC O157:H7 has rarely been implicated. This may reflect the fact that diagnostic laboratories do not routinely test for this STEC in patients with diarrhoea or may indicate that HUS in Australia is caused by different STEC than are commonly implicated in Europe and North America. The frequency of STEC associated disease in Australia is unknown. We undertook nationwide surveillance of HUS, as a marker of STEC related disease, through the Australian Paediatric Surveillance Unit (APSU).³⁰ Study aims were to determine the incidence, demographic, and clinical characteristics of HUS and to examine the aetiological role of STEC. During surveillance the first major outbreak of HUS in Australia occurred,^{31 32} providing a unique opportunity to compare clinical, epidemiological, and microbiological characteristics of sporadic and outbreak HUS.

Methods

CASE ASCERTAINMENT, CASE DEFINITION, AND CASE CLASSIFICATION

Cases were ascertained prospectively through the APSU by active, national surveillance between July 1994 and June 1998 inclusive.³⁰

All paediatric specialists in Australia were sent a monthly, reply paid report card and asked to indicate whether they had seen any child under 15 years of age in the previous month with any of the conditions listed on the card, including HUS. HUS was defined as "microangiopathic haemolytic anaemia (haemoglobin <100 g/l with microscopic evidence of fragmented red blood cells), thrombocytopenia (platelet count <100 000 × 10⁹), and acute renal impairment (oliguria or anuria with increased serum urea and creatinine)". During the study period over 900 clinicians were mailed and over 93% report cards were returned to the APSU each month. The APSU notified study investigators of reported cases of HUS each month. Investigators obtained further information on individual cases (clinical presentation, outcome, demographics, and laboratory results) from clinicians notifying cases by standardised postal questionnaire (96% of questionnaires were returned). Data provided was de-identified to maintain patient anonymity. Diarrhoea associated cases of HUS were classified as "sporadic" or "outbreak" cases. The single outbreak consisted of cases occurring in South Australia in January and February 1995 and in whom a common causative organism was subsequently confirmed. Cases with no significant gastrointestinal disease or isolation of STEC in the stool and with a family history of HUS, with an atypical or relapsing course or with confirmed systemic infection with a neuraminidase producing organism, were classified as "atypical". The ethics committee of the Children's Hospital at Westmead, Sydney approved the study.

MICROBIOLOGY

Clinicians reporting cases were asked to send paired serum samples and a stool sample or rectal swab and *E coli* isolates from faeces or implicated food, if available, to a centralised laboratory for isolation and characterisation of STEC isolates. Colonies from stool specimens, obtained as soon as possible after the diagnosis of HUS and cultured on MacConkey agar, were tested for production of Shiga toxin (Stx) using HeLa cells.^{26, 33} Stx1 or Stx2 was identified by neutralisation with monoclonal antibodies. Colony and Southern blotting was used to probe STEC isolates for genes encoding virulence associated factors, namely Stx1,

Stx2, *E coli* attaching and effacing gene (*eae*, intimin), and enterohemolysin.³³ Serotyping of STEC strains was performed at the Victorian Infectious Diseases Reference Laboratory.³⁴ The relatedness of O111:H- isolates from outbreak and sporadic cases was investigated by pulsed field gel electrophoresis (PFGE) as described previously.²⁶ The DNA patterns obtained were aligned with the aid of Gel Compar software (Applied Maths, Kortrijk, Belgium) to form a composite image, and were analysed as recommended by Tenover and colleagues.³⁵ Paired serum samples (acute and convalescent, collected approximately two weeks apart) were tested for agglutinating antibodies to O antigens 26, 111, and 157.³⁶ Detailed microbiological data on outbreak cases have been reported elsewhere.²³

ANALYSIS OF DATA

Incidence estimates and confidence intervals were calculated using Stat-Exact. Comparison of outbreak and sporadic groups was made using Student's *t* test, Mann-Whitney U test, and χ^2 or Fisher's exact tests. Incidence trends were analysed using the Cochran-Armitage trend.³⁷

Results

CATEGORISATION OF CASES, INCIDENCE, SEASONAL, AND GEOGRAPHIC DISTRIBUTION

In a 48 month period, 98 cases of HUS were identified, of which 84 (86%) were associated with gastrointestinal symptoms. Sixty four were sporadic cases and 20 comprised a single outbreak in South Australia occurring in January and February 1995.^{31, 32} Fourteen "atypical" cases were identified, of whom five had an atypical or relapsing course, four were familial, three were associated with pneumococcal infection, one was associated with immunodeficiency, and one had an STEC associated urinary tract infection.²⁶ The median age of cases at diagnosis was 31 months (interquartile range 16–70 months) and 70/98 (71%) children were under 5 years of age. Ninety one (92%) children were hospitalised for a median (interquartile range) of 14 (7–21) days.

The reported annual incidence of HUS (with 95% CI) was 0.64 (0.52 to 0.78) per 10⁵ children under 15 years. Incidence was significantly higher in children under 5 years (1.35 (1.06 to 1.72) per 10⁵) than in those aged 5–14 years (0.28 (0.18 to 0.39) per 10⁵; Fisher's exact test, *p* < 0.0001). When outbreak cases were excluded from the analysis, the overall incidence was slightly, though not significantly lower.

Figure 1 shows the distribution of cases by type and month of presentation. During the entire study period there was a significantly higher incidence of cases during summer months (χ^2 test for trend, *p* < 0.001). This seasonal distribution persisted when outbreak and atypical cases were excluded. However, when the data were analysed by individual year of occurrence, a significant increase in sporadic cases in the summer only occurred in 1994–95, at the time of the outbreak. All sporadic cases



Table 1 Laboratory and clinical features of diarrhoea associated HUS

| | Outbreak (n = 20) | Sporadic (n = 64) | p value (outbreak v sporadic) |
|---|----------------------|----------------------|----------------------------------|
| <i>Laboratory features</i> | | | |
| Minimum haemoglobin (median; g/l) | 73 (62–82) | 62 (52–73) | 0.06 |
| Minimum platelet count (median; × 10 ⁹ /l) | 29 (23–44) | 56 (28–96) | <0.01 ^m |
| Maximum neutrophil count (median; × 10 ⁵ /l) | 9.3 (7.3–23.7) | 8.6 (5.2–12.6) | 0.05 |
| Minimum serum sodium (mean; mmol/l) | 128 (4) | 131 (6) | <0.05 ⁱ |
| Maximum creatinine (median; μmol/l) | 463 (322–637) | 280 (140–593) | <0.05 ^m |
| <i>Clinical features</i> | | | |
| Age (median; mth) | 63 (34–99) | 30 (18–57) | <0.05 ^m |
| Age less than 5 years (n; %) | 9 (45%) | 49 (77%) | <0.05 ⁱ |
| Duration hospital stay (median; days) | 17 (10–22) | 12 (7–17) | <0.01 ^m |
| <i>Renal features</i> | | | |
| Anuria (n; %) | 16 (80%) | 31 (48%) | <0.05 ⁱ |
| Duration anuria (median; days) | 14 (10–20) | 5.0 (3–9) | <0.001 ^m |
| Dialysis (n; %) | 18 (90%) | 35 (55%) | <0.01 ⁱ |
| Duration dialysis (mean; days) | 14.8 (8.1) | 10.2 (6.3) | <0.05 ⁱ |
| Acute hypertension (n; %) | 13 (65%) | 25 (40%) | 0.07 |
| Elevated creatinine at discharge (n; %) | 17 (85%) | 12 (19%) | <0.0001 ⁱ |
| Hypertension at discharge (n; %) | 9 (45%) | 12 (20%) | <0.05 ⁱ |
| <i>Gastrointestinal features</i> | | | |
| Diarrhoea prodrome (n; %) | 20 (100%) | 64 (100%) | |
| Duration diarrhoea (median; days) | 3.0 (2.0–4.3) | 5.0 (4.0–10.0) | <0.01 ^m |
| Blood in stool (n; %) | 16 (84%) | 23 (36%) | <0.001 ^c |
| Severe colitis (n; %) | 6 (27%) | 2 (3%) | <0.01 ^f |
| Vomiting (n; %) | 17 (85%) | 50 (78%) | 0.75 |
| Acute pancreatitis (n; %) | 1 (5%) | 0 (0%) | 0.24 |
| <i>Other</i> | | | |
| Seizures (n; %) | 5 (25%) | 7 (11%) | 0.18 |
| Central nervous system events* | 9 (45%) | 8 (13%) | <0.001 ^c |
| Hyperglycaemia | 5 (25) | 3 (5%) | <0.05 ⁱ |
| Death <12 months after diagnosis | 1 (5%)† | 2 (3%)‡§ | 0.42 |

Results are shown as n (%), mean (SD), or median (interquartile range).

Comparison between outbreak and sporadic HUS was made using: †Student's *t* test; ^mMann–Whitney U test; ^cχ² test; ^fFisher's exact test.

*Includes children with one or more of the following: seizure, intracranial haemorrhage or infarct, retinal haemorrhage, encephalopathy.

Death was due to: †multiple cerebral haemorrhagic infarction; ‡myocarditis with pericardial effusion; §complications of renal failure.

occurring during this time period were microbiologically distinct from outbreak cases. There was a significant trend in the incidence of sporadic HUS, which decreased over the four year study period (χ² test for trend, *p* < 0.001). During the entire study period none of the sporadic cases was associated with another sporadic case.

COMPARISON OF OUTBREAK AND SPORADIC CASES WITH DIARRHOEA

Table 1 shows laboratory and clinical features of children with diarrhoea associated HUS. The outbreak group was significantly older than the sporadic group and the proportion of children under 5 years was lower. There was no gender bias. Duration of hospitalisation was significantly longer in the outbreak group.

The minimum haemoglobin and maximum neutrophil counts were not significantly different between groups (table 1). Maximum serum creatinine was significantly higher in the outbreak group and minimum platelet count and minimum serum sodium concentrations were significantly lower in the outbreak group.

Eighty per cent of outbreak and 48% of sporadic cases developed anuria; its duration was significantly longer in the outbreak group (table 1). Significantly more outbreak cases required dialysis and the duration of dialysis was significantly longer in the outbreak group. The frequency of acute hypertension in groups did not differ. However, at discharge from hospital a significantly greater proportion of the

outbreak group had persistent renal impairment (increased serum creatinine) and hypertension requiring treatment (table 1).

Although a diarrhoea prodrome was present in all 84 children, outbreak cases had a shorter prodrome and a much higher rate of macroscopic bloody diarrhoea (table 1). In outbreak cases the diarrhoea prodrome lasted between one and seven days and in sporadic cases, one to 30 days. Significantly more outbreak cases had severe haemorrhagic colitis. Left hemicolectomy was required in one outbreak case with ischaemic colon and partial colectomy in another. One sporadic case required appendectomy and omentectomy. Vomiting was reported in around 80% of both groups.

Acute pancreatitis occurred in one outbreak case and hyperglycaemia was significantly more frequent in the outbreak group (table 1). Insulin was required in two outbreak cases and one sporadic case. The proportion of children with central nervous system complications (intracranial bleed or infarct, retinal haemorrhage, encephalopathy, seizure) was higher in the outbreak group. One sporadic case had anterior compartment syndrome caused by rhabdomyolysis and required fasciotomy of the right leg. An iliac vein thrombosis occurred in one child in the outbreak group.

Three of 84 patients died (table 1). A 4 year old girl with outbreak HUS and acute renal failure, multiple cerebral haemorrhagic infarcts, and extensive colonic necrosis died two days after diagnosis. A 10 year old boy with sporadic HUS and STEC O111 in the stool died eight days after diagnosis, from cardiac complications (myocarditis with pericardial effusion and tamponade). A 1 year old boy with sporadic HUS died within 12 months of diagnosis with persistent hypertension and renal failure.

OUTCOME

Twelve month follow up data were available on all 19 survivors of the outbreak and 43 of 62 survivors of sporadic HUS. Six of 19 outbreak cases and four of 43 sporadic cases had chronic renal failure (persistent elevation of creatinine and/or glomerular filtration rate of <80 ml/min/1.73 m²) at follow up (*p* = 0.06). Hypertension persisted at 12 months in four sporadic but no outbreak cases. No survivors of the outbreak developed end stage renal failure, defined as requiring chronic dialysis (whether or not awaiting transplantation) or having had renal transplantation. Of the three sporadic cases that developed end stage renal failure by 12 months after diagnosis, one had had a kidney transplant, one was awaiting transplantation, and one had died. One sporadic case had persistent insulin dependent diabetes mellitus. Five children with sporadic HUS had seizures at follow up. Three of these developed seizures following discharge from hospital, so their aetiological relation to HUS is unclear.

MICROBIOLOGY

Stool and/or serum samples were available from all outbreak cases. STEC serotype O111:H– was isolated from 16/20 stools from

Table 2 Characteristics of STEC isolates in diarrhoea associated HUS

| No. of isolates | Serotype | Sorbitol fermentation | Presence of virulence associated factors | | | |
|-----------------|----------|-----------------------|--|-------|-----|------------------|
| | | | Stx1† | Stx2† | eae | Enterohaemolysin |
| Outbreak HUS* | | | | | | |
| 16 | O111:H- | + | + | + | + | + |
| Sporadic HUS | | | | | | |
| 8 | O111:H- | + | + | + | + | + |
| 1 | O26:H- | + | + | - | + | + |
| 4 | O113:H21 | + | - | + | - | + |
| 1 | O130:H11 | + | - | + | - | + |
| 1 | O157:H- | - | + | + | + | - |
| 1 | O157:H- | - | - | + | + | + |
| 1 | O157:H-‡ | ? | ? | ? | ? | ? |
| 1 | ONT:H- | + | - | + | - | - |
| 1 | ONT:H7 | + | - | + | - | + |
| 1§ | OR:H9 | + | + | - | - | - |

Specimens were investigated on 20/20 epidemic cases and 50/64 endemic cases.

*Paton *et al.*²³

†Shiga toxin produced and Shiga toxin genes present (+) or absent (-).

‡Virulence associated features not determined on this isolate.

§Obtained from a patient who was also infected with a Shiga toxin 2 producing strain of *Enterobacter cloacae*.³⁸

NT, non-typeable; R, rough.

these children.²³ Stool specimens or bacterial cultures were investigated from 50/64 children with sporadic HUS. STEC were isolated from 20/50 specimens (table 2). These included strains of serotypes O111:H-, O26:H-, O113:H21, O130:H11, O157:H-, O non-typeable:H-, O non-typeable:H7, and O rough (R):H9. All isolates fermented sorbitol except those of serogroup O157 (table 2). One child was infected with both STEC OR:H9 and an Stx2 producing strain of *Enterobacter cloacae*.³⁸ STEC O111:H- was also isolated from the stool of the asymptomatic identical twin brother of a culture negative case of endemic HUS.³⁹ The high number (30/50) of negative stool cultures in sporadic cases is not surprising in view of the long diarrhoea prodrome (up to 30 days) prior to diagnosis of some cases in this group.

Table 2 shows virulence associated characteristics of STEC isolates. All O111:H- strains from outbreak and sporadic cases produced Stx1 and Stx2 and carried the *eae* and enterohemolysin genes. However, PFGE analysis of the outbreak and sporadic O111:H- strains showed them to be different (fig 2). Furthermore, some O111:H- strains obtained from children with HUS in New South Wales around the time of the South Australian outbreak displayed a different phage type, colicin type and had different plasmid profiles compared to the epidemic strain.³⁹ Of the non-O111:H- strains isolated and examined for the presence of virulence associated factors, all but three carried the enterohemolysin gene and produced enterohemolysin (table 2). Two of the strains that did not produce enterohemolysin were of indeterminate serotype and produced only Stx1 or Stx2. All non-O111 STEC produced either Stx1 or Stx2 but only one produced both. Five strains did not carry the *eae* gene. One sporadic case without STEC infection had a *Salmonella* species, one *Salmonella* plus *Giardia lamblia*, and two *Campylobacter jejuni* in the stool. The STEC O5:H- isolate from an atypical case with urinary tract




Figure 2 Composite image of PFGE of whole cell DNA from serogroup O111 isolates of *E. coli* digested with *Xba*I. Lane 1, PFGE marker I (Boehringer Mannheim); lane 2, outbreak case; lanes 3-7, sporadic cases. Molecular weights of the markers are indicated on the left.

infection produced Stx1 and Stx2 and carried the gene for enterohemolysin but not that for *eae*.²⁶

Serum was available from 30 children with sporadic HUS, but as only one serum sample was available from most cases, we selected a high agglutinating antibody titre (>1/640) to indicate recent infection with *E. coli* strains of serogroups O26, O111, or O157. Five patients showed increased antibody to O157, one of whom was also culture positive for STEC O157:H-. All 25 other sera investigated for antibodies to serogroup O157 showed a titre of <1/80. Three patients who were culture positive for STEC O111:H-, had an antibody titre to O111 of >1/640. The other five O111 culture positive patients identified in this study did not undergo serological examination. One patient with negative stool culture had a titre of >1/2560 to O111. No patient showed increased antibody titres to O26.

Discussion

The epidemiology of HUS in Australia has not been described previously. Four years of national surveillance through the APSU identified 98 patients under 15 years of age with HUS. Approximately 90% had diarrhoea associated HUS and 20 were affected during an outbreak of STEC O111:H- transmitted by contaminated mettwurst. Diarrhoea associated HUS was associated with at least eight different STEC serotypes. Although up to seven cases were associated with STEC O157, no strains of O157:H7 were isolated, indicating major geographical differences in strain prevalence

worldwide. Comparison of clinical features of sporadic and outbreak cases of HUS suggests the outbreak strain caused more severe disease at presentation, although specific microbial factors accounting for these differences were not identified.

The active nature of the surveillance, high return rate of monthly report cards and questionnaires, and inclusion on our mailing list of all known paediatric specialists in Australia, suggest that we identified the majority of HUS cases. No other national data set is available for comparison. The incidence of HUS in children under 5 years (1.35 (95% CI 1.06 to 1.72) per 10⁵) was lower than that reported in Canada (3.11 per 10⁵ children <5 years)³ and the British Isles (3.3 per 10⁵ children).⁴ In Minnesota the incidence of HUS in children under 5 was 5.8 per 10⁵ in 1988 and 2.1 per 10⁵ in 1994–95.^{6,40} In the later study, one HUS case was reported for every 4.3–5.2 cases of STEC infection. This greatly overestimates the ratio of HUS to gastroenteritis as many cases of STEC induced gastroenteritis are unrecognised. This is borne out by a spate of recent outbreaks of STEC O157:H7 diarrhoea in Japan affecting 9845 children, of whom 149 developed HUS.⁴¹

Previous studies suggest a summer peak of STEC infection and HUS.⁸ Overall our study showed a seasonal (summer) peak in incidence. However, when analysed by individual year, a summer peak in sporadic HUS occurred only in the year of the outbreak. Interestingly, this increase was Australia wide and not restricted to the area of the outbreak. Further, the outbreak cases occurring at this time were microbiologically distinct from sporadic cases. This suggests that climatic or environmental factors operating throughout the country may have contributed to a high background incidence of STEC infection during this period, caused by a variety of STEC, and also increased the risk of an outbreak.

Comparison of sporadic and outbreak groups indicates the outbreak strain was associated with more severe disease at presentation, but the reason for this is unclear. Epidemiological studies suggest the young and elderly have increased susceptibility to diarrhoea associated HUS but that older children have a worse outcome.^{4,6,13} In this study, outbreak patients were previously well but significantly older than those with sporadic HUS. This may reflect the fact that younger children are less likely to eat fermented sausage. While this study concurs with previous studies in finding that a short diarrhoea prodrome and bloody diarrhoea were indicative of more severe disease, there was no association between neutrophil count and severity.^{3,11,42,43} Inoculum size may also be important in determining disease severity, although the South Australian data showed low numbers of the causative agent in the food vehicle, suggesting that this was not the explanation in this outbreak.²³

Microbial factors are likely to be important determinants of disease severity. In HUS, Stx is responsible for microvascular injury causing

bloody diarrhoea and the systemic manifestations.^{7,29} The extent of damage may relate to the amount of toxin absorbed, toxin type (Stx1 or Stx2) or subtype, and/or unrecognised host susceptibility factors.⁷ Expression of the *eae* gene is required for bacterial production of intimin, an outer membrane protein involved in the attachment of the STEC to epithelial cells.⁷ STEC frequently produce enterohemolysin but the roles of this and other plasmid encoded factors in disease are unknown.⁴⁴ The O111:H– strains, which caused the outbreak and those found to be associated with sporadic HUS, exhibited all the pathogenicity traits described above. However, comparison of outbreak O111:H– isolates with several O111:H– isolates from sporadic cases revealed differences in phage type and DNA restriction fragment length polymorphism, indicating that different O111:H– strains were responsible for outbreak and sporadic HUS. Genetic differences in the outbreak strain may contribute to differences in virulence by virtue of the presence of currently unknown virulence determinants or of different quantities of known determinants. A recent Minnesota study reported that molecular subtyping of strains isolated from the local community allowed the authors to identify mini outbreaks which would not have been recognised with standard microbial characterisation.⁴⁰ Molecular subtyping is also useful for O111 STEC and may provide a mechanism to identify strains with different degrees of virulence.

In this study, several non-O157:H7 strains were identified in children with HUS, as previously recognised.^{18,21,22,25} Apart from O111, only one of these produced both Stx1 and Stx2. Three strains lacked the *eae* gene and two of these were negative for enterohemolysin. The aetiological role of such strains in HUS is open to question, given that Stx1 and Stx2 are encoded by somewhat promiscuous bacteriophages which are readily transmitted between different strains of *E. coli*.⁴⁵ Thus, the ability to produce Stx may be acquired by a relatively harmless *E. coli* resident in the intestinal tract from a virulent STEC strain that was not detected when the stool sample was taken.⁴⁵ In such cases serological investigation may shed light on the aetiological agent by permitting detection of antibodies to O antigens associated with the most frequent STEC O serogroups. This applied to four patients in this study who were culture negative, but displayed increased antibody titres to O157. Detection of antibodies to Stx1 and Stx2,⁴⁶ as well as to intimin or functionally related, chromosomally encoded proteins,⁴⁷ enterohemolysin,⁴⁸ and flagella antigens may also provide useful retrospective information about the causative organism in culture negative patients. In this study, the relatively low isolation rate (40%) of STEC overall and of O157 strains in particular, from children with sporadic HUS may have been a result of an undue delay in obtaining samples for culture and transporting them to the laboratory. Isolation rates of O157 strains could conceivably have been improved by selective enrichment, for example, by using

immunomagnetic enrichment of faecal samples. Such strategies, however, would probably have biased the data against non-O157 strains.

Medium term prognosis for diarrhoea associated HUS in this study was similar to that reported previously.^{4,5} Despite differences in disease severity at presentation, follow up at 12 months indicated no significant differences between outbreak and sporadic groups in rates of chronic renal failure and end stage renal failure. Mortality rates and prevalence of hypertension, central nervous, and gastrointestinal complications in groups were also similar.

In conclusion, nationwide HUS surveillance through APSU, which is ongoing, provided new information on the burden of HUS in Australia and enabled linkage of epidemiological, clinical, and microbiological data. We identified 12 non-O157:H7 strains of STEC associated with HUS including one O111:H-, which was responsible for an outbreak. The heterogeneity of isolates and the absence of O157:H7 differ from the current situation in most countries. The observation that O157 isolates were non-motile and failed to produce the full array of virulence associated determinants suggests a comparatively reduced virulence of Australian representatives of this serogroup and may explain their failure to become established in this country. This study was unique in allowing comparison of outbreak and sporadic diarrhoea associated HUS occurring during surveillance and indicates that outbreak cases had more severe disease at presentation. Examination of microbial pathogenicity factors of isolated strains supported the notion that the outbreak of HUS may have been associated with a more virulent strain of STEC than strains causing sporadic disease.

We acknowledge all Australian paediatricians who contribute to the APSU, and the support of the Australian and New Zealand Paediatric Nephrology Association and the Division of Paediatrics of the Royal Australasian College of Physicians. We thank the following clinicians, who provided clinical information on cases reported to this study: Drs Anderson, Andrews, Brown, Burke, Choong, Craig, Donnelly, Duncan, Ebeling, Ferson, Fiumara, Forrest, Ford, Frischman, Goldwater, Gadd, Gorton, Gray, Harris, Harding, Henning, Hewitt, Hill, Hodson, Hogg, Hu, Jaensch, James, Jaquiere, Jenkins, Jureidini, Jones, Kelso, Kainer, Ketteridge, Knight, Lammi, Lewis, Lines, Lennon, McEniery, McLennan, Mulcahy, Munt, Num, O'Loughlin, Pearson, Powell, Prebble, Pallas, Petek, Rice, Roper, Ruben, Rosenberg, Roy, Shelton, Smith, Stuart, Sparnon, Taylor, Thesinger, Thies, Thomas, Tonge, Westphalen, White, and Wheaton. Laboratory assistance was provided by A Bigham, K Bettelheim, and D Lightfoot in Melbourne, and J Paton and P Goldwater in Adelaide. The APSU was funded by the Financial Markets Foundation for Children, the Commonwealth Department of Health and Aged Care, and the Clive and Vera Ramaciotti Foundation (administered by Perpetual Trustees). Laboratory studies were supported by the National Health and Medical Research Council of Australia.

- 1 Karmali MA, Steele BT, Petric M, Lim C. Sporadic cases of haemolytic-uraemic syndrome associated with faecal cytotoxin and cytotoxin-producing *Escherichia coli* in stools. *Lancet* 1983;1:1619-20.
- 2 Karmali MA, Petric M, Lim C, et al. The association between idiopathic hemolytic uremic syndrome and infection by verotoxin-producing *Escherichia coli*. *J Infect Dis* 1985;151:775-82.
- 3 Rowe PC, Orrbine E, Wells GA, McLaine PN. Epidemiology of hemolytic-uremic syndrome in Canadian children from 1986-1988. The Canadian pediatric kidney disease reference centre. *J Pediatr* 1991;119:218-24.
- 4 Milford DV, Taylor CM, Gutteridge B, et al. Haemolytic uraemic syndromes in the British Isles 1985-8: association with verotoxin producing *Escherichia coli*. Part 1: clinical and epidemiological aspects. *Arch Dis Child* 1990;65:716-21.
- 5 Brandt JR, Fouser LS, Watkins SL, et al. *Escherichia coli* O157:H7-associated hemolytic-uremic syndrome after ingestion of contaminated hamburgers. *J Pediatr* 1994;125:519-26.
- 6 Martin DL, MacDonald KL, White KE, et al. The epidemiology and clinical aspects of the hemolytic uremic syndrome in Minnesota. *N Engl J Med* 1990;323:1161-7.
- 7 Paton JC, Paton AW. Pathogenesis and diagnosis of Shiga toxin-producing *Escherichia coli* infections. *Clin Microbiol Rev* 1998;11:450-79.
- 8 Boyce TG, Swerdlow DL, Griffin PM. *Escherichia coli* O157:H7 and the hemolytic-uremic syndrome. *N Engl J Med* 1995;333:364-8.
- 9 Bell BP, Goldoft M, Griffin PM, et al. A multistate outbreak of *Escherichia coli* O157:H7-associated bloody diarrhea and hemolytic uremic syndrome from hamburgers. The Washington experience. *JAMA* 1994;272:1349-53.
- 10 Riley LW, Remis RS, Helgerson SD, et al. Hemorrhagic colitis associated with a rare *Escherichia coli* serotype. *N Engl J Med* 1983;308:681-5.
- 11 Pavia AT, Nichols CR, Green DP, et al. Hemolytic-uremic syndrome during an outbreak of *Escherichia coli* O157:H7 infections in institutions for mentally retarded persons: clinical and epidemiologic observations. *J Pediatr* 1990;116:544-51.
- 12 Anonymous. Outbreak of *Escherichia coli* O157:H7 infections associated with drinking unpasteurized commercial apple juice—British Columbia, California, Colorado and Washington, October 1996. *JAMA* 1996;276:1865.
- 13 Carter AO, Borczyk AA, Carlson JAK, et al. A severe outbreak of *Escherichia coli* O157:H7-associated hemorrhagic colitis in a nursing home. *N Engl J Med* 1987;317:1496-500.
- 14 Swerdlow DL, Woodruff BA, Brady RC, et al. A waterborne outbreak in Missouri of *Escherichia coli* O157:H7 associated with bloody diarrhea and death. *Ann Intern Med* 1992;117:812-19.
- 15 Keene WE, McAulty JM, Hoesly FC, et al. A swimming-associated outbreak of hemorrhagic colitis caused by *Escherichia coli* O157:H7 and *Shigella sonnei*. *N Engl J Med* 1994;331:579-84.
- 16 Deschenes G, Casenave C, Grimont F, et al. Cluster of haemolytic uraemic syndrome due to unpasteurised cheese. *Pediatr Nephrol* 1996;10:203-5.
- 17 Allaby MA, Mayon-White R. *Escherichia coli* O157: outbreak in a day nursery. *CDR (Lond Engl Rev)* 1995;5:R4-6.
- 18 Johnson RP, Clarke RC, Wilson JB, et al. Growing concerns and recent outbreaks involving non-O157:H7 serotypes of verotoxigenic *Escherichia coli*. *J Food Prot* 1996;59:1112-22.
- 19 Bokete TN, O'Callahan M, Clausen CR, et al. Shiga-like toxin-producing *Escherichia coli* in Seattle children: a prospective study. *Gastroenterology* 1993;105:1724-31.
- 20 Bopp CA, Greene KD, Downes FP, et al. Unusual verotoxin-producing *Escherichia coli* associated with hemorrhagic colitis. *J Clin Microbiol* 1997;25:1486-9.
- 21 Huppertz H-I, Busch D, Schmidt H, et al. Diarrhea in young children associated with *Escherichia coli* non-O157 organisms that produce shiga-like toxin. *J Pediatr* 1996;128:341-6.
- 22 Robins-Browne R, Elliott E, Desmarchelier P. Shiga-toxin producing *Escherichia coli* in Australia. In: Kaper JB, O'Brien AD, eds. *Escherichia coli O157:H7 and other Shiga toxin-producing E. coli strains*. Washington, DC: American Society for Microbiology, 1998:66-72.
- 23 Paton AW, Ratcliff RM, Doyle RM, et al. Molecular microbiological investigation of an outbreak of hemolytic-uremic syndrome caused by dry fermented sausage contaminated with shiga-like toxin-producing *Escherichia coli*. *J Clin Microbiol* 1996;34:1622-7.
- 24 Ong J, Zhi Li, Robins-Browne R, et al. Prevalence of verotoxigenic *Escherichia coli* serogroup O157:H7 in children with diarrhoea attending a Sydney hospital. *J Paediatr Child Health* 1993;29:185-7.
- 25 Kleanthous H, Smith HR, Scotland SM, et al. Haemolytic uraemic syndrome in the British Isles 1985-8: association with *Escherichia coli*. Part 2: microbiological aspects. *Arch Dis Child* 1990;65:722-7.
- 26 Starr M, Bennett-Wood V, Bigham A, et al. Hemolytic-uremic syndrome following urinary tract infection with enterohemorrhagic *Escherichia coli*: case report and review. *Clin Infect Dis* 1998;27:310-15.
- 27 Cabrera GR, Fortenberry JD, Warshaw BL, et al. Hemolytic uremic syndrome associated with invasive *Streptococcus pneumoniae* infection. *Pediatrics* 1998;101:699-703.
- 28 Fitzpatrick MM, Walters MDS, Trompeter RS, et al. Atypical (non-diarrhea-associated) hemolytic-uremic syndrome in childhood. *J Pediatr* 1993;122:532-7.
- 29 Neild GH. Haemolytic-uraemic syndrome in practice. *Lancet* 1994;343:398-401.
- 30 Elliott EJ, Williams K. Communicable diseases and the Australian Paediatric Surveillance Unit. *CDR (Lond Engl Rev)* 1997;7:R14-16.
- 31 Henning PH, Tham EBC, Martin AA, et al. Haemolytic-uraemic syndrome outbreak caused by *Escherichia coli* O111:H-: clinical outcomes. *Med J Aust* 1998;168:552-5.
- 32 Cameron AS, Walker CC, Beers M, et al. Enterohaemorrhagic *Escherichia coli* outbreak in South Australia associated with the consumption of mettwurst. *Commun Dis Intell* 1995;19:70-1.

- 33 Robins-Browne RM, Yam WC, O'Gorman LE, Bettelheim KA. Examination of archetypal strains of enteropathogenic *Escherichia coli* for properties associated with bacterial virulence. *J Med Microbiol* 1993;**38**:222-6.
- 34 Bettelheim KA, Thompson CJ. New method of serotyping *Escherichia coli*: implementation and verification. *J Clin Microbiol* 1987;**25**:781-6.
- 35 Tenover FC, Arbeit RD, Goering RV, et al. Interpreting chromosomal DNA restriction patterns produced by pulsed-field gel electrophoresis: criteria for bacterial strain typing. *J Clin Microbiol* 1995;**33**:2233-9.
- 36 Sack RB, Sack DA. Immunologic methods for the diagnosis of infections by Enterobacteriaceae and Vibrionaceae. In: Rose NR, de Macario EC, Fahey JL, et al, eds. *Manual of clinical laboratory immunology*, 4th edn. Washington, DC: American Society for Microbiology, 1992:482-8.
- 37 Armitage P, Berry G. Further analysis of categorical data. In: *Statistical methods in medical research*, 3rd edn. Oxford: Blackwell Science, 1995:410-11.
- 38 Paton AW, Paton JC. Enterobacter cloacae producing a Shiga-like toxin II-related cytotoxin associated with a case of hemolytic-uremic syndrome. *J Clin Microbiol* 1996;**34**:463-5.
- 39 Miles TA, Bird P, Bettelheim KA. Haemolytic-uraemic syndrome in the Hunter: public health implications. *Aust N Z J Public Health* 1996;**20**:457-62.
- 40 Bender JB, Hedberg CW, Besser JM, et al. Surveillance for *Escherichia coli* O157:H7 infections in Minnesota by molecular subtyping. *N Engl J Med* 1997;**337**:388-94.
- 41 Michino H, Araki K, Minami S, et al. Recent outbreaks of infection caused by *Escherichia coli* O157:H7 in Japan. In: Kaper JB, O'Brien AD, eds. *E. coli O157:H7 and other Shiga toxin-producing E. coli strains*. Washington, DC: American Society for Microbiology, 1998:73-81.
- 42 Lopez EL, Devoto S, Fayad A, et al. Association between severity of gastrointestinal prodrome and long-term prognosis in classic hemolytic-uremic syndrome. *J Pediatr* 1992;**120**:210-15.
- 43 Bell BP, Griffin PM, Lozano P, et al. Predictors of hemolytic uremic syndrome in children during a large outbreak of *Escherichia coli* O157:H7 infections. *Pediatrics* 1997;**100**:12.
- 44 Burland V, Shao Y, Perna NT, et al. The complete DNA sequence and analysis of the large virulence plasmid of *Escherichia coli* O157:H7. *Nucleic Acids Res* 1998;**26**:4196-204.
- 45 Acheson DW, Reid J, Keusch GT, et al. In vivo transduction with Shiga toxin 1-encoding phage. *Infect Immun* 1998;**66**:4496-8.
- 46 Karmali MA. Human immune response and immunity to Shiga toxin (Verocytotoxin)-producing *Escherichia coli* infection. In: Kaper JB, O'Brien AD, eds. *Escherichia coli O157:H7 and other Shiga toxin-producing E. coli strains*. Washington, DC: American Society for Microbiology Press, 1998:236-48.
- 47 Jarvis KG, Kaper JB. Secretion of extracellular proteins by enterohemorrhagic *Escherichia coli* via a putative type III secretion system. *Infect Immun* 1996;**64**:4826-9.
- 48 Schmidt H, Beutin L, Karch H. Molecular analysis of the plasmid-encoded hemolysin of *Escherichia coli* O157:H7 strain EDL 933. *Infect Immun* 1995;**63**:1055-61.



What's in the next issue

Future content

See which articles have just been accepted for publication and preview the table of contents for the next issue a month before it is published

www.archdischild.com