

Improving mental health through parenting programmes: block randomised controlled trial

J Patterson, J Barlow, C Mockford, I Klimes, C Pyper, S Stewart-Brown

Arch Dis Child 2002;**87**:472–477

See end of article for authors' affiliations

Correspondence to:
Dr J Patterson, Health Services Research Unit, Institute of Health Sciences, Old Road, Headington, Oxford OX3 7LF, UK; jacob@ukgateway.net

Accepted 20 May 2002

Aims: To assess the effectiveness of a parenting programme, delivered by health visitors in primary care, in improving the mental health of children and their parents among a representative general practice population.

Methods: Parents of children aged 2–8 years who scored in the upper 50% on a behaviour inventory were randomised to the Webster-Stratton 10 week parenting programme delivered by trained health visitors, or no intervention. Main outcome measures were the Eyberg Child Behaviour Inventory and the Goodman Strengths and Difficulties Questionnaire to measure child behaviour, and the General Health Questionnaire, Abidin's Parenting Stress Index, and Rosenberg's Self Esteem Scale to measure parents' mental health. These outcomes were measured before and immediately after the intervention, and at six months follow up.

Results: The intervention was more effective at improving some aspects of the children's mental health, notably conduct problems, than the no intervention control condition. The Goodman conduct problem score was reduced at immediate and six month follow up, and the Eyberg Child Behaviour Inventory was reduced at six months. The intervention also had a short term impact on social dysfunction among parents. These benefits were seen among families with children scoring in the clinical range for behaviour problems and also among children scoring in the non-clinical (normal) range.

Conclusion: This intervention could make a useful contribution to the prevention of child behaviour problems and to mental health promotion in primary care.

Mental health problems of clinical severity affect up to 20% of all children aged 5–15 years in Great Britain,¹ and these are now the commonest cause of severe disability in childhood.² Mental health promotion is a priority for public health in the UK.³ The importance of parenting as a risk factor for mental illness, both in childhood^{4–5} and in adulthood is well recognised.^{6–7}

The costs to society of childhood behaviour problems (the most common form of mental illness in children) is high.⁸ It has been proposed that there is an urgent priority to shift from reactive intervention to prevention, since the later the intervention, the costlier and less effective it is.⁹

Group based parenting programmes, run both by professionals and by parents, are becoming increasingly popular in the UK and a range of different programmes are available.¹⁰ Four recent systematic reviews, one focusing entirely on group based programmes,¹¹ and three covering these programmes within wider reviews of mental health promotion and behaviour problem prevention^{12–14} have provided evidence that group based parenting programmes are an effective and cost effective way to improve parenting, and that such changes have a beneficial effect on children's mental health and behaviour. A further systematic review has shown that group based parenting programmes can also be beneficial in improving parents' mental health.¹⁵

Most trials of the effectiveness of parenting programmes have been conducted in North America, and have taken a secondary preventive (indicated) approach (working with parents of children who already have behaviour problems) or a selective primary preventive approach working with parents at high risk.¹¹ A recent study in the UK has shown the effectiveness of the Webster-Stratton parenting programme in an indicated approach among parents of children who had been referred to child mental health services.¹⁶

Research into the effectiveness of group based parenting programmes in the UK is only just beginning. There are few

well designed trials in diverse populations and settings,¹⁷ and to our knowledge there are no randomised controlled trials of these interventions in UK primary care. In this study, we assessed the effectiveness of a primarily behavioural parenting programme, the Webster-Stratton Parents and Children Series,¹⁸ of proven effectiveness in UK clinical populations, delivered by health visitors in a general practice setting, in terms of its impact on child behaviour problems and parental mental health. This sample included both parents of children with behaviour problems in the clinical range, and parents whose children scored in the normal range.

METHODS

A postal survey was administered to the parents of all the children aged 2–8 years inclusive, registered at three general practices in Oxford (n = 1788).¹⁹ This survey comprised the Eyberg Child Behaviour Inventory (ECBI)²⁰ and demographic information. A total of 1105 questionnaires were returned, giving information on children in 800 families (response rate 70%). The mean score on the Eyberg Intensity Scale in this population was 102.7 and the median value was 100. A total of 193 children (17.9%) scored above the clinical cut off point of 127.

The parents of all children scoring above the median value (487 children in 391 families) were invited to participate in a randomised controlled trial of the Family Nurturing Network's parenting programme (Webster-Stratton's Parents and Children Series). The median rather than the mean was used because of the skewed distribution of behaviour scores.

Abbreviations: ECBI, Eyberg Child Behaviour Inventory; GHQ, General Health Questionnaire; PSI, Parenting Stress Index; SDQ, Strengths and Difficulties Questionnaire

Parents of either sex and any ethnic group were eligible. Children already receiving treatment for behaviour problems (27 children, 2.5%) and those with learning difficulties (78 children 7.1%) were excluded. These children were felt to be special cases warranting subgroup analysis for which the trial was not powered. A sample size calculation gave a total sample size of 100 to allow detection of an effect size of 0.6 (a moderate effect)²¹ on the Eyberg Child Behaviour Inventory Intensity Score with 80% power at the $p < 0.05$ level of significance, or 120 to detect an effect size of 0.5, and 116 parents were actually recruited into the trial. (See web supplement for full description of sample size calculation.)

Parents were invited to join the trial by a letter addressed to the parent who had returned the survey questionnaire. Parents who consented to join the trial were compared with eligible parents who did not consent, with regard to the mean intensity scores of their children, whether they were defined as a “case” on the intensity score or not, and by social class based on the respondent’s occupation.

We adopted a block randomisation strategy. The parents who consented to join the study ($n = 116$; uptake = 30%) were allocated to one of two groups with the aim of achieving groups balanced in terms of child “caseness” on the Eyberg intensity score, age, sex, social class, and ethnicity,²² as well as preferences for attendance. These two groups were randomised, by tossing a coin in the presence of an independent witness, to treatment or control. It is particularly important to take parents’ preferences into account in the implementation of health promotion interventions, and this method of randomisation allowed us to allocate parents according to the time of day and day of the week they were available to attend, as well as their need for a crèche, at the same time as producing six viable groups of 10 parents for delivery of the programme. Each consenting parent had an equal chance of being in the control or intervention group. The allocation to group was undertaken blind to general practitioner practice and area of residence, so neither of these factors influenced allocation to control or treatment group.

The intervention comprised sessions lasting about two hours each, once a week for 10 weeks. The activities included video vignettes of parent-child interactions, group discussion, role play, rehearsal of parenting techniques, and home practice. The techniques covered included play and positive interaction with the child, clear commands, limit setting, ignoring undesirable behaviour, praising and rewarding desirable behaviour, and following through on discipline. Four of the programmes were run at a health centre in the evenings between 7 30 and 9 30 pm. The other two programmes were run at a local community centre in the daytime, one between 9 30 and 11 30 am (with a crèche available) and the other between 12 noon and 2 00 pm. Each group was run by at least one health visitor with either a second health visitor or a nursery nurse as co-leader, all of whom had received three days of training from the Family Nurturing Network in Oxford, specifically to deliver this programme. The group leaders (seven in all) received weekly supervision meetings as a group during the delivery of the programmes, to provide support and ongoing training, and ensure equivalence of the six intervention programmes and integrity of programme delivery compared with the instruction manual.

The 56 control group parents received no extra intervention.

Self completion questionnaires were administered to all consenting parents prior to the intervention, immediately post-intervention, and at six months post-intervention. The primary outcome measure was the Eyberg Child Behaviour Inventory (ECBI). The ECBI was developed in the USA from parent reports of children’s problems behaviour and has two scales, one which measures the intensity or frequency of the problem behaviour, and the other which measures the extent to which this behaviour is a problem for the parents. The number of items and range of responses ensure that this scale

Table 1 Consenters are representative of all eligible families

	Consenters	Eligible non-consenters
Mean intensity score (SD)	125.3 (18.5)	121.5 (18.7)
Clinical range score	39.4%	29.5%
Social class I	13.3%	10.6%
Social class II	39.0%	32.4%
Social class III N	22.9%	28.9%
Social class III M	13.3%	13.1%
Social class IV	5.7%	10.5%
Social class V	5.7%	4.4%
Single parent	14.7%	12.7%
Asian	4.8%	5.5%
Black	1%	0.8%
Mixed race	2.9%	3.4%
White	91.4%	89.8%

None of the differences are statistically significant.

is sensitive to change, but it includes “maturational” items such as “wets the bed” which may change as the child matures. Secondary outcome measures included the Strengths and Difficulties Questionnaire (SDQ).²³ This was developed in the UK and includes prosocial as well as antisocial behaviours. This scale does not have the “maturational” items, but focuses on temper tantrums, fighting, and stealing. It also contains an “impact” measure of the effect of the child’s behaviour on their friendships, learning, leisure, and family life. It is generally recommended for use in children aged 4 years and above. The General Health Questionnaire (GHQ)²⁴ was used to capture the parents’ level of anxiety and depression; the Parenting Stress Index (PSI)²⁵ was used to measure changes in the specific stresses associated with the parenting role; and the Self Esteem Scale²⁶ was used to measure the parents’ self esteem.

The data were analysed using SPSS version 9. All analyses were carried out on an intention to treat basis, including in the intervention group both parents who had attended the parenting programme and those who had not. The statistical significance of changes in scores from baseline to post-intervention and six month follow up, within the control and intervention groups separately, was calculated using paired *t* tests for normally distributed outcome measures (the hyperactivity subscale of the SDQ and the SDQ total score, the PSI parent and difficult child domains, and the PSI total score). The Wilcoxon matched pairs signed rank sum test was used for outcome measures which were not normally distributed (ECBI intensity and problem scores, SDQ conduct, emotional, peer problems, prosocial and impact scores, PSI parent child interaction domain, GHQ somatic symptoms, anxiety, social dysfunction, depression and total scores, and the SES). Grouped *t* tests were used to compare the mean change in scores in the control and intervention groups where the differences were normally distributed (ECBI intensity score, SDQ total score, PSI parent child interaction, and parent domains), and Mann-Whitney U tests for the mean change in scores in the two groups where the differences were not normally distributed (ECBI problem score, SDQ conduct, hyperactivity, emotional, peer and prosocial scales, GHQ somatic anxiety, social, depression and total scores, PSI difficult child domain and total score, and SES).

RESULTS

The parents who consented to join the trial were representative of all the eligible families, in terms of the behaviour scores, social class, single parents, and ethnicity, as shown in table 1. However owing to over representation of non-manual




Figure 1 Attrition.

families in the survey, manual families were under represented compared with population norms. Consenting parents had slightly more children with behaviour in the clinical range, but this difference was not significant.

Thirty four of the 60 parents randomised to the intervention group attended 50% or more of the sessions (fig 1). Reasons for dropping out of the programme included increased work

commitments, moved away from the area, depression, other life stress, holiday fell at start of group sessions, missed too much to begin. Only one potential participant stated that the group was not helpful as a reason for dropping out. Forty six control group and 46 intervention group participants (79% of all initial participants) returned questionnaires at the six month follow up. There was no difference in the proportion of attenders or non-attenders returning questionnaires at follow up.

Changes between baseline and follow up scores in the intervention and control groups separately

Children in the intervention and control groups showed a significant ($p < 0.05$) reduction in ECBI intensity score (the primary outcome measure) at immediate and six month follow ups (table 2). The mean ECBI problem score was also significantly reduced among both the intervention and control groups at six months.

Intervention group scores on the SDQ were reduced at both the immediate follow up and at six months. While there was no reduction in the control group score post-intervention, there was a significant reduction at six months. Intervention group SDQ conduct scores were significantly reduced at both time points whereas control group conduct scores showed no change. Changes on the emotional subscale were significant in the intervention group at the immediate follow up and in the

Table 2 Mean (SD) scores and number of respondents at each time point

	Baseline		Immediate follow up		6 month follow up	
	Control	Intervention	Control	Intervention	Control	Intervention
<i>Eyberg</i>						
Intensity score	126.6 (16.9)	125.8 (22.8)	117.9* (21.8)	113.3* (23.5)	118.8* (20.3)	110.2* (21.6)
	56	60	50	46	46	45
Problem score	10.2 (7.2)	9.9 (9.5)	9.4 (6.8)	8.1 (7.7)	8.8* (6.5)	6.2* (6.3)
	54	52	47	40	41	38
<i>SDQ</i>						
Total difficulties	12.0 (5.4)	12.4 (6.2)	11.1 (5.7)	9.4* (5.7)	10.1* (4.7)	9.5* (6.4)
	48	47	50	46	45	46
Conduct problems	2.6 (1.8)	2.8 (1.9)	2.5 (1.6)	2.0* (1.5)	2.4 (1.6)	1.9* (1.5)
	50	52	50	46	45	46
Emotional score	2.9 (2.3)	2.8 (2.1)	2.6 (2.0)	1.8* (1.8)	2.1* (1.9)	2.2 (2.3)
	50	51	50	46	45	46
Hyperactivity	4.4 (2.5)	5.1 (3.0)	4.2 (2.6)	4.1* (2.5)	3.9* (2.3)	4.0* (2.3)
	48	51	50	46	45	46
Peer problems	2.0 (1.7)	1.8 (1.9)	1.9 (1.9)	1.4 (1.7)	1.7 (1.7)	1.5 (1.9)
	50	49	50	46	45	46
Prosocial score	6.7 (1.4)	7.3 (1.7)	6.8 (1.5)	7.8* (1.7)	7.0 (2.0)	7.7* (1.6)
	50	51	50	46	45	46
Impact	1.0 (1.2)	0.9 (1.5)	0.7* (1.2)	0.7 (1.4)	0.7* (1.0)	0.7 (1.4)
	50	51	50	43	46	45
<i>GHQ</i>						
Total score	4.3 (4.9)	5.1 (4.9)	4.7 (5.6)	3.9 (5.3)	3.0 (4.6)	2.7* (4.2)
	49	51	49	45	46	45
Anxiety	1.5 (1.9)	1.6 (1.9)	1.8 (2.2)	1.2 (1.9)	0.9* (1.7)	1.0* (1.8)
	50	51	50	46	46	45
Somatic symptoms	1.7 (2.2)	1.7 (1.8)	1.3 (1.9)	1.7 (2.2)	1.2 (1.9)	1.2* (1.8)
	50	51	49	46	46	45
Depression	0.2 (0.6)	0.7 (1.5)	0.4 (1.1)	0.3 (0.8)	0.09 (0.4)	0.07* (0.3)
	49	51	50	45	46	45
Social dysfunction	1.0 (1.6)	1.0 (1.7)	1.2 (1.9)	0.7* (1.5)	0.8 (1.8)	0.4* (1.3)
	49	51	50	46	46	45
<i>Parenting Stress Index</i>						
Total	86.5 (18.4)	85.0 (20.4)	83.3 (17.5)	79.8* (18.7)	83.4 (17.0)	79.0* (20.9)
	50	51	50	46	46	46
Parent domain	29.4 (7.2)	29.5 (9.2)	28.9 (7.6)	27.9 (7.9)	29.0 (7.1)	27.7 (8.6)
	50	51	50	46	46	46
Difficult child	34.4 (8.9)	32.2 (8.3)	32.3* (8.7)	30.4 (8.4)	32.2 (8.3)	30.0* (9.1)
	50	51	50	46	46	46
Parent-child interaction	22.7 (5.9)	23.3 (6.2)	22.1 (5.7)	21.5 (5.5)	22.2 (5.4)	21.7* (6.4)
	50	51	50	46	46	46
<i>Self Esteem Score</i>						
	29.7 (4.7)	29.2 (5.0)	30.7* (5.5)	29.8 (4.7)	30.3 (4.7)	29.5 (4.4)
	50	51	50	46	46	46

For prosocial and SES scores, increase = improvement; *significant change from baseline at $p < 0.05$.

Table 3 Results of grouped *t* tests and Mann-Whitney U tests to show *t* or *z* values, degrees of freedom, and *p* values for the significant differences in the changes in scores between the control and intervention group

	Immediate follow up	6 month follow up
<i>Eyberg</i>		
Intensity score	<i>p</i> =0.39	<i>t</i> =2.29, 89 df, <i>p</i> =0.024
Problem score	<i>p</i> =0.29	<i>p</i> =0.95
<i>SDQ</i>		
Conduct problem score	<i>t</i> =2.08, 86 df, <i>p</i> =0.041	<i>t</i> =2.15, 80 df, <i>p</i> =0.034
Emotional score	<i>p</i> =0.50	<i>p</i> =0.57
Hyperactivity	<i>p</i> =0.26	<i>p</i> =0.57
Peer problems	<i>p</i> =0.84	<i>p</i> =0.83
Prosocial score	<i>p</i> =0.17	<i>p</i> =0.14
Total difficulties	<i>p</i> =0.09	<i>p</i> =0.34
Impact	<i>p</i> =0.15	<i>p</i> =0.46
<i>GHQ</i>		
Total	<i>p</i> =0.13	<i>p</i> =0.24
Anxiety	<i>p</i> =0.18	<i>p</i> =0.92
Somatic symptoms	<i>p</i> =0.89	<i>p</i> =0.45
Depression	<i>p</i> =0.26	<i>p</i> =0.09
Social dysfunction	<i>z</i> =2.5, <i>p</i> =0.012	<i>p</i> =0.20
<i>Parenting Stress Index</i>		
Total	<i>p</i> =0.76	<i>p</i> =0.20
Parent domain	<i>p</i> =0.43	<i>p</i> =0.48
Difficult child domain	<i>p</i> =0.72	<i>p</i> =0.38
Parent child interaction	<i>p</i> =0.76	<i>p</i> =0.09
<i>Self Esteem Score</i>		
	<i>p</i> =0.52	<i>p</i> =0.84

p values shown for non-significant results. All differences favour the intervention group over the control group.

control group at the six month follow up only. Changes on the hyperactivity subscale were significant at both time points in the intervention group, but only at six months in the control group. On the peer relationships subscale there were no significant differences at either time point in either group. The SDQ prosocial subscale, measuring increases in the child's positive behaviours, showed a benefit in the intervention group at both immediate and six month follow ups. The control group did not show a significant difference at either time point on this subscale. The control group however did show significant improvement on the SDQ impact score at both time points.

Intervention group scores on the GHQ total scores were significantly lower than baseline at the six month follow up. This reflected significant reductions in all the subscales at six months. A reduction in the intervention group social dysfunction subscale was also observed at the immediate follow up. The control group showed a significant reduction on one measure only, the GHQ anxiety score at six months.

The intervention group showed significant improvement on the Parenting Stress Index total score at both follow ups. This reflected primarily improvements in the difficult child and parent-child interaction domains at the six month follow up. The control group improved on the difficult child domain at the immediate follow up, but this was no longer significant at six months.

The control group showed a significant improvement in the Self Esteem Scale at the immediate follow up, but this had disappeared by six months.

Differences between the intervention and control groups

The main aim of the trial was to detect a difference between the control and intervention groups in the changes observed in the Eyberg child behaviour inventory intensity score.

Although the intensity scores of the intervention group fell more than the control group at the immediate follow up, this difference was initially not significant (table 3). However, by the six month follow up, grouped *t* tests to compare the mean differences in scores showed that the intervention group ECBI intensity scores had changed significantly more than the control group (*p* = 0.024).

This difference was mirrored by the significant differences favouring the intervention group on the Goodman conduct problem score at both time points.

None of the other between group comparisons reached significance, except for the social subscale of the GHQ which favoured the intervention group immediately after the intervention. This difference was no longer significant by six months.

This trial was not powered to detect a difference between subgroups in the control and intervention groups. However, it was noted that the mean Eyberg Intensity scores of the 21 intervention group children whose initial scores fell in the clinical range decreased by 26.1 points from preintervention to six month follow up (*p* < 0.001) and that those of the 39 children initially scoring in the normal range decreased by 9.2 points (*p* = 0.002) over this period. Scores of the 25 control group children with initial scores in the clinical range decreased by only 9.3 points (*p* = 0.001), and those in the normal range (*n* = 31) decreased by 5.9 points (not significant).

DISCUSSION

Our initial survey had a good response rate of 70%. While all social groups were well represented in the trial, the sample from which the study was drawn was not entirely representative. Manual class families were under represented and ethnic minority families over represented. The families who consented to enter the trial were representative of eligible families from a socioeconomic point of view and they were also more likely to have a child with clinical level behaviour problems than were those who refused. This single study cannot provide evidence about what would happen were such a programme to be offered in other areas, and the under representation of manual groups detracts from the generalisability of the results. However, the fact that parents from all social groups, including unskilled manual workers, minority ethnic groups, and single parents did attend and that the programme preferentially attracted parents of children with behaviour problems must be viewed as encouraging.

After excluding children with learning disabilities and previous treatment for behaviour problems, 30% of families consented to enter the trial, to be randomised to receive a 10 week parenting intervention or not, and to complete several questionnaires. Given the number of other commitments that families of children in the 2–8 year age range are likely to have, this represented a significant level of interest in the intervention on offer. If the intervention were to be rolled out as a universal service, uptake might be expected to be higher than the 30% achieved for a trial, since parents would not be making a commitment to filling in questionnaires or to being randomised. In addition, if parenting programmes were regularly available within general practices, uptake would be cumulative with additional chances for parents to participate later if they could not attend the first group to which they were invited. This could also reduce the dropout rate because interested parents who found one group inconvenient to attend could participate in another. Although the prevalence of unhelpful parenting practices is high,²⁷ some parents will not need this intervention because they have already developed the skills and attitudes required of helpful parenting. Conversely it is likely that some parents, particularly those living in the most stressful social circumstances, will never be reached by this type of intervention; and for this group, alternative, more intensive, tailored programmes may be required.

Thus a goal of 100% attendance is likely to be both unrealistic and unnecessarily high.

Allocation to group was achieved by block rather than individual randomisation. This method, although possibly inferior from a statistical point of view, was considered important from the point of view of this trial. Health promoting interventions such as this are different from clinical interventions. In the latter, parents who have consulted a doctor about their child's behaviour can be expected to accept whatever treatment is offered and to make time to attend a group. Our trial invited parents who did not necessarily recognise that they had a problem to attend a group. In this situation it is more important to ensure that invitations take their preferences for attendance (for example, time of day and day of week) into account. In addition the randomisation process needed to deliver groups of 10–12 parents who could attend on the same day and time. The stratification process ensured an even distribution of potential confounding variables between the control and intervention groups.

Only 57% of those randomised to the intervention attended, but the trial had a good level of follow up, with responses from 79% of original participants at six months. The analyses were completed on an intention to treat basis so that responses from non-attenders were included in the intervention group, along with those of parents who attended the groups. This approach avoids bias attributable to unidentified differences between attenders and non-attenders and tends to an underestimate of the effect which would have been achieved had all those randomised into the intervention group actually attended.

The improvements we observed in control group children's scores on the main outcome measure (ECBI) are potentially attributable to regression to the mean. We selected children for invitation to the trial on the basis of their above average scores on this measure. The ECBI, however, also includes items such as "wets the bed" which tend to resolve as the child gets older. Comparison with control group changes in SDQ scores, which do not include such items, suggests that some of the control group improvement we observed is likely to be attributable to a maturation effect. Unlike the ECBI improvement, the improvement in SDQ scores in control group children did not reach significance. On the other hand the SDQ questionnaire is intended for children over the age of 4 years, and some of the children in the trial were aged 3. We ran all the analyses on the SDQ excluding children under 4 years of age at recruitment, and the results remained as reported.

Our observation that the improvement in both the ECBI intensity score, a measure based primarily on problem behaviours, and the SDQ (conduct) scores was significantly greater in the intervention than the control group provides confidence that the intervention was effective, at least as far as these aspects of children's mental health was concerned. The fact that the decline in scores on both measures in the intervention group was maintained over the six months following the intervention suggests that the intervention had an enduring effect on behaviour. It also suggests that the changes are unlikely to be caused by an immediate "feel good" factor attributable to attending a supportive group, or to the impact of lack of blinding of participants.

Although the changes shown in table 2 are suggestive of a beneficial impact on other aspects of children's mental health, the evidence here is not robust. When the small changes in the control group were taken into account there was no significant differences between groups. The Webster-Stratton Parents and Children Series is predominantly a behaviour management programme and it is therefore to some extent reassuring that it was this aspect of children's mental health which was most effected by the intervention. It would appear that there is a need to develop alternative or additional strategies or programmes to improve anxiety, hyperactivity, and prosocial behaviour.

Improvements observed in the mental health scores of mothers in the intervention group were also, with the

exception of social dysfunction at immediate follow up, not statistically robust. They are reassuring to the extent that mothers' mental health did not suffer as a result of the intervention and, in terms of social dysfunction, the effect was beneficial. The latter finding could, however, be attributable, not to any specific effect of the programme, but to the "feel good" factor referred to above. The Webster-Stratton Parents and Children Series, in contrast to some other parenting programmes, does not focus on parents' mental health and ways in which it could be improved. Any effect on parents' mental health would therefore be secondary to improvements in the parent-child relationship or to the non-specific effect of group support.

The trial findings on parents' mental health contrast with those obtained in a parallel qualitative study in which participants in this trial reported feeling more in control, with greater ability to cope with their children and a better relationship with them as a result of the intervention.²⁸ It is possible that such changes were not detected in the trial because the instruments used are not particularly sensitive to improvement in positive mental health.

Self report measures were selected owing to the resource intensive nature of independent observations, which were not practical for the purposes of this study. Although all measures of child problem behaviour were based on parental reports, these were all valid and reliable measures, and the specificity of the beneficial results to the predicted scales argues against the results being attributed to biased positive reporting by parents with elevated mood or improved social support as a result of the programme. In addition, a number of studies have shown adequate correlation between the results from parent reports and independent observations.¹⁶

Parents were invited to attend this trial if their children's scores on the ECBI were above average. Although parents of children whose scores fell in the clinical range were slightly more likely to accept the invitation than parents whose children's scores were normal, both groups were well represented in the trial, and our results suggest that both groups benefited. These results lend support to the feasibility of a population approach to the promotion of mental health through parenting programmes. They are consistent with results from an observational study,¹⁹ suggesting that the parents of children most in need are more likely to accept an invitation to a group than parents of other children, and that these two groups can derive benefit from attending the same group together. There remains a need, which this trial cannot fulfil, of showing improvements in positive mental health as a result of parenting programmes among children whose mental health is already above average. In order to establish the benefits of such a truly population approach, a much larger trial would be needed, powered to detect small changes. The conduct of such a trial would be greatly facilitated by the development of outcome measures which captured the full range of the components of positive mental health.

ACKNOWLEDGEMENTS

The authors would like to thank the participating parents, and the health visitors and nursery nurses who ran the parenting groups in this trial. Both the project and the Health Services Research Unit were supported by grants from the NHS Executive South East Regional Office. Dr Jane Barlow and Dr Cecilia Pyper are supported by grants from the Department of Health. This work was approved by the Oxford Applied & Qualitative Research Ethics Committee (formerly NAPREC).

.....

Authors' affiliations

J Patterson, J Barlow, C Mockford, C Pyper, S Stewart-Brown, Health Services Research Unit, Institute of Health Sciences, Old Road, Headington, Oxford OX3 7LF, UK

I Klimes, John Radcliffe Hospital, Oxford, UK



Supplementary information (expanded introduction and reference list) can be viewed on the ADC website (www.archdischild.com)

REFERENCES

- 1 **Campbell SB.** Behavior problems in preschool children: a review of recent research. *J Child Psychol Psychiatry* 1995;**36**:113–49.
- 2 **Bone M, Meltzer H.** *The prevalence of disability among children, OPCS Surveys of Disability in Great Britain, Report 3.* London: HMSO, 1989.
- 3 **Government White Paper.** *Saving lives: our healthier nation.* London, 1999.
- 4 **McCord W, McCord J, Howard A.** Familial correlates of aggression in nondelinquent male children. *Journal of Abnormal and Social Psychology* 1961;**62**:79–93.
- 5 **Patterson GR, DeBaryshe BD, Ramsey E.** A developmental perspective on antisocial behavior. *Am Psychol* 1989;**44**:329–35.
- 6 **Parker G.** Parental 'affectionless control' as an antecedent to adult depression: a risk factor delineated. *Arch Gen Psychiatry* 1983;**40**:956–60.
- 7 **Oakley-Browne MA, Joyce PR, Wells JE, et al.** Adverse parenting and other childhood experience as risk factors for depression in women aged 18–44 years. *J Affect Disord* 1995;**34**:13–23.
- 8 **Scott S, Knapp M, Henderson J, et al.** Financial cost of social exclusion: follow up study of antisocial children into adulthood. *BMJ* 2001;**323**:191–4.
- 9 **Hoghugh M.** The importance of parenting in child health. *BMJ* 1998;**316**:1545.
- 10 **Wolfendale S, Enzig H.** *Parenting education and support.* London: David Fulton Publishers, 1999.
- 11 **Barlow J, Stewart-Brown S.** Behaviour problems and group based parenting education programmes. *Developmental and Behavioral Pediatrics* 2000;**21**:356–70.
- 12 **Serketich WJ, Dumas JE.** The effectiveness of behavioral parent training to modify anti-social behavior in children: A meta-analysis. *Behavior Therapy* 1996;**27**:171–86.
- 13 **Lister-Sharp D, Chapman S, Stewart-Brown S, et al.** Health promoting schools and health promotion in schools: two systematic reviews. *Health Technol Assess* 1999;**3**(22).
- 14 **Marshall J, Watt P.** *Child behaviour problems.* Perth: Interagency Committee on Children's Futures, 1999.
- 15 **Barlow J.** Individual and group-based parenting programmes for improving psychosocial outcomes for teenage parents and their children. Cochrane protocol. *The Cochrane Library*, Issue 3. Oxford: Update Software, 2000.
- 16 **Scott S, Spender Q, Doolan M, et al.** Multicentre controlled trial of parenting groups for childhood antisocial behaviour in clinical practice. *BMJ* 2001;**323**:194–7.
- 17 **Webster-Stratton C.** Commentary: nipping conduct problems in the bud. *BMJ* 2001;**323**:197–8.
- 18 **Webster-Stratton C, Hollinsworth T, Kolpacoff M.** The long-term effectiveness and clinical significance of three cost-effective training programs for families with conduct problem children. *J Consult Clin Psychol* 1989;**57**:550–3.
- 19 **Patterson J, Mockford C, Barlow J, et al.** Need and demand for parenting programmes in general practice. *Arch Dis Child* 2002;**87**:468–77.
- 20 **Robinson EA, Eyberg SM, Ross AW.** The standardization of an inventory of child conduct problem behaviors. *J Clin Child Psychol* 1980;**9**:22–9.
- 21 **Cohen J.** *Statistical power analysis for the behavioural sciences.* New York: Academic Press, 1977.
- 22 **Altman DG, Bland JM.** How to randomise. *BMJ* 1999;**319**:703–4.
- 23 **Goodman R.** A modified version of the Rutter parent questionnaire including extra items on children's strengths: a research note. *J Child Psychol Psychiatry* 1994;**35**:1483–94.
- 24 **Goldberg DP, Hillier VF.** A scaled version of the General Health Questionnaire. *Psychol Med* 1979;**9**:139–45.
- 25 **Abidin R.** Parenting Stress Index: a measure of the parent-child system. In: Zalaquett C, Wood R, eds. *Evaluating stress: a book of resources.* Lanham, MD: Scarecrow Press, 1997:277–91.
- 26 **Rosenberg M.** *Conceiving the self.* New York: Basic Books, 1979.
- 27 **Nobes G, Smith M.** Physical punishment of children in two parent families. *J Clin Child Psychol Psychiatry* 1997;**2**:271–81.
- 28 **Patterson J, Mockford C, Barlow J, et al.** The impact of parenting programmes: a qualitative study. Forthcoming.

Website supplement for: Improving mental health through parenting programmes: Block randomised controlled trial (Patterson et al)

Introduction

Mental health problems of clinical severity affect up to 20% of all children aged 5-15 years in Great Britain¹ and these are now the commonest cause of severe disability in childhood.² Mental health promotion is a priority for public health in the UK³ and mental health in childhood is an important determinant of mental health in adult life.⁴ It is also an important determinant of criminality and violence.

The importance of parenting as a risk factor for mental illness both in childhood^{5, 6} and in adulthood is well recognised.^{7, 8} The parenting practices identified as important for the development of mental health have been investigated in a number of longitudinal studies. Authoritative parenting styles (firm, consistent discipline combined with warmth and support), in contrast to authoritarian (characterised by 'affectionless control') or permissive (neglecting) parenting, supports the development of positive mental health predicting social competence, educational success, better self-esteem and less aggression in later childhood and adolescence.⁹

Conversely, parenting practices characterised by little positive parental involvement with the child, harsh and inconsistent discipline, and poor monitoring and supervision predict behaviour problems in early childhood.⁶ These practices are also an important predictor of delinquency, criminality

and violence.⁵ The costs to society of such behaviour problems is high.¹⁰ It has been proposed that there is an urgent priority to shift from reactive intervention to prevention, since the later the intervention, the costlier and less effective it is.¹¹

Different approaches have been developed to improve parents' insight, skill and behaviour, including books, magazines such as Practical Parenting, videos¹² and television programmes such as the Triple P programme (Sanders, personal communication), telephone advice lines, drop in centres and parenting programmes. Two main approaches have informed the development of parenting initiatives, derived from social learning theory (behavioural programmes) and psychotherapeutic theory (relationship programmes). These approaches are used by psychologists and health visitors to coach parents in clinics and at home, and in mutual support from trained parents¹³ as well as in group based programmes.

Group based parenting programmes, run both by professionals and by parents, are becoming increasingly popular in the UK and a range of different programmes are available.¹⁴ Four recent systematic reviews, one focusing entirely on group based programmes¹⁵ and three covering these programmes within wider reviews of mental health promotion and behaviour problem prevention^{16, 17, 18} have provided evidence that group-based parenting programmes are an effective and cost-effective way to improve parenting, and that such changes have a beneficial effect on children's mental health

and behaviour. A further systematic review has shown that group based parenting programmes can also be beneficial in improving parents' mental health.¹⁹

Most trials of the effectiveness of parenting programmes have been conducted in North America, and have taken a secondary preventive (indicated) approach (working with parents of children who already have behaviour problems) or a selective primary preventive approach working with parents at high risk.¹⁵ A recent study in the UK has demonstrated the effectiveness of the Webster-Stratton parenting programme in an indicated approach among parents of children who had been referred to child mental health services.²⁰

The symptoms of mental illness show a unimodal distribution in population surveys both in children and in adults²¹ with no clear cut-off point between 'cases' and 'normal' people. The high prevalence of mental health problems and the unimodal distribution of symptom scores provide theoretical evidence to support a universal or population based approach to prevention.²² Such an approach would be in keeping with the view of many non-government organisations that mental health promotion should aim for the promotion of positive mental health – improving the mental health of 'psychiatrically normal' people as well as preventing mental illness.²³

There are strong practical arguments in favour of universal approaches. These include the inefficiency of selective approaches and the avoidance of stigma associated with programmes for parents who are perceived as 'failing'.²⁴ They are also supported on the basis of observations that parenting practices which are demonstrably unhelpful for children's development^{25, 13} are very common.²⁶

Several of those writing on the subject¹⁸ have suggested that it is important to provide a range of approaches - universal as well as selective and indicated.

A problem encountered by those who attempt to evaluate universal approaches to provision, however, is the lack of widely accepted measures of positive mental health. Most validated measures have been developed among clinical populations, and these measures have important 'floor' effects when used in trials of the universal approach (meaning that too many children return an optimum score at the beginning of the trial, leaving no room to measure improvement).

Research into the effectiveness of group based parenting programmes in the UK is only just beginning. There are few well designed trials in diverse populations and settings,²⁷ and to our knowledge there are no randomised controlled trials of these interventions in UK primary care. In this study, we assessed the effectiveness of a primarily behavioural parenting programme,

the Webster-Stratton Parents and Children Series,²⁸ of proven effectiveness in UK clinical populations, delivered by health visitors in a general practice setting, in terms of its impact on child behaviour problems and parental mental health. We aimed to test the effectiveness of a universal approach to provision, but because of concerns with the limitations of available outcome measures we invited parents of children whose behaviour fell in the upper (worse) half of the behaviour distribution. This sample included both parents of children with behaviour problems in the clinical range, and parents whose children scored in the normal range.

Sample size calculation to determine numbers of participants required

The sample size calculation was based on a study of parent-child interaction therapy (PCIT).ⁱ This is similar to the intervention used in our study. The study provided sufficient data to determine the effect sizesⁱⁱ for the effects demonstrated. In that study, children aged 3 to 6 years were randomly allocated to intervention or a waiting list control group. The mean length of the intervention was 13 weeks. Data at 4 months on 10 treatment versus 6 waiting list children showed a change in mean Intensity score on the ECBI from 159.5 (SD 16.6) to 117.5 (SD 18.8) in the treated group and 170.7 (SD 40.3) to 177.2 (SD 62.0) in the control group ($p < 0.02$). This study therefore showed a change of over two effect sizes in the treatment group.

However, their study was examining effectiveness among children whose mean score was initially extremely high (a clinical population; mean score 159.5 compared with a clinical cut-off score of 127 and a mean total population score of 100) rather than our sample which was to be drawn from the general population. Our sample size calculation using the program 'Power and Precision'ⁱⁱⁱ calculated the required number to demonstrate a change in the treatment group of one effect size (17.7 points on the Intensity scoring scale, using the pooled standard deviations). If 30 parents were recruited to each of the intervention and control groups, an effect of this magnitude would be demonstrated with 97% power (at a significance level of 0.05 (two-sided)) for a single intervention group. As there were multiple intervention groups in our study, since the intervention is recommended to be delivered in small groups of no more than 10 participants, this figure was increased to 36.

The average drop-out rate for parent training programmes is 28%.^{iv} Allowing for this drop-out rate, the sample size (reduced to 26) would still be sufficient to obtain a power of 90% to detect an effect of 2 effect sizes.

A similar procedure, allowing for an even more conservative estimate of the likely effect size to be obtained, gave a total sample size of 100 to allow detection of an effect size of 0.6 with 80% power or 120 to detect an effect size of 0.5 (moderate effect^v) with 80% power. This was chosen as the target sample size, which we estimated could be obtained from the three general practices initially selected for the survey.

References

1. Campbell SB. Behavior problems in preschool children: A review of recent research. *Journal of Child Psychology and Psychiatry* 1995; **36(1)**: 113-149.
2. Bone M, Meltzer H. *The prevalence of disability among children, OPCS Surveys of Disability in Great Britain, Report 3*. London: HMSO, 1989.
3. Government White Paper. *Saving Lives: Our Healthier Nation*, 1999.
4. Robins LN, Rutter M, (Eds). *Straight and devious pathways from childhood to adulthood*. Cambridge: Cambridge University Press, 1990.
5. McCord W, McCord J, Howard A. Familial correlates of aggression in nondelinquent male children. *Journal of Abnormal and Social Psychology* 1961; **62(1)**: 79-93.
6. Patterson GR, DeBaryshe BD, Ramsey E. A developmental perspective on antisocial behavior. *American Psychologist*, 1989; **44(2)**: 329-335.
7. Parker G. Parental 'affectionless control' as an antecedent to adult depression: a risk factor delineated. *Archives of General Psychiatry* 1983; **40**: 956-960.

8. Oakley-Browne MA, Joyce PR, Wells JE, Bushnell JA, Hornblow AR. Adverse parenting and other childhood experience as risk factors for depression in women aged 18-44 years. *Journal of Affective Disorders* 1995; **34**: 13-23.

9. Baumrind D. Parenting styles and adolescent development. In R Learner, AC Petersen, J Brooks-Gunn (Eds). *The encyclopaedia on adolescence* (pp. 746-758). New York: Garland, 1991.

10. Scott S, Knapp M, Henderson J, Maughan. Financial cost of social exclusion: follow up study of antisocial children into adulthood. *BMJ* 2001; 323: 191-194.

11. Hoghugh M. The importance of parenting in child health. *BMJ* 1998; 316: 1545.

12. Webster-Stratton C. Videotape modelling: A method of parent education. *Journal of Clinical Child Psychology* 1981; **10**: 93-98.

13. Henricson C, Katz I, Mesie J, Sandison M, Tunstill J. *National mapping of family services in England and Wales – A consultation document*. National Family and Parenting Institute, 2001.

14. Wolfendale S, Enzig H. *Parenting education and support*. London: David Fulton Publishers, 1999.

15. Barlow J, Stewart-Brown S. Behaviour problems and group based parenting education programmes. *Developmental and Behavioral Pediatrics*. 2000; **21**: 356-370.

16. Serketic WJ, Dumas JE. The effectiveness of behavioral parent training to modify anti-social behavior in children: A meta-analysis. *Behavior Therapy* 1996; **27**: 171-186.

17. Lister-Sharp D, Chapman S, Stewart-Brown S, Sowden A. Health promoting schools and health promotion in schools: two systematic reviews. *Health Technology Assessment* 1999; **3**(22).

18. Marshall J, Watt P. *Child Behaviour Problems*. Perth: Interagency Committee on Children's Futures, 1999.

19. Barlow J. (In press) Individual and group-based parenting programmes for improving psychosocial outcomes for teenage parents and their children. *Cochrane protocol*. Issue 3, Oxford: Update Software. In: The Cochrane Library. 2000

20. Scott S, Spender Q, Doolan M, Jacobs B, Aspland H. Multicentre controlled trial of parenting groups for childhood antisocial behaviour in clinical practice. *BMJ* 2001; 323: 194-197.

21. Offord DR, Kraemer HC, Kazdin AE, Jensen PS, Harrington R. Lowering the burden of suffering from child psychiatric disorder: Trade-offs among clinical, targeted, and universal interventions. *Journal of the American Academy of Child and Adolescent Psychiatry* 1998; **37 (7)**: 686-694.

22. Rose G. *The Strategy of Preventive Medicine*. Oxford: Oxford University Press, 1992.

**23. Mental Health Foundation,
www.mentalhealth.org.uk**

24. Stewart-Brown S. The public health importance of child behaviour problems. In: Buchanan A, Hudson B. *Parenting, Schooling and Children's Behaviour*. Guildford: Ashgate, 1998.

25. Leach P. *The physical punishment of children: Some input from recent research*. NSPCC Policy Practice Research Series, 1999.

26. Nobes G, Smith M. Physical punishment of children in two parent families. *Journal of Clinical Child Psychology and Psychiatry* 1997; **2**: 271-281.

27. Webster-Stratton C. Commentary: nipping conduct problems in the bud. *BMJ* 2001; 323: 197-198.

28. Webster-Stratton C, Hollinsworth T, Kolpacoff M. The long-term effectiveness and clinical significance of three cost-effective training programs for families with conduct problem children. *Journal of Consulting and Clinical Psychology* 1989; **57(4)**: 550-553.

29. Robinson EA, Eyberg SM, Ross AW. The standardization of an inventory of child conduct problem behaviors. *Journal of Clinical Child Psychology*. 1980; **9(1)**: 22-29.

30. Cohen J. *Statistical power analysis for the behavioural sciences*. New York: Academic Press, 1977.

31. Altman DG, Bland JM. How to randomise. *British Medical Journal* 1999; **319**:703-704.

32. Goodman R. A modified version of the Rutter parent questionnaire including extra items on children's strengths: A research note. *Journal of Child Psychology and Psychiatry*, 1994; **35(8)**: 1483-1494.

33. Goldberg DP, Hillier VF. A scaled version of the General Health Questionnaire. *Psychological Medicine*. 1979; **9(1)**: 139-145.

34. Abidin R. Parenting Stress Index: a measure of the parent-child system. In: Zalaquett C, Wood R (Eds). *Evaluating stress: A book of resources* (pp. 277-291). Lanham, MD, USA: Scarecrow Press, Inc., 1997.

35. Rosenberg M. *Conceiving the Self*. New York: Basic Books, 1979.

Patterson J, Stewart-Brown S, Mockford C, Barlow J, Pyper C. Need and demand for parenting programmes in a general practice setting. [Companion paper 1]

Patterson J, Mockford C, Barlow J, Stewart-Brown S. The impact of parenting programmes: A Qualitative study. [Forthcoming paper]

ⁱ Eyberg SM, Boggs SR, Algina J. Parent-child interaction therapy: A psychosocial model for the treatment of young children with conduct problem behavior and their families. *Psychopharmacology Bulletin* 1995; 31: 83-91.

ⁱⁱ Kazis LE, Anderson JJ, Meenan RF. Effect sizes for interpreting changes in health status. *Medical Care* 1989; 27 (3) Supplement: S178-S189.

ⁱⁱⁱ Computer program. *Power and Precision*. Release 1.20 September 24, 1997.

^{iv} Forehand R, Middlebrook J, Rogers T, Steffe M. Dropping out of parent training. *Behaviour Research and Therapy* 1983; 21(6): 663-668.

^v Cohen J. *Statistical power analysis for the behavioural sciences*. 1977, New York: Academic Press.