ORIGINAL ARTICLE

Type 2 and other forms of diabetes in 0–30 year olds: a hospital based study in Leeds, UK

R G Feltbower, P A McKinney, F M Campbell, C R Stephenson, H J Bodansky

Arch Dis Child 2003;88:676–679

See end of article for
authors' affiliations

Correspondence to:
Dr P A McKinney, Unit of
Epidemiology and Health
Services Research,
University of Leeds,
30 Hyde Terrace, Leeds
LS2 9LN, UK;
p.a.mckinney@leeds.ac.uk

Accepted
9 December 2002

Background and Aims: Following recent reports of increased numbers of adolescents being diagnosed with the adult or type 2 form of diabetes we aimed to describe the prevalence of both type 2 and other forms of diabetes in an urban population of children and young people in northern England.

Methods: A hospital based cross sectional study was performed in patients aged ≤ 30 years attending diabetic clinics in Leeds during the year 2000.

Results: A total of 677 subjects were identified, of whom 621 (92%) and 37 (5%) had type 1 and type 2 diabetes respectively. Four patients had confirmed maturity onset diabetes of the young, while the cause was uncertain for four. Median age of all patients was 22 years, with 396 (58%) aged 20–30; 32/37 patients with type 2 diabetes were aged 20–30. The prevalence of type 2 diabetes was 0.13 per 1000 overall, compared to 2.2 per 1000 for patients with type 1 diabetes. Of all type 2 diabetes patients, 24% were south Asian compared to 5% of the background population; 87% were categorised into the two least affluent tertiles of the Townsend score. This link with deprivation was not explained by the proportion of Asian patients across tertiles (approximately 25%).

Conclusions: This study shows extremely low prevalence of type 2 diabetes in 10–19 year olds, but will provide a baseline for future comparisons. Overall, type 2 diabetes is seen more commonly in south Asians, and an association with deprivation is suggested.

There is a clear need to assess the quality and provision of care for children and young people with diabetes, given that this is a condition that has long term complications and on which the UK National Health Service (NHS) currently spends almost 10% of its annual budget.¹ The focus on young people is particularly crucial because of the clear evidence that improved diabetic control through the maintenance of blood sugar levels leads to fewer complications and lower morbidity and mortality.^{2–4}

In order to maintain equity in the provision of care the planning of resource allocation needs to be based on reasonable estimates, not only of the number of individuals representing the overall burden of disease, but also quantifying the likely use of NHS services. At local level good information systems and diabetes registers can facilitate the provision of this type of information.⁵

Leeds is the second largest health authority in England with a densely urbanised population of 725 000 living in a compact geographical area of 560 km². South Asians comprised 5% of the local population in the 1991 census. Diabetes services for those diagnosed aged 30 and under are based at four hospital sites which function as part of the Leeds Teaching Hospitals NHS Trust, one of the largest Trusts in England with a population of 300 000 aged 0–30 years.

In the light of reports suggesting an impending epidemic of adult or type 2 diabetes in our teenage population,^{6–9} we aim to determine the prevalence of all forms of diabetes, including type 2, in children and young adults attending the hospital clinics in a large UK city. Since the risk of type 2 diabetes is known to be higher in Asian adult populations,^{10–11} we wanted to determine whether or not this was the case for a younger population. Finally we wished to examine the influence of socioeconomic status on the prevalence of type 2 diabetes.

METHODS

Information was collected on all children and young people aged 30 years and under who attended a hospital diabetes

clinic within the city of Leeds during the year 2000. Patients were identified using diabetes clinic lists for the paediatric, adolescent, and adult clinics from the following hospitals: the General Infirmary at Leeds (LGI), Wharfedale General Hospital, St James's University Hospital (SJUH), and Seacroft Hospital. Within the health authority it is well established practice for all newly diagnosed children and young adults, irrespective of their type of diabetes, to be referred to a hospital clinic; the clinics have a policy of not discharging patients into primary care. The two major teaching hospitals (LGI and SJUH) have dedicated adolescent and young adult clinics to which patients in the study age group were referred. Data were collected by abstracting information from hospital records onto a customised form, designed by both paediatric and adult diabetologists, by a single experienced research assistant (CS). Items included both demographic details and clinical information, which were computerised using a customised database. The amount of clinical information that was systematically available for every patient was limited; partial recording in the notes of height and weight, for example, did not allow us to analyse body mass index or to investigate relations with obesity.

The diagnostic category of diabetes was determined from the hospital records and classified as type 1,¹² type 2,¹³ maturity onset diabetes of the young (MODY), impaired glucose tolerance, and other forms of diabetes. Patients with other forms of diabetes included those categorised as "uncertain type 1 or type 2", diabetes ascribed to a known cause (either secondary to pancreatotomy, haematoma, or lipodystrophy), insulin treated "J" type (West Indian) diabetes, and unclassified diabetes. This group was excluded from the analysis. Patients whose initial diagnosis was uncertain or MODY had their medical records examined by two clinicians (FC, HJB) who evaluated all the available information and reached a consensus diagnosis. Criteria for the diagnosis of MODY were (a) genetic testing and/or (b) clinical (that is, diagnosed under 25 years of age, five years off insulin treatment from diagnosis,

Table 1 Comparative prevalence of diabetes (per 1000 population) in Leeds by age group

Type 1 diabetes			Type 2 diabetes			Other* types of diabetes		
Age (y)	n	Prevalence†	Age (y)	n	Prevalence†	Age (y)	n	Prevalence†
0-4	12	0.29	0-4	0	0.00	0-4	0	0.00
5-9	52	1.11	5-9	0	0.00	5-9	1	0.02
10-14	92	1.97	10-14	1	0.02	10-14	0	0.00
0-14	156	1.16	0-14	1	0.01	0-14	0	0.00
15-19	111	2.38	15-19	4	0.09	15-19	8	0.17
20-24	156	2.47	20-24	9	0.14	20-24	2	0.03
25-30	198	4.41	25-30	23	0.51	25-30	8	0.18
15-30	465	3.01	15-30	36	0.23	15-30	18	0.12
0-30	621	2.15	0-30	37	0.13	0-30	19	0.07

*Includes "uncertain type 1 or type 2 diabetes" (n=1), maturity onset diabetes of the young (n=4), impaired glucose tolerance (n=4), secondary diabetes (n=6), insulin treated "J" type (West Indian) diabetes (n=1), and "other" unclassified forms of diabetes (n=3).

†Denominator data was 2000 mid-year population estimates for Leeds District.

and family history of a similar condition, usually with an autosomal dominant pattern).

A measure of deprivation, the Townsend score,¹⁴ was derived for each individual based on their address and validated postcode using information from the 1991 decennial census. This index of material deprivation is derived from the following four variables: unemployment, car ownership, housing tenure, and household overcrowding. Townsend scores for all postcodes in Leeds were ranked and then divided into tertiles. Identical cut offs were then used to divide the case data into tertiles, with a higher score representing a more deprived area. This areal based measure of socioeconomic status has been shown to be a fairly robust measure of an individual's social class.¹⁵

A computer program¹⁶ was used to assign ethnic group to patients by identifying names of south Asian ethnic origin. This is based on full name and followed by visual inspection from an expert. The program has been shown to be accurate (90% sensitive, 99% specific) for the assignment of ethnic group to individuals in this way,¹⁷ and has been applied in other studies of ethnicity and paediatric chronic disease.^{18, 19}

Data were extracted from the customised database into Microsoft Access for further validation and checking, and then analysed using STATA.²⁰ Prevalence was calculated for each age group using mid-year (2000) population estimates for the district of Leeds, which is co-terminus with Leeds Health Authority. Prevalence figures for south Asians and non-south Asians were derived using data from the 1991 Census. Exact 95% CI were calculated assuming prevalence followed a binomial distribution.

RESULTS

A total of 677 patients attending hospital clinics in Leeds during the year 2000 were identified. The majority (86%) were assigned to the care of six consultants at the two major Leeds hospitals (LGI and SJUH), two of whom were paediatricians. Of these patients, 231 individuals (40%) were designated as being under the care of a specialist paediatrician, with a median age of 12.5 years (quartiles 9.6, 15.9). Patients attending the smaller outlying clinics without specialist paediatric support were all over 15 years of age.

The majority of patients (396/677, 58%) were aged between 20-30 years with the median age of the whole group being 22.1 years (quartiles 15.5, 26.9). The median duration of diabetes was 7.0 years (quartiles 3.0, 12.5). The median age at diagnosis was 11.5 (quartiles 6.7, 17.2), with two thirds of patients diagnosed under 15 years.

Table 1 shows the numbers of clinic attendees according to their type of diabetes and the estimated prevalence of these conditions by age group. A total of 621/677 (92%) patients had type 1 diabetes and 37/677 (5%) had type 2 diabetes, while a further 19/677 (3%) had their type classified as "other". This included uncertain type 1 or type 2 (n = 1), MODY (n = 4, aged 18-21), impaired glucose tolerance (n = 4), secondary diabetes (n = 6), insulin treated "J" type (West Indian) diabetes (n = 1), and other unclassified forms of diabetes (n = 3).

The prevalence of diabetes increased steadily with age, with a peak in 25-30 year olds of 4.4 and 0.5 per 1000 for type 1 and type 2 diabetes respectively. Overall prevalence was 2.2 and 0.13 for type 1 and type 2 diabetes. Prevalence of type 2 diabetes was significantly higher for south Asians than non-south Asians, both for 0-30 year olds combined (0.63 versus 0.10 per 1000; 95% CI 0.29 to 1.2 versus 0.067 to 0.15), and for 15-30 year olds (1.4 versus 0.17 per 1000; 95% CI 0.65 to 2.7 versus 0.11 to 0.25).

Thirty three (4.9%) of the 677 hospital attendees were south Asians (95% CI 3.4 to 6.8%), which was identical to the proportion of south Asians aged 0-30 years identified in the 1991 census. However, this proportion increased to 24.3% (95% CI 11.8 to 41.2%, n = 9/37) when we restricted the dataset to cases with type 2 diabetes (table 2). The proportion of patients with type 2 diabetes of south Asian origin was higher (40%) in the 10-19 year olds compared with the 20-30 year olds (22%), although there was no difference in the sex ratios by age. The median duration of disease for type 2 patients was typically two years, of which 5/37 of the patients with type 2 diabetes were diagnosed under the age of 15 years and 12/37 were diagnosed under 20. There was little variation in the proportion of Asians with age at onset, apart from the fact that no Asian patients were diagnosed between the ages of 20 and 24 years.

Table 2 Distribution of cases of type II diabetes by age, sex, and ethnic group in Leeds

Age group	Males	Females	Total	Median duration (y)	Proportion of South Asians (%)
10-19	2	3	5	2.2	40
20-24	4	5	9	2.9	22
25-30	12	11	23	2.0	22
10-30	18	19	37	2.1	24

Figure 1 Comparison of the socioeconomic profile (Townsend score) of patients with diabetes and the population of Leeds.

Figure 2 Distribution of patients with type 2 diabetes across Townsend deprivation tertiles by ethnic group.

Figure 1 shows that the distribution of the Townsend score for the diabetic patients is similar to the socioeconomic profile of the population of Leeds (Mann-Whitney test, $p = 0.73$), with a median score of -0.4 (quartiles $-0.4, 3.5$) for all hospital attendees. However, the profile of patients with type 2 diabetes was generally less affluent, with a median score of 4.3 (quartiles $-0.6, 6.9$). Figure 2 also shows the socioeconomic profile of patients with type 2 diabetes by ethnic group. This shows the highest numbers of type 2 patients occur in the two least affluent groups, an effect which was independent of ethnic group (Mann-Whitney test, $p = 0.99$).

DISCUSSION

This study provides timely prevalence figures of diabetes occurring in young people aged 0–30 in the second largest metropolitan district in England. These data are an important requirement for health service planning. The novel findings for type 2 diabetes are presented in the context of a dearth of published baseline data in the UK, and are particularly important in light of suggestions of an impending epidemic in

young people.^{7–21} Our data indicate that at present we are not facing a major public health problem with type 2 diabetes.

The prevalence of type 2 diabetes for young adults aged 15–30 in Leeds was estimated to be 0.23 per 1000 and 0.05 for those aged 10–19. In Birmingham, UK the estimated prevalence for under 18s was similar in 2000 at 0.038, showing this remains a rare chronic condition in this age range. Our data show clear differences in prevalence by ethnic group, with higher levels in south Asians aged 15–30 compared with non-south Asians (1.4 v 0.17, respectively) which taken together with the observations of Ehtisham and colleagues,²² show a consistent pattern of increased occurrence in young south Asians. This has clear implications for public health as we have now identified a high risk group who could be targeted for screening. Higher prevalence in other ethnic groups has been reported in the USA, ranging from 1 per 1000 in an American-Indian population²³ to 50 per 1000 in Pima Indians aged 15–19 years.²⁴

In considering whether our prevalence estimates represent the true burden of disease, a number of factors must be taken into account. A key issue is that diagnosis of type 2 diabetes is more likely to be misclassified, undiagnosed, or unreported than type 1.⁶ Our experience shows that clinicians treat diabetes empirically so that the aetiology is not always pursued. For example, few patients with type 2 diabetes had undergone testing for autoimmune antibodies in our study. Consequently, our classification of diabetes is based principally on clinical expert opinion, resulting in a few subjects for whom the diagnosis cannot be accurately defined. Despite this potential for misclassification, our results reflect what clinicians diagnose and treat and are therefore relevant to current healthcare in the city and can be extrapolated to other UK centres.

Another limitation is that our estimates may have excluded some eligible patients. This includes those who have never attended hospital, either because they were diagnosed and treated solely by general practitioners or they have failed to attend any appointments during the study period. In Leeds, the dominant patterns of referral from primary care are that virtually all paediatric patients are referred to hospital,²⁵ and

in the older age groups there is a well established tradition of referral to secondary care. Of 130 general practices in Leeds during 2000, only 47% reported having a clinic designed for adult diabetes patients. It is highly likely, particularly for teenagers, that they were referred to specialist hospital centres with experience of treating adolescents and young adults. Whilst accepting that we are more likely to have underestimated the numbers of older and therefore type 2 patients, we are confident that the figures are broadly representative, although possibly conservative.

A new observation is the association between type 2 diabetes and deprivation, with patients more likely to come from less affluent areas. If a relation existed between social class and obesity, this might explain the association between deprivation and type 2 diabetes, but there is little recent evidence to suggest a strong link between these two factors in children and young adults.^{26 27}

Obesity and lack of exercise, known risk factors for type 2 diabetes, are both increasing in adolescents in western populations,²⁸⁻³⁰ and appear to be coincident with a rise in type 2 diabetes. Unfortunately we were unable to directly investigate these risk factors in our dataset. To facilitate future research on young adult patients, particularly where the diagnosis is unclear, efforts should be made within all treatment centres to collect current information on weight and height and ethnic background, in addition to regularly evaluating the classification and treatment.

Our findings fail to show an "epidemic" of type 2 diabetes in adolescents in Leeds, but monitoring of trends will continue. A higher prevalence of type 2 diabetes in south Asian teenagers and young adults is consistent with observations in adults. The increased risk of type 2 diabetes associated with less affluent socioeconomic groups is a new finding and will require confirmation in other populations. Future work will investigate this cohort in relation to diagnostic specificity and clinical outcomes, including diabetic control.

Authors' affiliations

R G Feltbower, P A McKinney, C R Stephenson, University of Leeds, UK
F M Campbell, H J Bodansky, Leeds Teaching Hospitals NHS Trust, UK

REFERENCES

- 1 Currie CJ, Kraus D, Morgan CL, *et al.* NHS acute sector expenditure for diabetes: the present, future, and excess in-patient cost of care. *Diabet Med* 1997;14:686-92.
- 2 The DCCT/EDIC Research Group. Retinopathy and nephropathy in patients with type 1 diabetes four years after intensive therapy. The Diabetes Control and Complications Trial. *N Engl J Med* 2000;342:381-9.
- 3 The DCCT/EDIC Research Group. Effect of intensive therapy on the microvascular complications of type 1 diabetes mellitus. *JAMA* 2002;287:2563-9.
- 4 Edge JA, Ford-Adams ME, Dunger DB. Causes of death in children with insulin dependent diabetes 1990-96. *Arch Dis Child* 1999;81:318-23.

- 5 Vaughan NJA, Benedetti MM. A review of European experience with aggregated diabetes databases in the delivery of quality care to establish a future vision of their structure and role. *Diabet Nutr Metab* 2001;14:86-7.
- 6 Fagot-Campagna A, Pettitt DJ, Engelgau MM, *et al.* Type 2 diabetes among North American children and adolescents: an epidemiologic review and a public health perspective. *J Pediatr* 2000;136:664-72.
- 7 Zimmet P, Alberti KGMM, Shaw J. Global and societal implications of the diabetes epidemic. *Nature* 2001;414:782-7.
- 8 American Diabetes Association. Type 2 diabetes in children and adolescents. *Diabetes Care* 2000;23:381-9.
- 9 Ehtisham S, Kirk J, McEvilly A, *et al.* Prevalence of type 2 diabetes in children in Birmingham. *BMJ* 2001;322:1428.
- 10 Mather HM, Keen H. The Southall Diabetes Survey: prevalence of known diabetes in Asians and Europeans. *BMJ* 1995;291:1081-4.
- 11 Simmons D, Williams DRR, Powell MJ. The Coventry Diabetes Study: prevalence of diabetes and impaired glucose tolerance in Europids and Asians. *Q J Med* 1991;81:1021-30.
- 12 Alberti KG, Zimmet P. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: Diagnosis and classification of diabetes mellitus. Provisional report of a WHO consultation. *Diabet Med* 1998;15:539-53.
- 13 American Diabetes Association. Type 2 diabetes in children and adolescents. *Diabetes Care* 2000;23:381-9.
- 14 Townsend P, Phillimore P, Beattie A. *Health and deprivation: inequality and the North*. London: Croom Helm, 1988.
- 15 Danesh J, Gault S, Semmence J, *et al.* Postcodes as useful markers of social class: population based study in 26000 British households. *BMJ* 1999;318:843-5.
- 16 Bradford Health Authority, and City of Bradford Metropolitan District Council. *NAM PECHAN: South Asian Names (Indian Sub-continent)*. A computer program to identify and analyse south Asian names, version 1.1. Bradford, City of Bradford Metropolitan District Council Computer Services Division, 1996.
- 17 Cummins C, Winter H, Cheng KK, *et al.* An assessment of the Nam Pehchan computer program for the identification of names of south Asian ethnic origin. *J Public Health Med* 1999;21:401-6.
- 18 McKinney PA, Feltbower RG, Parslow RC, *et al.* Survival from childhood cancer in Yorkshire UK: effect of ethnicity and deprivation. *Eur J Cancer* 1999;35:1816-23.
- 19 Feltbower RG, Bodansky HJ, McKinney PA, *et al.* Trends in the incidence of type 1 diabetes in south Asians and other children in Bradford, UK. *Diabet Med* 2002;19:162-6.
- 20 StataCorp. *Stata Statistical software, release 7.0*. College Station, Texas, 2001.
- 21 Rosenbloom AL, Joe JR, Young RS, *et al.* Emerging epidemic of type 2 diabetes in youth. *Diabetes Care* 1999;22:345-54.
- 22 Ehtisham S, Kirk J, McEvilly A, *et al.* Prevalence of type 2 diabetes in children in Birmingham. *BMJ* 2001;322:1428.
- 23 Rios Burrows N, Acton K, Geiss L, *et al.* Trends in diabetes prevalence among American Indian and Alaskan Native children, adolescents and young adults, 1991-1997. 11th Annual Indian Health Service Research Conference, Albuquerque, New Mexico, 1999:19.
- 24 Dabelea D, Hanson RL, Bennett PH, *et al.* Increasing prevalence of type 2 diabetes in American Indian children. *Diabetologia* 1998;41:904-10.
- 25 Staines A, Bodansky HJ, Lilley HE, *et al.* The epidemiology of childhood diabetes in the United Kingdom: the Yorkshire Regional Childhood Diabetes Register. *Diabetologia* 1993;36:1282-7.
- 26 Sobal J, Stunkard AJ. Socioeconomic status and obesity: a review of the literature. *Psychol Bull* 1989;105:260-75.
- 27 Wang Y. Cross-national comparison of childhood obesity: the epidemic and the relationship between obesity and socioeconomic status. *Int J Epidemiol* 2001;30:1129-36.
- 28 Troiano RP, Flegal KM, Kuczmarski RJ, *et al.* Overweight prevalence and trends for children and adolescents: The National Health and Nutrition Examination Surveys, 1963 to 1991. *Arch Pediatr Adolesc Med* 1995;149:1085-91.
- 29 Chinn S, Hughes JM, Rona RJ. Trends in growth and obesity in ethnic groups in Britain. *Arch Dis Child* 1998;78:513-17.
- 30 Chinn S, Rona RJ. Prevalence and trends in overweight and obesity in three cross sectional studies of British children, 1974-94. *BMJ* 2001;322:24-6.

normal, delayed, or slow colonic transit times. In contrast in adults with constipation and a normal colonic transit time significantly more symptoms of psychological distress were found than patients with constipation of the slow transit type.²⁰ An important difference between adults and children with constipation lies in the absence of encopresis or soiling complaints in adults. We suggest that this agonising characteristic is primarily responsible for the high scores on the CBCL.

All study patients were referred to our paediatric motility unit to receive biofeedback training for their complaints. Prior to these sessions anorectal manometry was performed in all three patient groups. Clearly, anorectal manometry has no place in the diagnostic work up in children with recurrent abdominal pain. Total behaviour scores were not different for children with normal or abnormal defecation dynamics. These findings are in accordance with the study of Loening-Baucke *et al*, which also failed to show differences in behaviour profiles between children able or unable to defecate balloons.¹⁵

In conclusion, patients with defecation disorders have more behavioural problems than the normative sample. Since these behaviour problems are mild and disappear after adequate treatment, patients should be treated in a paediatric setting. Referral to a mental health service might be useful in patients with social withdrawal, a low self esteem, and depressive behaviour due to their defecation disorder.

Manometric or colonic transit patterns do not correlate with behavioural problems.

Authors' affiliations

M A Benninga, W P Voskuijl, J A Taminiau, Department of Paediatric Gastroenterology & Nutrition, Academic Medical Center, Amsterdam, Netherlands

G W Akkerhuis, Department of Child Psychiatry, Academic Medical Center, Amsterdam, Netherlands

H A Büller, Department of Paediatrics, Sophia Children Hospital, Rotterdam, Netherlands

REFERENCES

- 1 **Loening-Baucke V**. Constipation in early childhood: patient characteristics, treatment, and long-term follow up. *Gut* 1993;**34**:1400-4.
- 2 **Levine MD**. Encopresis: its potentiation, evaluation, and alleviation. *Pediatr Clin North Am* 1982;**29**:315-30.
- 3 **Wasserman AL**, Whittington PF, Rivara FP. Psychogenic basis for abdominal pain in children and adolescents. *J Am Acad Child Adolesc Psychiatry* 1988;**27**:179-84.
- 4 **Levine MD**, Mazanson P, Bakow H. Behavioural symptom substitution in children cured of encopresis. *Am J Dis Child* 1980;**134**:663-7.
- 5 **Landman GB**, Rappaport L, Fenton T, *et al*. Locus of control and self-esteem in children with encopresis. *J Dev Behav Pediatr* 1986;**7**:111-13.
- 6 **Gabel S**, Hegedus AM, Wald A, *et al*. Prevalence of behavior problems and mental health utilization among encopretic children: implications for behavioral pediatrics. *J Dev Behav Pediatr* 1986;**7**:293-7.
- 7 **Friman PC**, Mathews JR, Finney JW, *et al*. Do encopretic children have clinically significant behaviour problems? *Pediatr Radiol* 1988;**82**:407-9.
- 8 **Apley J**, Naish N. Recurrent abdominal pains: a field survey of 1,000 school children. *Arch Dis Child* 1958;**33**:165-70.
- 9 **Achenbach TM**. *Integrative guide for the 1991 CBCL/4-18, YSR and TRF profiles*. Burlington, VT: University of Vermont Department of Psychiatry, 1991.
- 10 **Verhulst FC**, Akkerhuis GW, Althaus M. Mental health in Dutch children: (I). A cross-cultural comparison. *Acta Psychiatr Scand* 1985;**323**(suppl):1-108.
- 11 **Achenbach TM**. *Manual for the Child Behaviour Checklist and Revised Child Behaviour Profile*. Burlington, VT: University of Vermont, Department of Psychiatry, 1983.
- 12 **Metcalf AM**, Phillips SF, Zinsmeister AR, *et al*. Simplified assessment of segmental colonic transit. *Gastroenterology* 1987;**92**:40-7.
- 13 **Arhan P**, Devroede G, Jehannin B, *et al*. Segmental colonic transit time. *Dis Colon Rectum* 1981;**24**:625-9.
- 14 **Benninga MA**, Büller HA, Taminiau JA. Biofeedback training in chronic constipation. *Arch Dis Child* 1993;**68**:126-9.
- 15 **Loening-Baucke V**, Cruikshank B, Savage C. Defecation dynamics and behaviour profiles in encopretic children. *Pediatrics* 1987;**80**:672-9.
- 16 **Nolan T**, Debelles G, Oberklaid F, *et al*. Randomised trial of laxatives in treatment of childhood encopresis. *Lancet* 1991;**338**:523-7.
- 17 **Young MH**. Functional encopresis: symptom reduction and behavioural improvement. *J Dev Behav Pediatr* 1995;**16**:226-32.
- 18 **van der Plas RN**, Benninga MA, Redekop WK, *et al*. Randomised trial of biofeedback training for encopresis. *Arch Dis Child* 1996;**75**:367-74.
- 19 **Garber J**, Zeman J, Walker LS. Recurrent abdominal pain in children: psychiatric diagnoses and parental psychopathology. *J Am Acad Child Adolesc Psychiatry* 1990;**29**:648-56.
- 20 **Wald A**, Hinds JP, Caruana BJ. Psychological and physiological characteristics of patients with severe idiopathic constipation. *Gastroenterology* 1989;**97**:932-7.

CORRECTION

The authors of the paper by Feltbower *et al* in the August issue (Type 2 and other forms of diabetes in 0-30 year olds: a hospital-based study in Leeds, UK, *Arch Dis Child* 2003;**87**:679-9) would like to fully acknowledge the help and support of the following groups and individuals, including the Leeds Diabetes Patient Record System Group.

MEMBERSHIP OF THE LEEDS DIABETES PATIENT RECORD SYSTEM GROUP (1999-2001)

Patricia McKinney, Carolyn Stephenson, Richard Feltbower (Paediatric Epidemiology Group), Trude Silman (former Community Health Council), Sue Jones, Sue Dodman (Clinical Services Managers), Linda Clapman, Virginia Nicholson (Diabetes Specialist Nurses), Fiona Campbell (Consultant Paediatrician), H Jonathan

Bodansky (Consultant Diabetologist), Deborah Quant (Patient and IT consultant), Sally Casley, Gillian Richardson (Clinical Effectiveness), Joe Eastman (Health Economist), Judith Knapton (Primary Care), Debra Gill (IT systems).

ACKNOWLEDGEMENTS

We are grateful to Dr Jon Fear from Leeds Health Authority for providing financial support towards the collection and analysis of the data. Novo Nordisk are thanked for providing the Diabetes Patient Record System software. We also appreciate the help and assistance of all local clinicians (Dr P Holland, Dr P Sheridan, Dr MU Afza, Dr S Gilbey, Dr M Mansfield, Dr JK Wales, Professor PJ Grant) and their staff. Diabetes specialist nurses were especially supportive, as were secretarial and medical records staff throughout the Trust. We also thank the Office for National Statistics for providing mid-year population estimates. Finally, we thank Roger Parslow for helpful comments on the manuscript.