Material We realized a prospective study on 87 patients diagnosed with AID. We analyzed the initial immune status, for all these patients.

For another group of 53 children, diagnosed with selective IgA immunodeficiency, we realized a six year period survey of the level of the T suppressor lymphocytes, the T helper/T suppressor ratio, and of the presence of auto antibodies: Anti DNA, rheumatoid factor (RF).

Results 8 patients (8%), from the group diagnosed with AID, were also identified with selective deficit in IgA at the moment of the initial diagnosis.

In the group of 53 patients with underlying IgA immunodeficiency, 2 patients developed over the 6 years of the survey, a significative titer of anti DNA antibodies. In one patient the presence of the RF was detected, 4 children presented a decrease of the T suppressor level, with a rise of the immune ratio. None of these patients presented clinical signs suggesting an AID.

Conclusions The Ig A immunodeficiency may be a risk factor for subsequent AID. There is a higher risk for AID in patients who develop anti DNA antibodies, RF or a persistent decrease of the T suppressor lymphocytes.

489 INTERLEUKIN PHENOTYPE IN PATIENTS WITH PFAPA

doi:10.1136/archdischild-2012-302724.0489

¹SI Iurian, ¹L Dobrota, ²S Iurian, ¹ML Neamtu, ³L Bera, ⁴A Rosenberg. ¹Research Department, Pediatric Clinic, Lucian Blaga University; ²Clinical Laboratory, Pediatric Hospital; ³Lucian Blaga University; ⁴Pediatric Hospital, Sibiu, Romania

Background PFAPA is a chronic condition including recurrent fever episodes, aphthous stomatitis, pharyngitis, adenitis. According to a previous study, even between febrile attacks, there is increasing of pro-inflammatory mediators.

Aims

1. To evaluate serum interleukin phenotype between febrile episode in PFAPA patients from our clinic;
2. To establish correlation between C reactive protein (CRP) and pro-inflammatory interleukins: tumor necrosis factor-alpha (TNF α), interleukin-8 (IL-8);
3. To evaluate link between CRP and anti-inflammatory interleukins: interleukin-10 (IL-10);
4. To identify a sensitive biological marker to estimate PFAPA evolution.

Methods Authors analyzed 2 groups: "PFAPA group" represented by 6 patients and "control group" containing 4 no-PFAPA patients. Inclusion criteria: patients up to 10 years of age that fulfilled PFAPA diagnosis criteria; patients between febrile attacks; negative procalcitonin (PCT) blood value in order to exclude bacterial infections for study patients. Exclusion criteria: patients during febrile attacks. Both groups patients were tested for serum levels of PCT, CRP, IL-8, TNF α , IL-10. Data was statistically analyzed using independent "t" test.

Results Both group patients have normal serum levels for interleukins 8/10 and high TNF α values. Mean value for TNF α was 11.26 pg/ml in PFAPA group and 13.2 pg/ml in no-PFAPA group. Regarding CRP values, mean value for PFAPA patients was 19.72 (range between 2.4–95) as compare to 5.04 in no-PFAPA patients.

Conclusions TNF α , IL-8, IL-10 aren't useful to appreciate PFAPA evolution pattern. CRP remains a sensitive marker for disease activity in PFAPA patients, even out of fever attacks. Our study didn't confirm previous study data.

490 RISK FACTORS OF ASTHMA EXACERBATION IN CHILDREN

doi:10.1136/archdischild-2012-302724.0490

DT Qosaj, S Hasani, H Ymeri. Pulmonology, Univeristy Clinical Centre of Kosova, Prishtina, Kosovo

Introduction Asthma is chronic disease which in recent years prevalence shows increase rate. In this study we analyze triggers that parents or child will tell as a risk factor for asthma exacerbation in children with asthma:

Aim Presenting cases treated in pulmonology clinic during their exacerbation and triggers that can lead to asthma exacerbation.

Method It is a prospective study, we include 92 children who came with symptoms of asthma exacerbation in pulmonology clinic. Asthma was clasified according to GINA classification and evaluate from parent/caregiver or child about risk factors that lead to exacerbation. Factors were listed by GINA.

Results Asthma exacerbation symptoms at children that were examined are cough(100%),difficult breathing(96%) and wheezing(24%) and chest tightness. From all children in our study 34% had one trigger, 16% 2 triggers and the others had more than 3 trigers that lead to asthma exacerbation. Triggers of asthma exacerbation are changing weather(cold air) (36%), viral infection(48%), passive smoking (36%), polen(16.6%) and 25% of parents don't know risk factors that lead to asthma exacerbation. Most of the children in the study lives in town (75%).

Conclusion Our study shows that viral infection and cold air are very common triggers of asthma exacerbation especially in small children while in children older than 5 years passive smoking is very present (36%) as risk factor.

491 PARAMETERS OF IMMUNE SYSTEM DISREGULATION DURING HERPES SIMPLEX VIRUS INFECTION IN CHILDHOOD

doi:10.1136/archdischild-2012-302724.0491

¹SM Hristina, ²L Saranac, ¹B Kamenov, ¹T Stankovic, ³D Djuric, ⁴L Brankovic, ⁴R Milicevic. ¹Immunology, Allergy and Genetic; ²Endocrinology; ³Gastroenterology; ⁴Molecular Medicine-Laboratory, Clinic of Pediatric, Medical University, Nis, Serbia

Introduction An important characteristic of the herpes simplex viruse, is their ability to persist in the tissues of their hosts for many years after initial infection as intracellular viruses. Characteristic life of virus (hronic persistent and ciclic replication) in organisms is often followed by immune dysregulation.

Materials and methods: Clinically manifestations in patients with herpesvirus infections were examined. We analysed: white blood cell count, hemoglobin level, serum immunoglobulins level, enzymes of cell destruction, oxidative metabolism of the peripheral blood phagocytes as ability of NBT reduction, serum level of IFN- γ IL-4 and DHEAS, cortisol were mesured by ELISA test.

Results Our patients had and all of them had positive ELISA test on viruse-HSV. Our parameters approved low level of hemoglobin, monocytosis, lymphocytosis, virocytosis and leukopenia. Our patients had high level LDH, CPK, low ability of NBT reduction. High levels of IFN γ followed high levels of LDH, CPK, GOT and GPT.

Conclusion Chronic activation of immune system is background of patogenetic mechanisms during herpes simplex virus infection. Different level of DHEAS and cortisol are part of regulatory mechanisms of immune response across endocrine system. Increase levels of DHEAS in our patients can display chronic inflammation. Absence of increase level of cortisol may suggestion that our patients had a little "acute" fase of infection opposite a lot of chronic disorders. Analyse of immunoregulatory mechanisms is essential to order level and place of damage cells, tissue and organs. It is important for therapy and prognosis of disease.

492 ATOPIC DERMATITIS ASSOCIATED WITH OBSTRUCTIVE BRONCHITIS IN EARLY CHILDHOOD

doi:10.1136/archdischild-2012-302724.0492

¹I Timovski, ¹V Angelovska, ²M Timovska. ¹PHI; ²PRD, Skopje, FYR Macedonia