were normal. Abdominal ultrasound unremarkable. Heart ultrasound: unremarkable. Neurological examination, muscle strength and reflexes are normal. EMG: no miopathic changes. ENG: normal. Rtg of lower extremities found calcifications of soft tissues. Bones with no destruction or osteolyses. Biopsy: on subcutan adipose tissue are found multiple and confluent foci of calcification called Calcinosi. She is without treatment to now and exacerbation of changes has been detected.

Conclusion Since there were no systemic metabolic disorders, tissue injury, or other founded reason, and the diagnose of Calcinosi cutis idiopathica was made.

Acknowledgements to nurses of nephrology department.

630 NEUROPSYCHOLOGICAL OUTCOME IN CONGENITAL HYPOTHYROIDISM IN AN ITALIAN COHORT: THE DEVELOPMENTAL QUESTIONNAIRE FOR CH (DQCH)

doi:10.1136/archdischild-2012-302724.0630

¹S Bargagna, ²A Olivieri, ³C Fazzini, ¹M Bozza. ¹Stella Maris Scientific Institute, Pisa; ²Istituto Superiore Sanità, Rome, Italy

Background and Aims Several studies on neuropsychological outcome in early treated children with congenital hypothyroidism (CH). have demonstrated the presence of developmental disorders, in particular motor abilities language, emotions 1.2.3 The aim of this study was to individuate the most vulnerable developmental domains and which risk factors are significantly associated with a poor outcome. To this end we have developed a developmental questionnaire on CH (DQCH).

Methods The DQCH has been created with dichotomous answers giving a score which hinders the dispersion of data and makes it easy to fill-in for a person knowing the child (clinical psychologist, physician, parents). It consists of 49 questions, in 7 domains - motor skills, personal autonomy, language development, social development, behaviour, biorhythms, and success in school. This questionnaire was designed for an easy way to collect data on developmental milestones and neuropsychological outcome in a large cohort of children with CH and their age-matched controls.

Results All domains of our questionnaire show major impairments in children with CH than in controls, in particular in personal autonomy for the group with thyroidal agenesis and social development for ectopic glands.

Conclusions We hypothesize that children with a more severe outcome are more frequently those of mothers affected by clinical and subclinical forms of hypothyroidism during pregnancy.

References Oerbeck B, et al. Congenital Hypothyroidism: Influence of Disease Severity and L-Thyroxine Treatment on Intellectual, Motor, and School-Associated outcomes in Young Adults. *Pediatrics* 2003; 4:923-930.

Rovet JF. Congenital Hypothyroidism: long-term outcome. *Thyroid* 1999; Neuropsychological Developmental Congenital C Child Neuropsychological 2002.

631 INFLUENCE OF ANTHROPOMETRICAL AND MENTAL STATUSES ON OCCURRENCE OF BEHAVIOUR DISEASES AMONG CHILDREN OF SCHOOL AGE

doi:10.1136/archdischild-2012-302724.0631

K Lepiatsila. *Belarussian State Medical University, Minsk, Belarus*

To estimate influence of the anthropometrical status and personal features on occurrence of psychosomatic frustration, in particular, eating disorder, among children of school age.

We studied 57 children (age 15±1.3 years). They were interviewed with Toronto Alexithymia Scale (TAS), scale of Rotter, standardized questionnaire "Aim-Means-Result" (AMR), scale of Rean and divided

into 3 groups on the basis of body mass index (BMI). 1) A (BMI< 19 kg/m², 2) B(19 kg/m²< BMI< 24kg/m²), 3)C (BMI>24kg/m²).

Results Based on the TAS 41% of boys showed higher than normal rates of alexithymia (A -40%,B -60%,C -100%). Among girls (75%,50%,43% respectively). AMR: 87%b and 82%g can't substantiate their goals. 50% of children had psychological barriers. Low-energy potential was recorded in Groups A and C (17% and 25%). 66% girls and 88% boys of all groups indicated optimal results. Scale of Rotter. Boys of Group C were dominated by external locus of control. Girls of such group were dominated by internal locus of control of life. Motivational pole is not strongly marked in 61% of girls.

The analysis of TAS showed an increasing trend of alexithymia among boys (from 59% to 100%) and lower among girls (from 75% to 43%) with an increase in BMI. Boys with a BMI> 24 kg/m² prevailed external locus of control, the girls - internal locus of control. In analyzing the results of the questionnaire of Rean boys tended to reduce the motivation to succeed and grow with the increase of negative motivation with increasing BMI (success from 70% to 25%).

632 EARLY INTERVENTION OF VISUAL IMPAIRMENT MAY PROTECT ADAPTIVE BEHAVIOUR IN DOWN SYNDROME?

doi:10.1136/archdischild-2012-302724.0632

¹A Dressler, ²M Bozza, ³V Perelli, ³S Bargagna. ¹Division of General Neuropediatrics and Neonatology, Medical University Vienna, Vienna, Austria; ²IRCCS-Stella Maris Institute; ³Stella Maris Institute, Pisa, Italy

Down syndrome (DS) is one the most common genetic disorders. Little is known about the impact of visual disorders in DS on daily life. Our aim was to study the relation between the incidence of ocular manifestations and adaptive behaviour.

Methods We performed a detailed medical history, including ocular disorders. We tested Hirschberg's corneal reflex method, observed eye movements during fixation of a slowly moving object, cover test, and assessed the Vineland Adaptive Behaviour Scales, as well as the Coloured Progressive Matrices. 52 DS individuals with an age range of 19 to 52 years. Results. We observed a high incidence of ocular anomalies (refractive errors in 59.2%, strabismus in 51%, motility disorders in 63.3% and congenital bilateral cataract in 16.2%), not differing with age. The occurrence of visual disorders did neither lead to total lower adaptive level nor show an influence on cognition. Daily living skills were significantly lower in individuals with not corrected disorders, but these individuals showed a significant high co-morbidity of autism with childhood onset. We observed a majority of individuals with an adaptive functioning above the average. Hyperopia and the presence of refractive errors were significantly more frequent in individuals with an adaptive level above the average. Conclusions. Visual disorders lead to a poorer performance in adaptive behaviour in individuals with DS and also autism. This suggests that early intervention on refractive errors and visuo-motor skills helps in the acquisition of daily living activities, which remain stable over the life-span. Therapeutic options for cataract need to be addressed early.

633 WHAT IS THE EFFECTIVENESS OF DIMETHYLGLYCINE IN TREATING AUTISTIC SYMPTOMS IN CHILDREN: A SYSTEMATIC REVIEW

doi:10.1136/archdischild-2012-302724.0633

M Haroon, G Ryan, M Randell, J Wilson, T Khatau, CHARGE: Child Health Applied Research Group, East-midlands. *Leicester Partnership Trust, Leicester, UK*

Background and Aims Dimethylglycine, an amino-acid derivative, has been proposed as a treatment for Autism. A systematic review was conducted to synthesize the best evidence relating to this topic.