to both aEEG pattern at 48h and MRI performed during the second week. Comparisons were done with ANOVA, SPSS19.

Results 21 newborns were monitored. During hypothermia there was a trend towards higher rSO₂ in newborns with predicted abnormal outcome. During rewarming there was a significant difference between the normal and abnormal outcome groups ($p<0.02$).

Abstract 1069 Table 1 rSO₂ values during hypothermia and rewarming

	rSO ₂	Normal (n=9)	Intermediate (n=4)	Abnormal (n=8)
Hypothermia	12 hours	76,3	74,5	79,4
	24 hours	83,1	77,0	88,1
	48 hours	84,6	88,0	89,5
	72 hours	82,1	79,0	87,4
Rewarming	34.5°C	77,3	84,0	89,8
	35.5°C	75,8	73,5	88,4
	35.5°C	74,6	77,8	87,4

Conclusions NIRS monitoring may improve the early recognition of newborns with abnormal outcome. Our data suggests that a rSO₂ value of 85 during rewarming can be used as a cutoff for predicting abnormal outcome.

1070 CEREBRAL BLOOD FLOW CHANGES IN PRETERM NEONATES REQUIRING RESUSCITATION AT BIRTH

doi:10.1136/archdischild-2012-302724.1070

¹N Ahmed, ²V Sundaram, ²P Kumar. ¹Neonatology; ²PGIMER, Chandigarh, India

Background Hypoxia in the perinatal period causes increase in cerebral blood flow followed by a decrease. CBF data is not available for preterms.

Objectives To study CBF changes in Anterior and Middle cerebral arteries (ACA & MCA) of asphyxiated preterm neonates (< 34 weeks).

Setting Level III Neonatal unit in Northern India, PGIMER, Chandigarh.

Subject and Interventions This is a subgroup analysis of a RCT comparing 21% vs 100 % oxygen for resuscitation. CBF was measured in ACA and MCA at 6±2, 12±2 and 24±4 hours. Peak systolic velocity (PSV) cm/s, End diastolic velocity (EDV) cm/s, Resistance index (RI), Pulsatility index (PI), Systolic diastolic ratio (S/D) and Velocity time integral (VTI) were measured using Philips ultrasonography machine.

Results CBF was measured in 57 neonates. Mean gestation and weight was 31±1.3 weeks and 1400±358 g. HIE was seen in 24 babies (42%). PSV and VTI showed an increasing trend whereas S/D showed a decreasing trend as age increased. PSV [29.2 (23.7–34.2), 28.3 (23.9–33.4) and 33.9 (29.6–37.8) cm/s, $p=0.02$], VTI (6, 6.5 and 8 cm, $p=0.002$) and S/D (4.1 (3.07–6.6), 4.03 (3.3–5.4) and 3.8 (3.1–4.7), $p=0.01$) in the MCA showed significant trend at 6, 12 and 24 hrs respectively. No such trend could be observed in the ACA. On comparing 21% and 100% oxygen groups no difference were observed.

Conclusion CBF indices (PSV, VTI & S/D) of MCA showed a significant increase from 6 to 24 hrs. No difference was observed on comparing 21% with 100% oxygen.

1071 AEEG DURING THE FIRST 72 HOURS AFTER BIRTH IN INFANTS WITH PRENATALLY DIAGNOSED CONGENITAL HEART DEFECT

doi:10.1136/archdischild-2012-302724.1071

¹H ter Horst, ¹G Benus, ²M Rooftoof, ¹E Kooi, ¹A Bos. ¹Pediatrics; ²Pediatric Cardiology, University Medical Center Groningen, Groningen, The Netherlands

Background and Objective Survival in infants with congenital heart defects (CHD) is associated with neurologic morbidity. Hypoxic-ischemia after closure of the ductus arteriosus is probably one of the factors resulting in brain damage. Prenatal diagnosis makes it possible to prevent closure of the ductus arteriosus and therefore cerebral perfusion can be preserved. Amplitude integrated EEG (aEEG) is a method to evaluate brain injury. Abnormal (a)EEG patterns and epileptic activity (EA) are frequently observed in infants with CHD. Our aim was to determine the course of aEEG patterns in infants with prenatally diagnosed CHD.

Methods Retrospective cohort study of infants prenatally diagnosed with CHD. aEEGs were assessed by pattern recognition: background pattern (BP), presence of sleep wake cycling (SWC) and EA were appraised.

Results Twenty infants (mean GA 39 wks; birth weight 3416 g) were included. Eleven infants had transposition of the great arteries, 7 infants had hypoplastic left heart syndrome, and 2 had aortic valve stenosis. At 6h after birth 80% of infants had normal BPs (continuous normal voltage (CNV)). Only 2 infants had severely abnormal BPs (continuous low voltage). At 24h, 94% had CNV. None of the infants showed EA. SWC was present in 85% and emerged at a median postnatal age of 10.4 hrs.

Conclusions aEEGs of infants with a prenatally diagnosed CHD are normal in the majority of infants, with normal emergence of SWC and absence of EA. This indicates that prenatal diagnosis can prevent brain damage in infants with CHD.

1072 REVIEW OF CURRENT OPINION AND AVAILABILITY OF CEREBRAL FUNCTION MONITORING AND THERAPEUTIC HYPOTHERMIA IN UK NEONATAL UNITS

doi:10.1136/archdischild-2012-302724.1072

¹CS Zipitis, ¹A Sylvan, ²T Harte, ¹S Hurst, ¹K Glover. ¹Royal Albert Edward Infirmary, Wigan; ²Salford Royal Hospital NHS Foundation Trust, Salford, UK

Background and Aims Therapeutic hypothermia has been established as an effective treatment modality for term neonates that have suffered an acute perinatal hypoxic ischaemic event. This is a time-critical procedure which increases pressure to commence active cooling as soon as possible after the injury.

This study aims to define the current levels of ownership of CFM and cooling equipment and capture some staff perceptions around cooling matters in neonatal units in the UK.

Methods Telephone survey conducted over March–November 2011 with the 205 active neonatal units listed on the British Association of Perinatal Medicine website.

Results 100% response-rate (205/205). 141 (68.8%) responders were Lead Nurse/Sisters, 59 (28.8%) were Nurses and 5 (2.4%) were Doctors. Overall, CFM was available at 106/205 (51.7%) units and 89/205 (43.4%) had cooling equipment. Equipment ownership was high in NICUs and progressively decreased in LNUs and SCBUs. The majority of responders were positive to the idea of therapeutic hypothermia as a standard of care whilst some asked for further research; only a small minority were negative. 91.6% were keen to consider a scheme where therapeutic hypothermia was initiated at the referring centre and continued in a dedicated cooling centre.

Conclusions Therapeutic hypothermia is widely recognised as effective. Availability of cooling equipment and CFM has increased in all levels of units in the UK. The time-constraints in initiating therapeutic hypothermia might mean that a scheme of initiating therapeutic hypothermia locally, continuing this during transport and completing it in a designated cooling centre is the way forward.

1073 EARLY EEG IN INFANTS WITH HYPOXIC ISCAEMIC ENCEPHALOPATHY: POSSIBLE INFLUENCES OF THERAPEUTIC HYPOTHERMIA

doi:10.1136/archdischild-2012-302724.1073