Methods 2 cases of PTPN11 gene mutation confirmed Noonan syndrome selected for review based on clinical course.

Results

1. Male born at 35+3 weeks with antenatal diagnosis of bilateral pleural effusion. Transferred to The Children's University Hospital on day 10 for management of malrotation; echo revealed structurally normal heart with mild pulmonary hypertension. Day 24 monocytosis and splenomegaly noted. Day 25 echo demonstrated increasing left ventricular hypertrophy (LVH) with normal function. Day 27 diagnosed with Juvenile Myelomonocytic Leukaemia and commenced on treatment with methylprednisolone. Day 32 repeat echo showed severe LVH with near obliteration of the left ventricle. Rate of acceleration queried to be secondary to glucocorticoids. Patient died day 32 secondary to multisystem organ failure.
2. Male born at 37+6 weeks with antenatal diagnosis of right side pleural effusion. Day 1 profound hypotension resistant to multiple inotrope support, chest drain inserted and commenced on inotrope resistant hypotensive dose of hydrocortisone. Echo day 1 moderate biventricular hypertrophy and structurally normal heart. Day 15 echo demonstrated severe left ventricular hypertrophy with significant cardiac compromise. Despite maximum efforts continued to deteriorate and died on day 17.

Conclusions Noonan syndrome is an uncommon condition with an association of hypertrophic cardiomyopathy in 20% to 30% of patients. In this case series complications of Noonan syndrome treated with glucocorticoids may have exacerbated cardiac function to an irreversible degree. This should be considered in the management of these patients.

1158 SYSTOLIC-DIASTOLIC FUNCTION IN CONGESTIVE HEART FAILURE SECONDARY TO CONGENITAL HEART MALFORMATIONS EVALUATED BY CLASSICAL AND TISSUE DOPPLER ECHOCARDIOGRAPHY

doi:10.1136/archdischild-2012-302724.1158

¹A Butnariu, ¹C Rusu, ¹D Iacob, ²C Munteanu, ¹M Marc. ¹University of Medicine and Pharmacy 'Iuliu Hatieganu'; ²Public Health Institute Iuliu Moldovan, Cluj-Napoca, Romania

Background and Aims Tissue Doppler velocities are relatively independent of ventricular geometry, particularly right ventricular geometry that is predominantly affected in the case of congenital heart disease (CHD).

Aims The evaluation of systolic and diastolic function in pediatric patients with congestive heart failure (CHF) secondary to CHD using classical echocardiographic parameters and pulsed tissue Doppler parameters.

Methods The study included 27 children diagnosed with CHF secondary to congenital heart malformations. The parameters of systolic and diastolic function were measured by 2D echocardiography, 2D guided M mode, color and pulsed Doppler, as well as by pulsed tissue Doppler at the level of the mitral and tricuspid annulus.

Results A relaxation alteration pattern or a pseudonormal pattern of E diastolic velocity compared to the A wave was found ($E=A$; $E>A$) in the group of subjects with heart failure. E wave deceleration time (EDT) had significantly increased values in the case of patients with CHF, being correlated with diastolic dysfunction. Left ventricular flow propagation velocity V_p was decreased in patients with heart failure. Associations between the severity of systolic dysfunction and the diastolic dysfunction evaluated by 2D echocardiographic parameters, M mode and Doppler and measured by pulsed tissue Doppler velocities at the mitral and tricuspid annulus were found in pediatric patients diagnosed with congestive heart failure ($p<0.05$).

Conclusions In children with heart failure, some conventional parameters of the diastolic function were maintained within

normal or pseudonormal values, diastolic dysfunction being confirmed in these cases by tissue Doppler measurements.

1159 LONG TERM GROWTH OF EXTREMELY LOW BIRTH WEIGHT INDIAN INFANTS AT CORRECTED AGE 1 YEAR

doi:10.1136/archdischild-2012-302724.1159

K Mukhopadhyay, R Mahajan, P Kumar. Neonatal Unit, Dept of Pediatrics, PGIMER, Chandigarh, India

Background and Aims Inspite of increasing survival of ELBW infants in India there is significant paucity of data on their long term growth. The aim of our study was to assess growth of ELBW infants at corrected age 1 year.

Methods Prospective observational study conducted in the follow up clinic of a level III neonatal unit. Forty eight ELBW infants discharged from neonatal intensive care unit were followed for weight, length and Head circumference at corrected age (CA) 1 year and z-scores were calculated.

Results The mean (SD) birth weight and gestation were 872 ± 82 g and 29.9 ± 2.3 weeks. At CA 1 year, as per WHO growth charts (2006), growth at $<3^{\text{rd}}$ centile was observed in 60.4% infants for weight, 54% for length and 58.3% for HC. Growth was more than 50th centile in 6.3% infants for weight, 4.2% for length and 2.1% for HC. Z-scores of 0 and above was recorded in 9% infants for weight and 7% for length.

Twenty eight babies (58%) were SGA. SGA infants were smaller in weight (856 ± 87 g vs 901 ± 66 g, $p=0.023$), higher in gestation (30.9 ± 1.9 wks vs 28.2 ± 1.8 wks, $p=0.000$) and discharged earlier ($p=0.003$) than AGA. There was no difference in growth parameters between SGA and AGA and male and female at CA 1 year.

Conclusion ELBW infants had significant growth failure at CA 1 year and intergroup differences between SGA and AGA were not observed.

1160 EFFECT OF CERVICAL COLONIZATION ON NEONATAL OUTCOME IN HIGH RISK PREGNANCIES: RESULTS FROM A TERTIARY MATERNITY CENTER IN TURKEY

doi:10.1136/archdischild-2012-302724.1160

¹MY Oncel, ²S Celen, ³G Demirel, ¹FE Canpolat, ¹E Calisici, ¹R Ozdemir, ¹SS Oguz, ³S Saygan, ²N Danisman, ¹U Dilmenc. ¹Neonatology; ²Perinatology; ³Microbiology, Zekai Tahir Burak Maternity Teaching Hospital; ⁴Pediatrics, Yildirim Beyazit University, Faculty of Medicine, Ankara, Turkey

Background and aim To evaluate and compare the morbidity and mortality of neonates born to pregnant women with positive and negative cervical cultures.

Methods The demographic and clinical features of mothers included in this study, along with details of the microorganisms isolated on maternal cervical cultures and the number of days between a positive cervical culture and delivery were recorded. Neonates were stratified into two groups based on cervical culture results of their mothers - Group 1, positive cervical culture; Group 2, negative cervical culture.

Results A total of 216 women who delivered 242 infants were included in the study. Group 1 consisted of 90 neonates while Group 2 had 152 newborns. Mean levels of the acute phase reactants, CRP and IL-6, obtained 6 hours after delivery were significantly higher in Group 1 compared to Group 2 ($p<0.05$ for CRP and $p<0.001$ for IL-6). Although there was no difference between groups in terms of duration of respiratory support, mean duration of hospitalization as well as mortality rate were significantly higher in Group 1 ($p<0.001$, $p<0.05$, respectively).

Conclusions Women diagnosed with a high-risk pregnancy should be treated with antibiotics immediately after a positive cervical culture result, and delivery should be delayed until the success of

antibiotic treatment can be evaluated. Early initiation of maternal antibiotic therapy is associated with shorter durations of hospital stay for newborns. Close follow-up of mothers with high risk pregnancies and extension of treatment duration are critical for determining prognosis in newborn infants.

1161 QUANDRY OVER THE USE OF ANTIFUNGAL PROPHYLAXIS IN PRETERM INFANTS: SURVEY OF CURRENT PRACTICE IN THE UNITED KINGDOM

doi:10.1136/archdischild-2012-302724.1161

¹K Durairaj, ²V Viswanath, ¹M Dyke. ¹Neonatal Medicine, Norfolk and Norwich University Hospital, Norwich; ²Neonatal Medicine, Luton & Dunstable University Hospital, Luton, UK

Introduction Prophylactic anti fungal use reduces the incidence of colonisation and invasive fungal infection in premature neonates. We surveyed the current regimens for the use of antifungal prophylaxis in the tertiary neonatal units across the UK.

Method We enquired about indications for use, drug of choice for prophylaxis, criteria for stopping the prophylaxis and drug used for suspected or proven fungal infection.

Results Out of 52 units 42 [81%] responded. 7 units [17%] did not use any prophylaxis. 26 units [62%] had guidelines on the use of anti fungal prophylaxis. 9 units [21%] used prophylaxis but did not have any guidelines. Of the units using prophylaxis, 43% used birth weight as a criterion ranging from < 750grams to < 1.2kilogram. 51% of units used gestation as a criterion ranging from < 25weeks to < 32 weeks. 20% of units used antibiotic use as their only criterion for starting prophylaxis. 31% used presence of longline as a criterion. Small number of units used abdominal surgery, prolonged intubation, NEC, Candida colonisation, postnatal steroids and ranitidine as a criterion. The commonest drug used for prophylaxis was fluconazole [50%]. 29% of units used nystatin and 12% of units used miconazole gel. 26% of units used the same prophylactic drug when treating suspected or proven fungal infection.

Conclusion Despite evidence of the efficacy of anti-fungal prophylaxis, 17% of tertiary units are not using antifungal prophylaxis for infants at high risk. There remains considerable heterogeneity in indications and the specific antifungal used for prophylaxis.

1162 COMPARATIVE ANALYSIS OF STAPHYLOCOCCUS EPIDERMIDIS STRAINS ISOLATED FROM NEWBORNS

doi:10.1136/archdischild-2012-302724.1162

¹E Hell, ²CG Giske, ¹K Gemzell Danielsson, ³G Marchini. ¹Dept. of Women's and Children's Health, Obstetrics and Gynecology Unit; ²Dept. of Microbiology, Tumor- and Cell Biology; ³Dept. of Women's and Children's Health, Neonatal Unit, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden

Staphylococcus epidermidis are important opportunistic biofilm forming pathogens, particularly causing infection in patients with indwelling medical devices. Preterm infants represent a high-risk group for device-related *S. epidermidis* infections since they require the delivery of nutrients and drugs over long periods.

The present study compared genetic and phenotypic characteristics of *S. epidermidis* strains isolated from blood stream infections of preterm infants (n=10) versus skin isolates obtained from healthy newborns (n=16). Two reference strains were also included the study. Insertion element IS256, as a marker for invasiveness, was analysed by PCR. Antimicrobial susceptibility was testing against cefoxitin, gentamicin and vancomycin. Pulsed-Field Gel Electrophoresis was performed to study clonal relationship among strains.

90% of the blood isolates were resistant to cefoxitin and gentamicin and all these carried IS256. All skin isolates were susceptible to both cefoxitin and gentamicin and all lacked IS256. All of the 28 strains included in the study were susceptible to vancomycin.

We conclude that the *S. epidermidis* strains isolated from blood stream infection in preterm infants are clonally not related to the normal colonizing *S. epidermidis* skin flora at birth, have different phenotypic features related to antimicrobial susceptibility, and have most probably originated from the hospital environment.

1163 EPIDEMIOLOGY OF PROVEN NOSOCOMIAL SEPSIS IN LOW BIRTH WEIGHT INFANTS ADMITTED IN THE LEVEL 3 NEONATAL INTENSIVE CARE UNIT

doi:10.1136/archdischild-2012-302724.1163

M González, C Gómez, MG Espinosa, M Rodriguez, R Santos, E Salguero. Neonatal Intensive Care Unit, Hospital Carlos Haya, Málaga, Spain

Background Nosocomial infections remains a leading cause of morbidity and mortality in Neonatal Intensive Care Unit(NICU).

Aim To assess incidence, etiology and outcome of culture-proven nosocomial sepsis in low birth weight(LBW) infants.

Methods Retrospective study of preterm infants with birth weight < 1500g and proven nosocomial sepsis admitted in NICU of Hospital Carlos Haya during 2011.

Results Sixty neonates experienced at least one or more episode of nosocomial sepsis out 160 LBW infants meaning an incidence of 37.5%. 61% positive blood culture. Table one. Mortality was 6.6% of all patient with proven sepsis and 4% of all positive blood culture. In our series fungal sepsis were the most aggressive being responsible of the 50% of deaths.

Abstract 1163 Table 1 Etiology of nosocomial infections during 2011

Gram Positives	Coagulase negative Staphylococcal (including <i>S.epidermidis</i>)	67%(38% <i>S.epidermidis</i>)
	Enterococcus spp	7%
	<i>S.aureus</i>	1%
Gram negatives	<i>Klebsiella pneumoniae</i>	7%
	<i>Serratia marcescens</i>	5%
	<i>E.coli</i>	1%
	<i>Enterobacter</i> spp	4%
	<i>Pseudomonas</i> spp	5%
	Other gram negatives	1%
Fungal	<i>Candida</i> spp	2%

Conclusions Overall infections rate is similar to the current reports for level 3 hospitals. Gram positives organisms were common being *S. epidermidis* the most frequent. Related to gram negative *K. pneumoniae*, *S. marcescens* and *Enterobacter* were the most frequently isolated. No cases of expanded spectrum of betalactamases bacterial. Mortality is less than previously reported.

1164 REDUCTION IN FUNGAL SYSTEMIC INFECTIONS IN PRETERM NEONATES WITH NYSTATIN PROPHYLAXIS

doi:10.1136/archdischild-2012-302724.1164

MA Ihsan, M Yar, F Sabbagh, M Abu-ouf, MM Abou Al-Seoud. Neonatal Division-Paediatric Department, King Fahad Armed Forces Hospital, Jeddah, Saudi Arabia

Background Invasive fungal infection is an important cause of mortality and morbidity in the very low birth weight (VLBW) and the extremely low birth weight (ELBW) infants. A policy of oral nystatin prophylaxis was introduced with the aim of reducing the incidence of invasive fungaemia among high risk neonates.

Aim To determine whether this policy had reduced the rates of invasive fungal infection.

Methods In December 2004 oral nystatin prophylaxis implemented for babies with birth weight equal or < than 1250 grams starting on 3rd day of life till they retain their birth weight. In 2010