Insofar, blood pressure and sleep-related breathing should be monitored beginning with an obstructive AHI of 3/hour sleep. Children with OSA should be treated early enough. Prevention and early treatment of obesity as a risk factor for OSA as well as for hypertension is becoming an important social challenge.

135 VENTILATORY SUPPORT IN PATIENTS WITH CENTRAL HYPOVENTILATION SYNDROME (CHS) - RESULTS OF THE EU-CHS NETWORK SURVEY

doi:10.1136/archdischild-2012-302724.0135

¹M Migdal, ²H Trang, ³J Peters, ⁴M Katz-Salamon, ⁵G Ottonello, ⁶M Samuels. ¹Children's Memorial Health Institute, Warsaw, Poland; ²Hopital Robert Debre, Paris, France; ³Klinikum Dritter Orden, Munich, Germany; ⁴Karolinska Institutet, Stockholm, Sweden; ⁵G Gaslini Institute, Genoa, Italy; ⁶University Hospital of North Staffordshire, Stoke on Trent, UK

Background and aim of the study Aim of the study was to identify the European centers for diagnosis and treatment of patients with CHS and to collect information on type of ventilatory support currently used for this group of patients.

Method The survey was performed in countries participating in the EU-CHS network project. Using a special template following information were collected: total number of already diagnosed patients, number of currently treated patients, type of ventilatory support (invasive ventilation via tracheostomy, mask ventilation, diaphragm pacing, negative ventilation).

Results The survey was performed during 2011. Replies were received from 15 centers located in 10 European countries. Number of CHS patients already diagnosed and currently treated varies significantly between centers (from 2 to 112 and from 1 to 70 respectively). Total number of already diagnosed CHS patients were 305 and 199 of them are currently treated. Most frequently used type of ventilator support is ventilation via tracheostomy (52% of patients). Diaphragm pacing alone or diaphragm pacing and ventilation is currently used in 42 patients. Only one CHS patient is currently treated with negative pressure ventilation.

Conclusions Results of this study have shown existing significant differences (mainly number of patients) but also some similarities (type of ventilatory support) between centers for CHS diagnosis and treatment located in 10 different European countries. All of the collected data should be used for further work on the European CHS network of reference centers.

The study was supported by the EU-CHS Network (grant number 2008 12 06).

136 SPONTANEOUS BREATHING PATTERNS OF TRANSITIONING PRETERM INFANTS IN THE DELIVERY ROOM (DR) AND INTERACTIONS WITH MANUAL POSITIVE PRESSURE VENTILATION

doi:10.1136/archdischild-2012-302724.0136

^{1,2,3}G Schmölzer, ²J Kaufman, ^{2,4}O Kamlin, ^{2,4}P Davis. ¹Department of Pediatrics, University of Alberta, Edmonton, AB, Canada; ²Neonatal Services, The Royal Women's Hospital, Melbourne, VIC, Australia; ³Department of Paediatrics, Medical University of Graz, Graz, Austria; ⁴Dept. of Obstetrics & Gynaecology, The University of Melbourne, Melbourne, VIC, Australia

Background Positive pressure ventilation (PPV) remains the cornerstone of respiratory support after birth. Effectiveness of PPV in the DR may be compromised by leak, obstruction and inappropriate tidal volume (V_T) delivery.

Aim Describe leak, obstruction and V_T during spontaneous breathing amongst preterm infants transitioning in the DR and interaction of these breaths with mask ventilation.

Methods Patients were enrolled from the control arm of a randomised trial investigating the use of a respiratory function monitor

(RFM) in the DR; infants < 32 weeks who were stabilised by paediatric trainees unaware of RFM data were included. During spontaneous breathing on mask continuous positive airway pressure (CPAP) and the delivery of PPV, airway pressures, gas flow and V_T were recorded. Data were analysed by breath type.

Results In 29 infants, a total of 3864 inflations (mechanical) and inspirations (infant breaths) were analysed (Table 1). Overall mask leak was highest during spontaneous inspirations occurring between manual inflations and those during CPAP. The average V_T was highest during combined inspiration and inflation.

Abstract 136 Table 1 Type of Inflation/Breath

	Number of Inflations	Mask Leak (%)	Tidal volume (mL/kg)
Manual Inflation	2055 (53%)	41 (32–53)	8.2 (6.1–10.7)
Assisted Inflation	335 (9%)	61 (46–81)	9.3 (4.5–12.6)
Spontaneous breath during PPV	181 (5%)	92 (65–100)	2.9 (2.1–4.3)
Spontaneous breath while on CPAP	1285 (33%)	66 (41–100)	4.4 (2.5–6.2)

Conclusion Preterm infants are active participants in stabilisation in the DR. V_T and mask leak vary depending on the interaction between the infant's breathing efforts and the inflations provided by the resuscitator.

137 HIGH FLOW NASAL CANNULAE (HFNC) OR NASAL CONTINUOUS POSITIVE AIRWAY PRESSURE (NCPAP) POST-EXTUBATION IN PREMATURE INFANTS? A RANDOMISED CONTROLLED TRIAL

doi:10.1136/archdischild-2012-302724.0137

¹C Collins, ¹JR Holberton, ¹C Barfield, ²PG Davis. ¹Neonatal Medicine, Mercy Hospital for Women; ²Newborn Research, Royal Women's Hospital, Melbourne, VIC, Australia

Background NCPAP facilitates successful extubation. It is unclear whether HFNC are as effective as NCPAP in preventing extubation failure. In addition to an alternative modality of respiratory support HFNC may result in less nasal trauma than NCPAP.

Methods 132 preterm ventilated infants were randomised and stratified by gestation (< 28 vs 28–32 weeks). Primary outcome was extubation failure defined by a composite of 3 pre-specified failure criteria in the 7 days post-extubation. Individual failure criteria were not mutually exclusive and are defined; Apnoea, > 6 episodes in 6 hours or 1 requiring IPPV, Acidosis, pH < 7.25 & pCO_2 > 66 mmHg, and > 15% increase in FiO_2 from extubation. A nasal trauma score was adapted from Kaufman [E-PAS 2007:61390].

Results

Abstract 137 Table 1

	HFNC N=67	NCPAP N=65	
Male n (%)	33 (49)	41 (63)	
Birthweight g mean (SD)	1123 (317)	1105 (374)	
Mean Completed Weeks gestation (SD)	27.9 (1.95)	27.6 (1.97)	
Failed Extubation in 1st week by Composite Criteria n (%)	15 (22)	22 (34)	
Apnoea: >6 in 6hrs or 1 needing IPPV n (%)	14 (21)	17 (26)	
Acidosis: pH < 7.25 & pCO_2 > 66 mmHg n (%)	0	3 (5)	
>15% increase in FiO_2 post-extubation n (%)	7 (10)	12 (18)	
Reintubated in 1st week n (%)	7 (10)	8 (12)	
Nasal Trauma Score 1st week mean (SD)	3.1 (7.2)	11.8 (10.7)	p < 0.001*

Conclusions Rates of extubation failure were not significantly different between the groups. HFNC resulted in significantly less nasal trauma than NCPAP. This benefit may need to be considered in post-extubation respiratory support for preterm infants.

138 MASK OR NASAL TUBE TO PROVIDE POSITIVE PRESSURE VENTILATION (PPV) TO PRETERM INFANTS IN THE DELIVERY ROOM (DR)-THE MONT TRIAL

doi:10.1136/archdischild-2012-302724.0138

¹COF Kamlin, ²K Schilleman, ¹JA Dawson, ²E Lopriore, ¹S Donath, ¹GM Schmoelzer, ²FJ Walther, ¹PG Davis, ²AB te Pas. ¹Royal Women's Hospital, Melbourne, VIC, Australia; ²Leiden University Medical Center, Leiden, The Netherlands

Background For initial PPV a face mask is used, but obstruction and leak often occurs. A nasal tube is an alternative interface in the DR to provide PPV. Its safety and efficacy have not been tested in a large RCT in extremely preterm infants.

Objective To determine whether a single nasal tube is more effective than face mask during the stabilisation of infants born between 24 and 29 weeks.

Design and methods An RCT performed in Melbourne and Leiden. Infants were randomized immediately prior to birth to receive PPV using a T-piece with either a nasal tube or round mask. Resuscitation guidelines were standardised. Criteria for intubation: cardiac compressions, apnea, CPAP >7cmH₂O and FiO₂ >0.4.

Primary outcome: intubation in the first 24 hours from birth.

Results In total 368 infants were randomized. Baseline variables, primary and secondary outcomes were similar (table).

Abstract 138 Table 1 Baseline characteristics and outcome

	Mask (N= 187)	Nasal Tube (N= 181)	OR (95% CI)
GA (wks)	27 (26–29)	27 (26–28)	-
Birthweight (g)	1000 (807–1196)	946 (780–1182)	-
Intubation in first 24hrs	102 (54)	97 (54)	0.96 (0.64–1.45)
Intubation in DR	58 (31)	41 (23)	0.65 (0.41–1.04)
Air Leak	14 (8)	11 (6)	0.80 (0.35–1.81)
Total (MV+CPAP) days of ventilation	13 (6–41)	16 (4–41)	NS
Death or BPD	68 (36)	64 (35)	0.96 (0.62–1.47)

Conclusions In infants < 30 weeks gestation receiving PPV in the DR, there were no differences in short term outcomes using the nasal tube compared to the face mask.

139 RESEARCH NETWORKS IN PEDIATRIC PRIMARY CARE: EAPRASNET

doi:10.1136/archdischild-2012-302724.0139

¹S del Torso, ²D van Esso, ³A Hadjipanayis, ⁴Z Grossman. ¹ULSS 16, Padova, Italy; ²Primary Care Center 'Pare Claret', Barcelona, Spain; ³Pediatric Department, Larnaca General Hospital, Larnaca, Cyprus; ⁴Maccabi Health Services, Tel Aviv, Israel

EAPRASnet is the European Academy of Paediatrics Research in Ambulatory Setting Network, whose members are European primary care paediatricians committed to research in their practices that expressed their will to join a research network.

The network is the research laboratory of the primary care setting and in the latest study, EAPRASnet has enrolled a total of 685 paediatricians from 21 countries.

A recruitment survey showed that areas of most interest for research were: quality of care indicators, communication with parents, obesity, attention deficit hyperactivity disorder and effective

well child care. Main incentives for participation in a research project were interest in the topic (81%) and effort to improve quality of care (71%). Lack of time was the leading reported obstacle (72%).

A second electronic survey assessing primary care paediatricians' estimations and practices regarding parents' vaccination refusal was sent to 395 EAPRASnet members, with a response rate of 87%. Of respondents who vaccinate in their clinic, 93% estimated the total vaccine refusal rate less than 1%. Sixty nine percent of all respondents prefer a shared decision making approach to handle parents' vaccine hesitancy.

A recent survey on Use of Antibiotics in Upper Respiratory tract infections has been performed to evaluate European primary care paediatricians' knowledge, attitudes and practice regarding antibiotic prescribing for upper respiratory infections in order also to identify targets for future intervention studies.

140 RESEARCH NETWORKS IN PEDIATRIC PRIMARY CARE: TRANSITION TO REALITY

doi:10.1136/archdischild-2012-302724.0140

T DeWitt. *Pediatrics, Cincinnati Children's Hospital Medical Center, University of Cincinnati College of Medicine, Cincinnati, OH, USA*

Research networks in pediatric primary care, often called practice-based research networks (PBRN) in the United States, have an increasingly important role in the spectrum of research that spans from basic research to practical and effective application in primary care settings. In addition to assessing efficacy, PBRN's are particularly critical in assessing effectiveness in the real world of community practice - the final step in translating new discovery into practice. Key elements of effective translation that can be assessed by PBRN's include rapidity of adoption, adherence, and quality improvement systems to assure consistent application. This presentation will discuss PBRN's in the context of the National Institute of Health's "Roadmap for Medical Research" and an emphasis on translational research. Examples of the use of pediatric PBRN's in this capacity, including ADHD and asthma, with a focus on effectiveness will be discussed.

141 TRANSCUTANEOUS ELECTRICAL STIMULATION (TES) AND INTRACTABLE CHRONIC CONSTIPATION IN CHILDHOOD

doi:10.1136/archdischild-2012-302724.0141

^{1,2,3}YI Yik, ^{3,4}JM Hutson, ³BR Southwell. ¹Department of General Surgery, Faculty of Medicine, University of Malaya, Kuala Lumpur, Malaysia; ²Department of Paediatrics, University of Melbourne; ³Gut Motility and Research Laboratory, Murdoch Children's Research Institute; ⁴Department of Urology, Royal Childrens Hospital Melbourne, Melbourne, VIC, Australia

Background and aims Intractable chronic constipation in the form of slow-transit constipation (STC) often resistant to medical treatment. Some children required appendicostomy for antegrade enemas to improve symptoms. Transcutaneous electrical stimulation (TES) was used by physiotherapists to overcome STC in children successfully. This study aimed to examine the effectiveness of home-based TES when STC children were trained by a naïve clinician. We hypothesized TES would improve symptoms with reduce laxative use.

Methods A prospective study (2009–2011) whereby a surgeon was trained to deliver TES method to STC children/parents, who then self-administered TES at home (1 hr/day x 6 months) using a battery-powered interventional stimulator. Daily continence diary (including laxative use) was recorded before and throughout TES; PedsQL4.0 questionnaires and gastrointestinal nuclear transit scintigraphy (NTS) were completed before and after TES. Appendicostomy for antegrade enemas was offered if TES failed to improve