physical activity inquiry, the connection between physical activity and food intake.

Results The overweight prevalence was 7.1% and that of obesity 12.2%. The overweight and obese children, as compared to the normal weight ones eat, daily, sweets (57.7%, 62.1% vs. 12.3%), soft drinks (23.9%, 39.9% vs. 5.2%), fast-food (22.8%, 36.7% vs. 9.4%), meat products (54.6%, 76.1% vs. 43.8%), fruits (16.8%, 7.3% vs. 48.5%), vegetables (41.3%, 13.9% vs. 84.5%), cereals (31.1%, 24.7% vs. 49.8%), milk products (18.1%, 32.2% vs. 62.3%). 54.7% of the overweight children and 60.1% of the obese do not perform any physical activity in their spare time, as compared to the normal weight (29.8%). The food consumption while watching TV or playing on the computer was associated with overweight and obesity in children.

Conclusions Following the nutritional and physical activity investigation in the school children, I noticed an alimentary abuse, both quantitative and qualitative, associated with a physical activity decrease.

1445 THE GROWTH AND NUTRITIONAL STATUS AMONG SCHOOL CHILDREN IN NOVI SAD IN SCHOOL YEAR 2010/2011

doi:10.1136/archdischild-2012-302724.1445

¹M Hadnadjev, ¹M Budakov, ²D Hadnadjev, ¹M Bratic. ¹Paediatrics Department, Health Centre Novi Sad; ²Center for Radiology, Clinical Center Vojvodina, Novi Sad, Serbia

Aim To examine physical growth and nutritional status of school children in Novi Sad, during the school year 2010/2011 and determine differences in relation to primary and secondary school children.

Methods On the medical examination of school children aged 7–14 (primary school) and 15–19 (secondary school) in Novi Sad we registered the total number of children: with normal weight (from –1.5SD to +1.5SD), obese (+1.5 to 3SD) and thin children (from –1.5SD to –3SD); of average height (from –1.5SD to +1.5SD), tall (+2SD, +3SD) and short (–2SD, –3SD) (nomogram provided by Vukovic).

Results From the total number of the enrolled pupils (n=40861) we have examined 37.25% (n=15222): there were 12783 (83.98%) of average height, 2325 (15.27%) tall pupils and low height pupils 114 (0.75%); 11990 (78.77%) of well-nourished, obese 2404 (15.79%) and underweight children 828 (5.44%). After comparing the data from primary and secondary schools, a statistically significant difference, considering a total number of tall, short, obese, and thin pupils, was in favor of primary school children (p<0.01). There were more children with normal weight in secondary schools (p>0.05), and more obese children in primary schools (p>0.05).

Conclusion During 2010/2011, the assessment of growth and nutritional status results of school children were in favor of primary school children. There were more children with normal weight children in secondary schools, and more obese children in primary schools, but the differences were not significant.

1446 INFANT FEEDING CHOICES MAY HAVE GENDER EFFECT ON GROWTH DURING INFANCY

doi:10.1136/archdischild-2012-302724.1446

K Nagahara, K Dobashi, K Itabashi. *Pediatrics, Showa University, Tokyo, Japan*

Background and Aims A retrospective analysis of healthy Japanese children was conducted to determine whether the growth pattern is altered by breast-feeding (BF) or formula-feeding (FF).

Methods The data of 204 elementary students, age 6 to 9 years were obtained from their parents by questionnaires. The BF and FF group were defined as those fed by only breast milk or formula at 4 months of age. Seventy-one children (31 boys, 40 girls) were in BF and 30 (19 boys, 11 girls) were in FF. The anthropometric data at

birth, 1, 4, 7, 10, 18, 36 months-old and investigation time in BF was compared to those in FF.

Result No significant differences were observed in the anthropometric data between BF and FF girls. The BF boys mean body weight (BW) at 18 months was almost 1,000g lower than those in FF. The BW-SD score was significantly smaller from 4 to 18 months, and BMI was lower from 10 to 36 months in BF boys. However, both BWSD and BMI had no differences between BF and FF boys from 6 to 9 years. Multiple regression analyses showed that the birth weight, prepregnancy BW, and infant feeding choice were significant factors associated with BWSD, and feeding choice was the only significant factor associated with BMI at 18 and 36 months.

Conclusion Infant feeding choices may have gender effect on growth during infancy. When we evaluate infant growth, we should consider not only infant feeding choices, but also gender, birth weight and prepregnancy BW.

1447 WEIGHTING THE FACTORS ASSOCIATED WITH CHILDREN OBESITY: AN INTERNATIONAL PERSPECTIVE TOWARD AN UNIFIED MODEL

doi:10.1136/archdischild-2012-302724.1447

D Gregori, I Baldi, OBEY-AD Study Group. *Cardiologic, Thoracic and Vascular Sciences, University of Padova, Padova, Italy*

Beside genetic predisposition, several factors have been proposed to promote overweight and eventually obesity in children, from the socio- and built- environment down to behavioural attitudes (Bouchard, 2007). How this model is shared by different cultural settings is however unclear and less investigated.

Using a unified protocol for data collection, a cross-sectional study has been performed on 960 children in India, Italy, Germany, France, UK, Argentina, Mexico and Brazil. Children CDC z-scores for BMI have been evaluated in association with several known factors influencing overweight and obesity (maternal and neonatal aspects, socio-economic familiar status, BMI of parents, physical activity, nutrition habits, screen activities). Based on a random-effect mixed effect model and the Kullback-Leibler Entropy Measure, the capability to explain variability in BMI of such factors has been computed. Percentages of explained variation are given as follows: India 41.4%, Italy 46.6%, Germany 65.6%, France 52.3%, UK 70.1%, Argentina 62.3%, Brazil 59.7%, Mexico 58.2%. Capability of proposed factors to capture variability in BMI is significantly higher in UK (p=0.03) than in other countries, being significantly lower in an emerging country like India (p=0.042). These results may suggest that more intense research should be specifically targeted to capture risk factors which are specific for that given cultural setting in addition to the general ones.

1448 EARLY FORMULA FEEDING PRACTICES AND THEIR POTENTIAL CONTRIBUTION TO LATER OBESITY RISK

doi:10.1136/archdischild-2012-302724.1448

¹R Tarrant, ²M Sheridan-Pereira, ³K Younger, ³J Kearney. ¹Department of Clinical Nutrition and Dietetics, Our Lady's Children's Hospital; ²Department of Paediatrics and Newborn Medicine, Coombe Women and Infants University Hospital; ³Department of Biological Sciences, Dublin Institute of Technology, Dublin, Ireland

Background and Aims Early feeding practices, including early introduction to solid foods and overfeeding, are known risk factors for childhood obesity. This study aimed to assess maternal formula feeding practices and infant formula feeding patterns, factors that are known to potentially contribute to later obesity risk.

Methods This Irish prospective observational study involved the recruitment and follow-up of 450 eligible mother-infant pairs to 6 weeks postpartum. Data related to formula milk consumption patterns, formula type/brand changing, additions of solids to bottle