Material and Method A retrospective statistic study of ADI in children aged 0–16 years, admitted to 2nd Pediatric Clinic, Emergency County Hospital Craiova, over a period of 6 years (01.01.2006–31.12.2011).

Results Of the total number of 14427 admitted children, aged 0–16 years, 645 presented acute intoxications with various etiologies, among which 252 had ADI; intoxication type: accidental in 154 (61.1%), voluntary in 98 (38.9%).

Accidental ADI: sex M/F=54/100, social environment U/R=93/61, age group (years): 0–1/1–3/3–6/6–10= 27/68/33/26.

Voluntary ADI: M/F=23/75, U/R=59/39, age (years): 6–10/10–14/14–16= 7/37/54.

Etiologic spectrum in the studied group: AINS/antialgics in 39 (15.5%), methoclopramid 29 (11.5%), anti-epileptics 24 (9.5%), tranquilizers/sedatives 15 (5.9%), neuroleptics 9 (3.6%), parasympatholitics 8 (3.2%), antibiotics/antiparasitary 17 (6.7%), drugs with cardio-vascular effect 8 (3.2%), drug combinations 41 (16.3%), other drugs 12 (4.8%), unknown 50 (19.8%). Clinical forms: mild in 127 (50.4%), moderate 101 (40.1%) and severe 24 (9.5%). No deaths were registered with ADI.

Average period of hospitalization: accidental ADI 3.33±2.54 (1–9), voluntary ADI 3.28±1.24 (1–6) days.

Conclusions ADI represented 39.1% of the total number of acute intoxications; 61.1% of ADI were accidental. Most ADI were caused by AINS, methoclopramid, anti-epileptics. ADI prevailed in females, in urban children, both in voluntary and accidental ADI. The clinical forms were mostly mild.

1654 RELATIONSHIP BETWEEN MYOCARDIAL INJURY, OXIDATIVE STRESS MECHANISM AND SEPSIS/SEPTIC SHOCK IN INFANTS SUBMITTED TO SURGERY FOR CONGENITAL HEART DISEASE

doi:10.1136/archdischild-2012-302724.1654

¹MS Oliveira, ²EM Floriano, ³PH Manso, ⁴RN Sgabieri, ⁵JG Klamt, ⁶EZ Martinez, ⁷WVA Vicente, ⁸SG Ramos. ¹Department of Pathology; ²Department of Pediatrics; ³Department of Surgery and Anatomy; ⁴Department of Biomechanics, Medicine and Locomotive Apparatus Rehabilitation; ⁵Department of Statistics, Faculty of Medicine of Ribeirão Preto, University of São Paulo, Ribeirão Preto, Brazil

Background and Aims A progressive ventricular dysfunction caused by ischemic myocardial injuries remains one of the leading causes of death during the postoperative course in congenital heart disease (CHD). The aim of this study was to investigate the role of oxidative stress in these myocardial injuries.

Methods Myocardial injuries and oxidative stress mechanisms were assessed by histopathology and immunohistochemistry and quantified by morphometrical analyses.

Results Myocardial injury was observed in pediatric patients submitted to surgery for CHD with cardiopulmonary bypass, followed by lethal exit. Oxidative stress mechanisms were directly related to these particular types of myocardial injuries. Importantly, 4-hydroxynonenal (4-HNE), a marker of lipid peroxidation, is strongly expressed, especially in irreversible myocardial lesions. Although morphologically similar, myocardial injuries observed in patients who evolved with sepsis in the peri-operative period exhibited a completely different set of oxidative stress mechanisms. Increased concentrations of nitrotyrosine protein adducts were observed in these patients, suggesting that peroxynitrite-mediated protein nitration may be the predominant oxidative stress mechanism found in these situations.

Conclusions The underlying mechanisms of these lesions seem to be related to the development of ischemia or ischemia/reperfusion followed by oxidative stress mechanisms that vary depending on whether sepsis was present. While the exact mechanism is not fully understood, it has been suggested that endogenous catecholamine release could have a role in this process.

1655 LEVOSIMENDAN AND MILRINONE: A SAFE COMBINATION?

doi:10.1136/archdischild-2012-302724.1655

¹A Morales Martínez, ²P García Soler, ³C Yun Castilla, ⁴L Romero Moreno, ⁵JM Camacho Alonso. ¹Pediatric Intensive Care Unit, Hospital Regional Universitario Materno Infantil Carlos Haya; ²Hospital Materno Infantil Carlos Haya, Málaga, Spain

Background/Aims Levosimendan is an inotropic and vasodilator drug. Most protocols suggest avoiding other vasodilators, inodilators after its introduction. The technical data recommends not using with other vasodilators. We report our experience of concomitant use of levosimendan and milrinone in a series of patients.

Methods Observational study. Review of medical records of concomitant use of levosimendan (24 hours infusion without loading dose) and milrinone, from June 2009 to December 2011. Multivariate analysis of epidemiologic data, pathology, type of heart failure, indication of the drug and side effects.

Results 81 cases received levosimendan. Among them, 64 (79%) received simultaneously milrinone. Mean age 14.8 months (5 days–112 months) 57.8% males. 87.5% were postoperative cardiac surgical patients (41% tetralogy of Fallot). Right ventricular failure was the most common indication (56.9%) followed by left ventricular failure (29.2%) and biventricular failure (12.3%). Diastolic dysfunction was reported in 49.2% of our patients, 46.1% systolic dysfunction and 4.6% both. In 31% of cases both drugs were initiated simultaneously (operating theatre). Milrinone was the first drug in 41.5% cases. The average dose of milrinone was 0.8 mcg/kg/min. 19/63 cases (30%) suffered from hypotension. In 7 cases (11% of total) milrinone was suspended (without association to type of pathology or dysfunction). In the other 12 patients, infusion of milrinone was decreased, but not suspended. Among the cases of right ventricular failure, hypotension appeared in 32.4%, in 10.5% with left ventricular failure and in 55.5% with biventricular. There were no complications associated.

Conclusions In our series, administration of levosimendan and milrinone was safe. The appearance of hypotension was controlled and was not associated with other complications. Patients at increased risk of hypotension were those with biventricular dysfunction. Larger prospective studies are necessary to test the safety of this combination.

1656 BIOMARKERS FOR SEPSIS, MULTIPLE ORGAN DYSFUNCTION SYNDROME (MODS) AND MORTALITY AFTER OPEN HEART SURGERY IN CHILDREN

doi:10.1136/archdischild-2012-302724.1656

¹T da Rocha, ²L Mülle, ³C Ricachinevsky, ⁴A Botta, ⁵P Carvalho. ¹Uti Pediátrica Hospital Santo Antonio, Universidade Federal do Rio Grande do Sul; ²Uti Pediátrica Hospital Santo Antonio, Santa Casa de Misericórdia de Porto Alegre, Porto Alegre, Brazil

Background and Aims Biomarkers can help to predict risk of unfavorable outcomes after open heart surgery in children.

Methods We performed a retrospective cohort of 121 children after open heart surgery. We analyzed the serum blood lactate, base excess (BE), blood glucose, central venous oxygen saturation (SATvc), troponin I, C-reactive protein (CRP), and leukocyte counts in different postoperative days (POd).

Results There were 7.4% deaths, 27.3% of sepsis and 60.3% of MODS. For death, showed better power PO1d and PO2d lactate (OR = 24.1 [CI 4–112]) and (OR = 9.7 [CI 1.2 to 85.7]); PO1d EB (OD = 30.6 [CI 2.6 to 351]); PO1d total leukocytes (OD = 5.8 [1.2 to 29.8]) For sepsis, showed better power: PO6h glucose (OD = 2.4 [1.03 to 5.7]); POI and PO3d SATvc (OD = 2.4 [1.09 to 5.8]) and (OD = 25.6 [2.2–298]); PO6h troponin I (OD = 2.8 [1.1 to 6.8]) and PO1d bastonades (OD = 6.5 [1.4 to 29.6]). For MODS, showed better power: PO6h and PO1d SATvc (OD = 12.2 [2.6 to 55.7]) (OD = 2.87 [1.1 to 7.4]) and, POI/PO6h/PO1d troponin-I (OD = 3.2 [1.6 to 8.0]); POI/PO6h CPR (OD = 3.7 [1.3 to 10.8]).

Conclusions Blood lactate, base excess, leukocyte count and percentage of bastonades are suitable markers of mortality. Blood glucose, central venous oxygen saturation and cardiac troponin-I are valued predictors of sepsis, and the last two were also reliable markers of MODS and CPR marker only the latter.

1657 MORTALITY IN POST-CARDIAC SURGERY IN CHILDREN: THE ROLE OF MULTIPLE ORGAN DYSFUNCTION

doi:10.1136/archdischild-2012-302724.1657

¹T da Rocha, ²A Botta, ³C Ricachinevsky, ⁴LD Mülle, ⁵P Carvalho. ¹Uti Pediátrica Hospital Santo Antonio, Universidade Federal do Rio Grande do Sul; ²Uti Pediátrica Hospital Santo Antonio, Santa Casa de Misericórdia de Porto Alegre, Porto Alegre, Brazil

Background and Aim The multiple organ dysfunction syndrome (MODS) is associated to worse prognosis in critical setting. Its role after open heart surgery is still purely studied.

Methods We made a retrospective cohort of 121 children to study factors associated to mortality after open heart surgery such as systemic inflammatory response syndrome (SIRS), MODS, sepsis, low cardiac output syndrome (LCOS), respiratory and cardiac support time.

Results 7.4% of non survivors occurred. The presence of sepsis in the first postoperative day had the highest odds ratio (OR) = 31.71 (2.6 to 393.8), followed by renal dysfunction on the third day, OD=14.1 (2.9 to 66.6); uni ventricular correction, OD=14.2 (2.9 to 66.66); the presence of MODS on the third day, OD=10.0 (1.9 to 50.9); presence of LCOS on the fifth day, OD=9.1 (2.1 to 40.2); and cardiac and respiratory dysfunction in the fifth day, OD=6 (1.4 to 25.6). On the other hand, the absence of SIRS in the immediate postoperative period was protective, with OD=0.92 (0.87 to 0.97). Furthermore, the mean time of cardiac support was higher in non survivors (98.8 vs. 53.7 hours), also respiratory support (42.2 h vs. 87 hours) and time of hypotension (85.2 h vs. 62.2 hours) than in survivors ($p < 0.05$).

Conclusions The increased risk of mortality due to MODS appears to be caused by primary cardiac dysfunction, as there is also an association with LCOS, renal and respiratory dysfunctions secondary to the first.

1658 ACUTE KIDNEY INJURY AFTER HEART SURGERY IN CHILDREN

doi:10.1136/archdischild-2012-302724.1658

^{1,2}MGDS El Halal, ¹PRA Carvalho, ²CV Bohrer, ²AP Pearson. ¹Universidade Federal do Rio Grande do Sul; ²Hospital da Criança Santo Antônio, Porto Alegre, Brazil

Background and Aims Acute kidney injury (AKI) has been associated to adverse outcomes in children after heart surgery, but previous studies have used several different definitions of AKI. After publication of the pediatric RIFLE (pRIFLE) criteria, it has been widely used as a definition for AKI in children. This study aims to investigate association between occurrence of AKI according to pRIFLE criteria and adverse outcomes in children after heart surgery.

Methods Children submitted to open heart surgery in a hospital in Brazil were followed from arrival until death or discharge from the Pediatric Intensive care Unit (PICU). The exposition variable was occurrence of AKI according to pRIFLE criteria, which divides AKI in three categories: R-Risk, I-Injury, F-Failure, according to changes in urine output or the estimated glomerular filtration rate. The outcomes studied were death, length of mechanical ventilation (MV) and length of PICU stay.

Results Eighty five children were studied. Forty seven (55.3%) did not have AKI during PICU stay, while 22 (25.9%), 7 (8.2%) and 9 (10.6%) were classified as R, I and F, respectively. Comparing to children who did not develop AKI, the relative risk for death was

1.07 (0.09–12.48), 16.87 (2.14–132.50) and 11.25 (1.55–81.61) in the R, I and F group, respectively ($p = 0.001$). Lengths of MV and of PICU stay were significantly higher in those with any degree of AKI.

Conclusions Occurrence of AKI according to pRIFLE criteria is associated to adverse outcomes in children after open heart surgery.

1659 POST - OPERATIVE ICU COURSE OF INFANT BELOW 2.2 KG UNDERGOING CARDIAC SURGERY

doi:10.1136/archdischild-2012-302724.1659

A Mehmood, SR Ismail, MS Kabbani, RM Abusuliman, HK Najm. *Cardiac Sciences-Pcicu, King Abdulaziz Medical City for National Guards - Pediatric Cardiac Intensive Care, Riyadh, Saudi Arabia*

Introduction Infants with low body weight (LBW) are major challenges for post cardiac surgery care. We conducted this study to compare post-operative course and outcome of infant weighing 2.2 Kg or less with matching group of infants with normal body weight who underwent similar cardiac surgery.

Methods We reviewed retrospectively all infants below 2.2 kg who underwent cardiac operation at our institution from January 2001 to March 2011. Cases with LBW (group A) were compared with matching group (Group B) of normal body weight infants who had similar cardiac surgery and matching surgical risk category. We compared demographic, ICU parameters, complications and short-term outcome of both groups.

Results Thirty seven patients were included in group A and 39 in group B. Except for Weight (2.13 ± 0.08 kg in Group A vs. 3.17 ± 0.2 kg in group B); there was no statistical difference in demographic data between both groups. Cardiac procedures included coarctation repair, Arterial switch, VSD repair, tetralogy of Fallot repair, systemic to pulmonary shunt and Norwood procedures. Patients in group A had statistically significant difference from group B in term of bypass time ($p = 0.01$), duration of inotropes ($p = 0.01$), duration of mechanical ventilation ($p = 0.004$), number of re-intubations ($p = 0.015$), PCICU length of stay ($p = 0.007$) and mortality (13.5% in group A vs. 0% in group B, p value 0.02).

Conclusion Patients with LBW below 2.2 Kg can go for cardiac surgery with overall satisfactory result but with increase risk of ICU morbidity and mortality.

1660 RISK FACTORS FOR MORTALITY AFTER TRANSCATHETER INTERVENTION AFTER CARDIAC SURGERY IN CHILDREN WITH CONGENITAL HEART DISEASE

doi:10.1136/archdischild-2012-302724.1660

¹M Amin, ²A Hussain, ¹A Kouatli, ^{1,3}S Deraz, ⁴H Abozeid, ⁴J Al Ata. ¹CVD, King Faisal Specialist Hospital & Research Centre Gen. Org; ²CVD, King Faisal Specialist Hospital & Research Centre, Jeddah, Saudi Arabia; ³Pediatrics, Minoufiya University, Shibin El Kom, Egypt; ⁴King Faisal Specialist Hospital & Research Centre Gen. Org, Jeddah, Saudi Arabia

Aim Risk stratification for mortality after cardiac catheterization in early post-operative children.

Introduction Transcatheter intervention has evolved overtime to expose high risk, sick children to such procedures. The risk of dying, rises significantly in emergent and critically sick patients. Rarely published, we reviewed records of 35 patients, 43 procedures during immediate/early post operative period after cardiac surgery for congenital heart disease.

Methods Multivariate analysis for age, weight, sex, time between surgery/intervention, and the intervention on natural vasculature/synthetic material, was done, as possible risk factors for mortality.

Results Thirty patients survived and cleared from ICU in due course. None of the risk factors could be attributed to the mortality.