Conclusion In a small cohort of children undergoing CPB surgery, increased coronary flow and, to less extent, prolonged CPB appeared to be associated with heightened risk for delayed myocardial repolarization after surgery.

1664 IMPACT OF CHYLOTHORAX ON THE POST OPERATIVE OUTCOME AFTER PEDIATRIC CARDIOVASCULAR SURGERY

doi:10.1136/archdischild-2012-302724.1664

SR Ismail, M Kabbani, H Najm, G Shaath, A Jijeh. *Cardiac Science, King Abdulaziz Medical City for National Guards, Pediatric Cardiac Intensive Care, Riyadh, Saudi Arabia*

Background Chylothorax is the accumulation of Chyle in the pleural cavity, usually develops after disruption of the thoracic duct along its intra-thoracic route. In the majority of cases this rupture is secondary to trauma (including cardio thoracic surgeries). Chylothorax is a potentially serious complication after cardiovascular surgeries that require early diagnosis and adequate management.

Methods A retrospective study of all cases Complicated with Chylothorax after pediatric cardiac surgery in King Abdulaziz Cardiac Center between January 2007 and December 2009 was conducted. The study aim to determine the risk factors, and the impact on the post operative course.

Results We have 1135 cases operated during the study period, 57 cases (5%) were complicated with chylothorax in the post operative period, The most common Surgeries complicated with chylothorax were the single ventricle repair surgeries (Glenn- Fontan) 15 cases (27%), followed by the arch repair cases 10 cases (18%), the ventricular septal defect cases 10 cases (18%), the Atrioventricular septal defect cases 7 cases (12%), the arterial switch cases 6 cases (11%), and others 8 cases (14%). The ICU stay, the length of hospital stay and the bypass time were significantly longer in the chylothorax group, also the ventilation time, the inotropes duration and number were higher in the chylothorax group.

Conclusion Chylothorax after pediatric open heart surgery is not an uncommon complication, it Occurs more commonly with single ventricle repair and Aortic arch repair surgeries, it has a significant impact on the post operative course and post operative morbidity.

1665 NONINVASIVE MONITORING OF HEMOGLOBIN CONCENTRATION IN PEDIATRIC CARDIAC POSTOPERATIVE PATIENTS

doi:10.1136/archdischild-2012-302724.1665

P García Soler, JM Camacho Alonso, A Morales Martínez, G Milano Manso. *Pediatric Intensive Care Unit, Hospital Regional Universitario Materno Infantil Carlos Haya, Málaga, Spain*

Background/Aim One of the major concerns after cardiovascular surgery is to detect bleeding and optimize hemoglobin (Hb) to improve cardiac output, leading to repeated blood tests. We describe our experience with a new method of a noninvasive and continuous measurement of Hb concentration.

Methods Partial results of a prospective study in children after cardiac surgery, from January to March 2012. SpHb monitoring was performed during the first 48 hours after surgery (Pulse Co-Oximeter Radical-7™/7.8.0.1, Masimo, CA). When each blood sample was taken, we collected the data from Radical-7™, using central laboratory as a reference method (SiemensADVIA2120i). Data are reported as mean values and SD (normally distributed) and as median values and minimum-maximum range (distribution non-normal).

Results 78 blood samples were drawn from 21 patients, with a median age of 1.3 months (0.23–56), a median weight of 3.7 kg (3–16.6) and a perfusion index (PI) of 1.2 (0.21–13). All the measurements were performed in sedated subjects, 79.5% of them were intubated. Mean Hb on the laboratory analyzer was 12.47±2.28 g/dl and mean Hb on the pulse oximeter (SpHb) was 13.2±2.26 g/dl; its

correlation coefficient was 0.75 and R^2 was 0.55 ($p < 0.05$). The mean of differences between both methods was 0.75 ± 1.6 g/dl. Bland-Altman plot shows that 65% had a difference < 1 g/dl in comparison with laboratory hemoglobin.

Conclusions SpHb offers moderately acceptable accuracy in pediatric cardiac postoperative patients, maybe influenced by the low weight and PI of our patients. It has the advantages of providing continuous measurements, noninvasively, which may facilitate hemoglobin monitoring in the intensive care unit.

1666 INCIDENCE AND RISK FACTORS FOR POST CARDIAC SURGERY ARRHYTHMIAS IN A TERTIARY PAEDIATRIC CARDIAC REFERRAL CENTRE

doi:10.1136/archdischild-2012-302724.1666

¹K Adhvaryu, ²R Johnson, ¹N Shetty. *¹Paediatric Intensive Care Unit; ²Department of Cardiology & Cardiothoracic Surgery, Alder Hey Children's Hospital, Liverpool, UK*

Background and Aims Arrhythmias in the immediate post-operative period are a significant cause of morbidity. Its incidence varies between 15–79% and depends on case mix of surgical patients and methods used to identify them. Our aim was to determine the incidence and risk factors for their occurrence in our regional cardiac referral centre.

Methods A retrospective study was conducted including all paediatric (newborn to 18 years) patients who underwent a surgical procedure requiring cardio-pulmonary bypass between September 2010 and August 2011. Patients were identified and data gathered from the cardiac databases, electronic patient records and typed operation notes. Patients were excluded if they had pre-operative arrhythmias or if the arrhythmias during surgery were so transient such that they only occurred in the operation theatre.

Results Of the 346 cases that met inclusion criteria, 15% developed arrhythmias of which half required intervention. Risk factors for arrhythmias included surgery involving the septum (21% vs 8% non-septal), male (17% vs 14%) and age less than one year (20% vs 9%). Those with arrhythmias had significantly longer bypass (160min vs 116min, $p < 0.05$) and cross clamp times (101 min vs 77min, $p < 0.05$). Transient rhythm problems that required intervention were sinus bradycardia, junctional rhythm, junctional ectopic tachycardia and supra ventricular tachycardia, whilst 43% of those with heart block required permanent pacing.

Conclusion Despite advances in surgical techniques arrhythmias after cardiac surgery were seen in a significant number (15%) of our patients although only a small proportion ($< 1\%$) require long term interventions.

1667 PERIOPERATIVE CHANGES IN RENAL BIOMARKERS IN CHILDREN UNDERGOING OPEN HEART SURGERY

doi:10.1136/archdischild-2012-302724.1667

J Krastins, Z Straume, J Auzins. *Pediatric Intensive Care, University Children's Hospital, Riga, Latvia*

Introduction Cardiac surgery with cardiopulmonary bypass (CPB) is commonly perceived as a risk factor for decline in renal function. Hypothermia, hypoxia, hypotension, non-pulsatile blood flow during CPB, use of ACE inhibitors, inotropic and (or) vasoactive support affects kidney and contributes to the acute kidney injury (AKI).

Objective To evaluate dynamics of renal biomarkers - serum creatinine (SCr) and glomerular filtration rate (GFR) in children undergoing open heart surgery.

Methods We conducted prospective, non-randomized observational study at the tertiary care University Children's Hospital Pediatric ICU. We enrolled 30 patients, 12 boys and 18 girls with CHD. Their median body weight was 6.8 kg, (IQR 5.2–8.2 kg) and median