

(SIE); or as a further, poorly characterised form of AD skin peeling, termed epidermolysis bullosa simplex superficialis (EBSS), previously described in two families.

We report 6 affected individuals from 2 generations with generalised AD skin peeling. All presented neonatally with erosions at trauma-prone sites including the axillae, back and thighs, with ongoing skin fragility caused by friction. All 4 affected children reported slightly dry skin in the first decade of life with very mild hyperkeratosis of the axillae and neck. Some individuals had peeling of fingertips and soles, and one adult had mild diffuse plantar hyperkeratosis. There was no erythema, mucosal, nail or hair involvement. Initially, EBSS was considered based on AD inheritance, the generalised distribution, and lack of inflammation and ichthyosis at presentation.

A biopsy of rubbed, uninvolved skin from one affected individual showed a thickened stratum corneum but no signs of blistering or ultrastructural abnormalities at the dermal-epidermal junction or within the epidermis. Sequencing of *KRT5* and *KRT14* (keratins 5 and 14) showed no mutations, but whole exome sequencing demonstrated a heterozygous missense mutation in *KRT1* encoding keratin 1, p.Ser338Pro, in the 4 probands tested. This amino acid substitution is located within the L12 linker region, close to where other pathogenic mutations in keratin 1 have been reported in unrelated individuals with EI. Therefore, the most likely diagnosis in this family is EI due to a novel mutation in *KRT1*.

This clinically mild disorder and new *KRT1* gene pathology extends genotype-phenotype correlation in EI and underscores the value of next generation sequencing in diagnosing clinically atypical genodermatoses.

#### G29 A CASE OF PHYLLOID HYPOMELANOSIS – A RARE BUT SPECIFIC PRESENTATION OF CHROMOSOMAL MOSAICISM

doi:10.1136/archdischild-2013-304107.042

<sup>1</sup>L Solman, <sup>1</sup>M Glover, <sup>1,2</sup>VA Kinsler. <sup>1</sup>Paediatric Dermatology Department, Great Ormond Street Hospital, London, UK; <sup>2</sup>Clinical and Molecular Genetics Unit, UCL Institute of Child Health, London, UK

This 18 month old boy presented with a history of hypopigmented macular lesions, mild neurodevelopmental delay and recurrent episodes of acrocyanosis with tachycardia and increased tone. The cutaneous lesions developed from the age of 3 months, affecting the right trunk and right upper and lower limbs. Examination revealed large well-defined hypopigmented macules in a classical phylloid pattern, with a midline cut-off anteriorly and posteriorly on the trunk, compatible with a diagnosis of phylloid hypomelanosis. Mild facial dysmorphism was noted. Neurodevelopmental assessment at the age of 12 months suggested mild global delay; EEG and MRI of the CNS are pending. Cardiovascular examination was normal, however 24 hour ECG revealed non-specific ST segment changes. Ophthalmologic assessment was normal. Array comparative genomic hybridisation on a peripheral blood sample was normal. Karyotyping of affected and unaffected skin fibroblasts is underway.

Phylloid hypomelanosis (Greek phyllon = leaf, eidos = form) is characterised by congenital hypopigmented macules resembling a floral ornament, with round, oval or oblong patches (1), distinct from the commoner Blashko-linear distribution. It is a rare but highly specific sign of chromosomal mosaicism, universally associated thus far in the literature with mosaicism for duplications of 13q (1.2). Associated extracutaneous anomalies vary, and can include neurological, ocular, dental and skeletal defects (2). Cardiovascular abnormalities have not been reported thus far, and follow-up in our patient will clarify whether this is an associated or incidental feature. Affected individuals require multi-disciplinary assessment and long-term follow-up. As this is a somatic mosaic condition the possibility of fully affected offspring from the probands should be addressed at an appropriate age.

#### REFERENCES

1. Phylloid hypomelanosis is closely related to mosaic trisomy 13. Happle R. *Eur J Dermatol*. 2000 Oct-Nov; 10(7):511–2.
2. Phylloid hypomelanosis and mosaic partial trisomy 13: two cases that provide further evidence of a distinct clinicogenetic entity. González-Enseñat MA, Vicente A, Poo P, Catalá V, Mar Pérez-Iribarne M, Fuster C, Geán E, Happle R. *Arch Dermatol*. 2009 May; 145(5):576–8.

G30

#### DEVELOPING PAEDIATRICIANS AS FUTURE CLINICAL LEADERS: ENABLING DOCTORS IN QUALITY IMPROVEMENT AND PATIENT SAFETY (EQUIP) PROGRAMME DESIGN AND EVALUATION

doi:10.1136/archdischild-2013-304107.043

<sup>1</sup>J Runnacles, <sup>2</sup>L Linkson, <sup>2</sup>P Lachman. <sup>1</sup>Paediatrics, Kingston Hospital NHS Trust, Kingston-upon-Thames, UK; <sup>2</sup>Quality, safety and transformation, Great Ormond Street Hospital for Children NHS Foundation Trust, London, UK

**Aims** Paediatric postgraduate training needs to prepare paediatricians for the future delivery of high quality care. Doctors in Postgraduate Training (DrPGT) are often best placed to identify safety/quality concerns and can innovate across organisational boundaries. To address this, a programme was developed at a large tertiary centre providing a supportive educational environment. Its aims are to allow experiential learning on an improvement project alongside teaching of quality improvement (QI) and systems theory.

**Methods** EQUiP (Enabling Doctors in Quality Improvement and Patient Safety) supports DrPGTs through a QI project within their department, aligned to Trust's objectives. A three level approach to the programme ensures DrPGT engagement. All DrPGTs participate in a 1hr workshop to understand the importance of QI (level 1). Level 2 is a 6-month rotational programme with 2 full day workshops on improvement methodology, project surgeries facilitated by managers, and mentoring with senior clinicians. Level 3 is more intensive, over a 9 month period, to develop expertise and deliver level 1 workshops. The innovation involves a peer-designed programme while being work-based, delivering organisational strategies. Pre- and post-programme questionnaires allow Kirkpatrick 4-level evaluation.

**Results** All 40 participants agreed that the project was a valuable learning experience and that the programme met their expectations (level 1, reaction). Level 2, learning, was demonstrated by an improvement in QI definitions post programme, awareness of QI resources and confidence in using methodologies including PDSA and process mapping ( $P < 0.001$ ). Post programme, all but one participant said they are planning another QI project and that they are more aware of improvement work in their unit (behaviour change, level 3). Benefits to the organisation (level 4) are evident from successful projects presented to the executive team showing reduction in DNA rates, improved theatre efficiency, improved quality of medical notes etc.

**Conclusion** EQUiP changes the way DrPGTs view healthcare as they become quality champions for their department. The design and evaluation of EQUiP may inform similar educational programmes in other organisations. This capacity building is crucial to ensure future Paediatric leaders have the skills and motivation to improve the effectiveness of our healthcare system.

G31

#### USE OF TRANSLATED VERSIONS OF ROYAL COLLEGE OF PAEDIATRICS AND CHILD HEALTH (RCPCH) APPROVED PREM TOOL FOR PATIENT FEEDBACK IN AN ACCIDENT AND EMERGENCY DEPARTMENT DEALING WITH A MULTIETHNIC POPULATION

doi:10.1136/archdischild-2013-304107.044

C Singh, L Alsford. *Department of Paediatrics, North Middlesex University Hospital NHS Trust, London, UK*

**Aims** The aim of our study was to collect feedback by using the RCPCH PREM tool for paediatric urgent and emergency care (A&E),

and to try translated versions of the tool to maximise the participation of respondents with limited knowledge of English in a diverse ethnic population

**Background** Patient/parent satisfaction surveys are important monitoring tools used in the national health service (NHS). Before the Urgent-and-Emergency-Care PREM tool was published by the RCPCH in October 2012 there was no standard feedback form for paediatric A&E. This new tool is a well researched and standardised tool for obtaining children/parent feedback.

**Methods** Demographic data from the census was collected which showed that our NHS trust caters for a population with a large percentage of Turkish/Greek Cypriot, African and Somalian ethnicity. There is an annual attendance of about 38000 to children's A&E. While doing the survey, we eliminated the bias due to language-barrier by translating the RCPCH tool in Turkish and Somalian, the two commonly spoken languages in our ethnic population-group. The PREM tool was translated by doctors with knowledge of the languages and was colour coded for adult or children versions. Questionnaires were given to consecutive willing parents and/or children while waiting in the department. The feedback forms were analysed on Microsoft-Excel using common statistical methods.

**Results** Total of 50 feedback forms were collected. Feedback was given by 12(24%) children, 29(64%) parents and 4(9%) by both. The study group had 19(41%) European, 9(20%) Asian, 12(26%) African and 2(4%) mixed. The main languages spoken were English 25(57%), other European 10(22%), and all other 10(22%). The main highlights of the survey was that 43 of 46 (94%) respondents were satisfied by the services, 24(49%) waited longer than expected, 15(32%) wanted better information while waiting and 4(8%) were not given adequate privacy.

**Conclusions** The introduction of feedback-forms in multiple languages has perhaps given a more unbiased feedback with more involvement of the ethnic subgroups. The results were overall satisfying but a few specific areas that need improvement were identified. The responses to individual questions will also serve as a baseline for serial monitoring after implementing changes and training in problem areas.

G32

### DEVELOPING PRINCIPLES FOR DELIVERING AND COMMISSIONING BETTER HEALTH OUTCOMES AND EXPERIENCES FOR CHILDREN AND YOUNG PEOPLE SO THEY ARE COMPARABLE WITH THE BEST IN THE WORLD

doi:10.1136/archdischild-2013-304107.045

<sup>1,2</sup>M Clements, <sup>1,3</sup>K Evans, <sup>1</sup>T Cogan, <sup>1,4</sup>C Upton, <sup>1,5</sup>D Williams, <sup>1,6</sup>S Evans. <sup>1</sup>Strategic Network for Child Health and Well-Being, NHS East of UK, Cambridge, UK; <sup>2</sup>Department of Paediatrics, West Suffolk Hospital NHS Foundation Trust, Bury St Edmunds, UK; <sup>3</sup>Children & Young People's Programme, NHS Institute for Innovation and Improvement, Coventry, UK; <sup>4</sup>Department of Paediatrics, Norfolk and Norwich Hospital NHS Foundation Trust, Norwich, UK; <sup>5</sup>Department of Paediatrics, Addenbrookes Hospital NHS Foundation Trust, Cambridge, UK; <sup>6</sup>Child Health, Eastern Region Public Health, Cambridge, UK

**Aim** The development of a single set of principles to be used by providers and commissioners, across the whole healthcare system wherever a child or young person is seen. The aim is to improve the health outcomes and experiences for children and young people (CYP).

**Methods** A multi-professional team of GPs, Health Visitors, School Nurses, Paediatricians, Public health, Children's Nurses, patient and family representatives came together over a period of 6 months in a number of facilitated work-shops.

CYP and family engagement was central to this work with visits to Children's Centres, primary and secondary schools to listen to families experiences of health services. A graphic illustrator captured the key messages from each engagement event.

The principles were referenced against the Children and Young People's Outcome Forum Report and the NHS Mandate.


**Results** A set of 6 principles;

1. Child and Family focused
2. Health Promotion
3. Transformation
4. Settings
5. Information and Communication
6. Evidence Based and Sustainable

Each principle has an aspirational statement and then indicators to be used to evidence achievement towards a principle. They can be used to assess an existing service or to develop a new service and can be used for a condition across a pathway e.g asthma or for a service e.g. GP practise.

The principles have been developed as a single A5 poster for ease of use and are colourful and visual. A postcard has also been developed which summarises the feedback from young people and families but also translates the principles for families so that they know what they can expect from services. Figures 1, 2, 3.

Local organisations are encouraged to add their own and health-watch websites to the postcards to allow continuing feedback from families.



**Abstract G32 Figure 1** Stillbirths Figures Mbarara Hospital 2012