O-207

THE STATUS OF EARLY CHILDHOOD DEVELOPMENT OF "LEFT-BEHIND CHILDREN": A CROSS-SECTIONAL SURVEY IN RURAL CHINA

¹H Guan, ¹XW Tao, ¹ZY Fan, ²YR Zhao. ¹Early Childhood Development, Capital Institute of Pediatrics, Beijing, China; ²Department of Developing Pediatric, Shengjing Hospital China Medical University, Shenyang, China

10.1136/archdischild-2014-307384.277

Objective Approximately 58 million of children living in rural areas in China have been left behind after their parents migrated to cities to find work. The impacts of parental migration on the early development of left-behind children (LBC) are uncertain. We conducted a preliminary cross-sectional survey to investigate the prevalence of these left-behind children, their status of early childhood development and associated risk factors.

Methods A total of 1107 children were enrolled from two counties in Anhui province, China. Their caregivers completed questionnaires on demographics, the Early Childhood Development Questionnaire and the Parenting Self-efficacy Scale.

Results The estimated prevalence of left-behind children in the area was 86.6% (959/1107), comprising 30.8% (341/1107) and 55.8% (618/1107) children as a result of migration of either parent or both parents, respectively. The duration of breastfeeding for LBC was significant lower than those staying with mothers (8.74 vs. 10.14; $p < 0.01$). LBC had fewer reading books than non-left-behind children (NLBC) ($p < 0.05$). LBC also showed fewer times of physical check and higher case rate of diarrhoea in the last two weeks than NLBC. Male LBC who were fostered by caregivers with lower levels of parenting efficacy, and came from poorer families with less social support, experienced more mental health problems than other children.

Conclusions The study confirmed some negative outcomes for children who were left-behind in their early childhood. As the huge population proportion of LBC in rural China, more research on intervention strategies to improve the early development of LBC is urgently needed.

O-208

RANDOMISED CONTROLLED TRIAL OF DAY CARE VERSUS HOSPITAL CARE OF SEVERE CHILDHOOD PNEUMONIA AND SEVERE MALNUTRITION IN DHAKA, BANGLADESH

¹H Ashraf, ¹NH Alam, ²N Gyr. ¹Centre for Nutrition and Food Security, International Centre for Diarrhoeal Disease Research Bangladesh, Dhaka, Bangladesh; ²Faculty of Medicine, University of Basel, Basel, Switzerland

10.1136/archdischild-2014-307384.278

Background and aims Management of severe pneumonia and malnutrition relies on hospital-based treatment but practical barriers often prevent children in areas with highest disease burden from receiving hospital care. A RCT compared clinical and cost of Day Care Approach (DCA) of management with that of hospital management.

Methods Children aged 2–59 months with severe pneumonia with severe malnutrition received either DCA at outpatient clinic with antibiotics, micronutrients, diet and supportive treatment from 08:00–17:00 daily, while mothers were educated on continuation of care at home, or hospital care with similar 24-hours treatment.

Results Four hundred seventy children aged 12.2 (8.8) months were equally assigned randomly to either DCA or hospital care, 13% had hepatomegaly and 11% had hypoxemia. The durations of day care and hospital were 7.9 (5.5) and 7.1 (3.1) days, respectively. Successful management was possible for 184/235 [78.3% (95% CI 72.6–83.1%)] in DCA and 201/235 [85.5% (95% CI 80.5–89.5%)] in hospital. Thirty-six [15.3% (95% CI 11.3–20.5%)] in DCA and 22 [9.4% (95% CI 6.3–13.8%)] in hospital required referral to hospitals. There were no deaths among DCA, however, 2 [0.9% (95% CI 0.2–3.0%)] died in hospital. The average cost of treatment per episode per child was 165 US\$ for DCA and 256 US\$ for hospital.

Conclusions Severe pneumonia with severe malnutrition can be treated successfully and safely at reduced cost by DCA, but as effectively as hospital care, which can have an enormous impact in countries where recourses are scarce and hospital beds are limited for inpatient treatment.

ESPNIC – Young Investigators Presentations

O-209

CONTINUOUS INCISIONAL LIDOCAINE INFUSION IN PAEDIATRIC PATIENTS FOLLOWING OPEN HEART SURGERY

B Toledo, S Fernández, S Mencía, R González, E Sánchez, J López-Herce. *Pediatric Intensive Care Unit, Hospital General Universitario Gregorio Marañón, Madrid, Spain*

10.1136/archdischild-2014-307384.279

Study objective To evaluate the efficacy of local incisional lidocaine infusion after heart surgery.

Methods Observational prospective study including post cardiac surgery children in whom a 0.5% lidocaine elastomer was intra-operatively placed within the surgical wound. Lidocaine doses and plasma levels 48 h after administration, adverse effects, additional analgesic or sedative drugs and analgesia and sedation score were registered. All patients received intravenous metamizol and paracetamol.

Results 105 patients were analysed. Median age was 66 months, with 21% of patients under 2 years of age. The most frequent surgical interventions were: Fontan surgery (26.2%), atrial septal defects (20.4%) and ventricular septal defects (10.7%).

Median lidocaine administration duration was 58 h. 52.4% received fentanyl infusion (mean dose 1 mcg/kg/h, mean duration 24 h), 18.1% received midazolam infusion (1.5 mcg/kg/h, 8 h) and 12% received propofol (1.5 mg/kg/h, 4 h). Mean comfort score was 21.8 points and mean analgesia score was 1.9 (62% of patients required additional analgesia).

Fentanyl infusion was less often needed in patients over 2 years old than in younger patients (45.8% vs. 77.3%; $p = 0.015$). Midazolam infusion was also less often required in patients over 2 years old (13.3% vs 36.4%; $p = 0.025$).

Lidocaine plasma levels were recorded 48 h after administration in 66 patients, 46% ($n = 31$) were within therapeutic range (1.4–5 mcg/ml). No important side – effects attributable to lidocaine were observed.

Conclusions Incisional analgesia with lidocaine is safe and effective after cardiac surgery. Children over 2 years of age require less additional analgesia and sedation.