

effect is always lower than in singletons. Given the high prevalence of twins among preterm infants, this matter should be further investigated.

**PS-319 CORD BLOOD VITAMIN D STATUS AND NEWBORN BODY COMPOSITION**

<sup>1</sup>E Malchau Carlsen, <sup>2</sup>KM Renault, <sup>3</sup>V DeKnecht, <sup>4</sup>K Nørgaard, <sup>3</sup>D Cortes, <sup>5</sup>KF Michaelsen, <sup>5</sup>C Mølgaard, <sup>6</sup>O Pryds. <sup>1</sup>Department of Pediatrics, Hvidovre University Hospital University of Copenhagen, Hvidovre, Denmark; <sup>2</sup>Department of Obstetrics and Gynaecology, Odense University Hospital University of Southern Denmark, Odense, Denmark; <sup>3</sup>Department of Pediatrics, Hvidovre University Hospital University of Copenhagen, Hvidovre, Denmark; <sup>4</sup>Department of Endocrinology, Hvidovre University Hospital University of Copenhagen, Hvidovre, Denmark; <sup>5</sup>Department of Sports Exercise and Nutrition, University of Copenhagen, Copenhagen, Denmark; <sup>6</sup>Department of Sports Exercise and Nutrition, Hvidovre University University of Copenhagen, Hvidovre, Denmark

10.1136/archdischild-2014-307384.617

**Background and aims** Previous studies in newborns have found a positive association between adiposity and maternal and cord blood 25-hydroxy-vitamin D status (vit-D). It has been hypothesised that vit-D play a role in adipocyte metabolism and that this can be linked to obesity. Cord blood vit-D is closely associated with maternal vit-D concentration and is further affected by maternal obesity. The aim with our study was to examine the association between newborn cord blood vit-D and body composition where analyses were stratified by maternal pre-pregnancy obesity since this is a strong determinant of newborn body composition.

**Methods** Pre-pregnancy obese and normal weight mothers were included. Cord blood was collected at birth and newborn body composition was assessed using dual-energy X-ray absorptiometry within 48 h of birth. Sufficient cord blood and scans were obtained in 173(55%) mother-newborn dyads. Multiple linear regressions with vit-D was as a dependent variable were performed.

**Results** We included 41 pre-pregnancy normal weight-newborn dyads and 132 pre-pregnancy obese-newborn dyads. There was no difference in mean cord blood vit-D between offspring of normal weight and obese mothers (59.6 vs. 62.4 nmol/l (p = 0.64)). Vit-D was significantly associated with normal weight offspring body composition, positively with lean mass (p = 0.031) and inversely with fat% (p = 0.037) but was not associated with obese offspring body composition. See Table 1.

**Conclusion** We found an association between vit-D and newborn body composition in normal weight offspring. This implies that the effect of vit-D on newborn body composition may be determined by maternal weight.

**Abstract PS-319 Table 1** Association between newborn cord blood vitamin D (nmol/l) and maternal age and parity and newborn body composition in normal weight and obese mothers offspring

Dependant	Vitamin D (nmol/l)		Vitamin D (nmol/l)	
	Normal weight offspring (n = 41)	P- value	Obese offspring (n = 132)	P- value
Determinants	β*		β*	
Maternal age (years)	2.5	0.026	0.5	0.46
Primiparity (y/h)	34.5	0.004	9.8	0.10
Lean mass (g)	0.04	0.031	0.01	0.083
Fat (%)	-3.5	0.037	0.21	0.76

**Perinatology**

**PS-319a EXTREMELY LOW EXCLUSIVE BREAST FEEDING (EBF) RATE AMONG THE SYRIAN REFUGEE COMMUNITIES IN JORDAN**

G Fander, N Beck, H Johan, W Dymnt. *International Programm, Medair, Lausanne, Switzerland*

10.1136/archdischild-2014-307384.618

A total of 31,485 Syrian refugee households have been reached by the Medair nutrition programme covering 35% of the estimated number of refugees in Jordan urban areas. In April of 2014, a household survey was conducted to estimate exclusive breastfeeding (EBF) rates among those households reached by Medair with IYCF messages. 990 out of 31,485 households were selected by systematic random sampling for a telephone survey. Among those sampled, 12.9% (128 households) had children less than 6 months of age; 24.2 ± 7.7% of these reported practicing exclusive breastfeeding within the previous 24 h. This value is much lower than levels of EBF reported in Syria prior to the crisis (42.6%) despite effective IYCF promotion activity reflected by the fact that 71.3% (± 6.4%) of lactating mothers surveyed were able to state more than two benefits of EBF covered. Considering EBF is one of the most effective way to save the lives of young children, this deterioration in EBF among refugees placing young children at an increased risk of death should serve as a warning to the humanitarian community that measures to improve EBF is needed immediately. Additionally the survey implies other interventions are needed to address the other potential obstacles to EBF practices such as cultural and social barriers. Group sessions to monitor the barriers and promoters of IYCF as well as developing a self-supporting system among caregivers influencing the social and cultural aspects of EBF may be helpful to facilitate dissemination of lessons learned among the refugee communities.

**Pneumonia**

**PS-320 THE RELATION OF TRAFFIC LOAD TO TRAFFIC-DEPENDENT POLLUTANTS AND CROUP**

L Stanislavchuk. *Pediatric Infection Disease, Vinnytsya National Medical University N.A. M. Pyrogov, Vinnytsya, Ukraine*

10.1136/archdischild-2014-307384.619

**Objective** Some studies have compared the occurrence of croup with air pollution. Results have so far remained contradictory. Motor vehicles represent the principal source of air pollution in the city of Vinnytsya, Ukraine. The objective of this study was to determine the relation of traffic load to traffic-dependent pollutants and croup.

**Methods** Among a population of 8.067 children in residence near areas of high traffic density (>1.500 motor vehicles/ hour) and 2.473 children in residence near areas of low traffic density (<300 motor vehicles/ hour) cases of croup were registered by physicians during a 4 years period in 2000–03. Air pollution by sulfur dioxide (SO<sub>2</sub>), nitrogen dioxide (NO<sub>2</sub>), particulate matter, carbon monoxide (CO) were locally measured.

**Results** Areas with high traffic load are characterised by higher concentration of traffic-dependent pollutants and higher annual incidence of croup (Table).