replacement therapy which significantly increased the survival of these patients.

PO-0044 A REVIEW OF THE CLINICAL PRACTICE OF PAEDIATRICIANS WITH EXPERTISE IN CARDIOLOGY (PECS) IN THE UK

¹Y Singh, ²H Andrews, ¹W Kelsall. ¹Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK; ²College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK

10.1136/archdischild-2014-307384.719

Background Recently there have been significant developments in PEC training and service aims including the recognition of Paediatrician with Expertise in Cardiology Special Interest Group (PECSIG) by the RCPCH and British Congenital Cardiac Association (BCCA), the creation of an agreed SPIN-module training curriculum for PECs and the development of PEC-specific service standards by the BCCA.

Aim To find out which clinical services are provided by PECs and to determine the extent of variability in their clinical practice.

Methods An internet-based questionnaire was sent out via PEC-SIG and NICHe (Neonatologists with Interest in Cardiology and Haemodynamics) contact databases and an NHS directory. Non-responders were followed up via telephone.

Results The response rate was 72% (129 of 179 hospitals). PECs carried out echocardiography in all hospitals in which they were employed (69%). Support for this service was provided by echo-technicians in 36% of hospitals and by neonatologists in 27%.

PEC-led outpatient clinics were held at least fortnightly in 66% of hospitals. The mode duration of appointment for a new patient was 30 min (range <20 min to >45 min) while for the follow-up appointment the mode was 20 min (range <20 min to >45 min).

Telemedicine facilities were established in only 45% centres, where sharing echocardiogram images via PACS was used most commonly.

Conclusion There remains significant variation in PEC services especially regarding the frequency of clinics, the duration of appointments and telemedicine utilisation. It was reassuring to see a relatively high number of hospitals offering PEC-led echocardiography and other PEC service provision.

Abstract PO-0044 Table 1 PEC service provision

Type of service	Percentage of centres providing service
Paediatric echocardiography	87%
12-lead ECG	98%
24-hour Holter ECG	91%
24-hour ambulatory BP monitoring	74%
Long term external cardiac monitoring (e.g. cardiac memo)	56%
Exercise testing	47%
Other services*	4%

* a handful offered TILT test and fetal echocardiography facilities

PO-0045 A REVIEW OF THE AVAILABILITY OF PEC (PAEDIATRICIAN WITH EXPERTISE IN CARDIOLOGY) SERVICES IN THE UK

¹H Andrews, ²W Kelsall, ²Y Singh. ¹College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK; ²Addenbrooke's Hospital, Cambridge University Hospitals NHS Foundation Trust, Cambridge, UK

10.1136/archdischild-2014-307384.720

Background A national survey completed in 2008 reported that PEC services were available in only 35% of non-specialist paediatric cardiology hospitals in the UK. In 2012, the NHS *Specialised Services Commission* recommended that there should be an increased level of PEC services in all hospitals as part of their 'Safe and Sustainable' review.

Aim To determine the availability of PEC services in the UK.

Methods An internet-based questionnaire was sent out via PEC-SIG and NICHe (Neonatologists with Interest in Cardiology and Haemodynamics) contact databases and an NHS directory. Non-responders were followed up via telephone.

Results The response rate was 72% (129 of 179 hospitals). Of these hospitals, 69% had established PEC services with at least 1 PEC employed and 19% had two or more PECs. In all centres echocardiograms were performed by either PEC consultants or specialist paediatricians, with support from echo-technicians in 36%.

Local PEC-led outpatient clinics ran at least fortnightly in two-thirds (66%) of hospitals, whilst 63% of hospitals held out-reach clinics with a paediatric cardiologist from a specialist centre at least monthly. However, 12 of the hospitals without any PEC services reported that they never held out-reach clinics either.

Conclusion There has been a substantial increase in PEC availability in non-specialist paediatric cardiology hospitals (69% as compared to 35% in 2008) but still 31% had no established PEC services. In most but not all hospitals without PEC services, support is offered by tertiary-centres for paediatric cardiology through out-reach clinics.

PO-0046 CLINICAL AND ECHOCARDIOGRAPHIC EFFECTS OF HYPOCALCEMIA SECONDARY TO SEVERE VITAMIN D DEFICIENCY (VDD) AND EFFECT OF TREATMENT

A Soliman, S Saeed, A Khellah. *Pediatrics, Hamad Medical Centre, Doha, Qatar*

10.1136/archdischild-2014-307384.721

Introduction Hypocalcemia (HC), without an underlying myocardial disease, is a relatively uncommon but reversible cause of congestive heart failure.

Objective We studied the cardiac functions (heart rate, blood pressure, ECG and Echocardiographic parameters (Fractional shortening (FR), left ventricular end diastolic diameter (LVEDD) of 14 children who presented with hypocalcaemia due to VDD before and 2-4 weeks after treatment with an IM dose of vitamin D 3 (VD) (10,000 IU/kg).

Results Correction of HC and VDD was associated with marked improvement of the LVEDDS (3.2 ± 4.4 to 1.1 ± 2.8) and slowing of the heart rate (from 101 ± 34 to 94.7 ± 30/min). The FS and QTc did not change. The LVEDDS was negatively correlated with serum calcium level (r = -0.46, p = 0.03) and PTH concentrations (r = 0.44, p = 0.032) but not with 25OHD level (r = -0.2, p = 0.2).