groups were compared with regard to their epidemiological characteristics.

Results We identified 627 diabetic children. They were 332 boys and 295 girls. The incidence was estimated at 8.5/100 000 children under 15 years and 9.6/100 000 children (0–4 years). The incidence was 7.7/100 000 children (0–15 years) in 2009, it passed to 8.93/100 000 in 2011.

The patients were aged 0–4 years in 33% of cases, 5–9 years in 34.1% of cases and 10–15 years in 32.9% of cases.

The discovery of diabetes was in winter in 35% of cases. Parental consanguinity was noted in 31.2% of cases.

Conclusion Type 1 diabetes is a public health problem in Tunisia, its incidence increases and the age of diagnosis shifts to ages younger. Winter predominance of discovery supports the hypothesis of a triggering viral infection.

PO-0055 INSULIN PUMP IN CHILDREN WITH TYPE 1 DIABETES IN A REGIONAL HOSPITAL IN IRELAND

A Pavel, O Oyediji, J Jennings, J Chukwu. *Paediatrics, Our Lady of Lourdes Hospital, Drogheda, Ireland*

10.1136/archdischild-2014-307384.728

Introduction Initiation of insulin pump therapy in children with type 1 diabetes results in better glycaemic control, reduction in the short-term and long term complications and in better quality of life.

Objectives To determine the impact of insulin pump on glycaemic control (HbA1c), BMI and occurrence of severe complications in children with IDDM in a secondary care centre.

Methods A retrospective study of children with type 1 diabetes on insulin pump therapy for at least one year at the time of the study was conducted. HbA1c, BMI and frequency of severe complications one year before and after introduction of insulin pump were compared.

Results Twelve out of the thirty children (40%) on insulin pump therapy met our inclusion criteria. Their mean age at the time of the study was 12.6 years. Seven boys (58.3%) and five (42%) girls were studied. The mean duration of diabetes was 5.5 (± 2.2) years. The mean HbA1c before the introduction of pump therapy was 8.1% vs. 7.1% one year after; while the mean BMI z-score was 0.79 before and 0.88 after. Severe hypoglycaemia and DKA were noted in two children before but none after the initiation of pump therapy. The mean HbA1c decreased by 0.4% at 3 months (p 0.05) and by 1% at 2 months (p 0.013) of pump therapy.

Conclusions Initiation of insulin pump therapy results in significant reduction in the HbA1c within the first twelve months of therapy with a decrease in the frequency of occurrence of severe complications.

PO-0056 CLINICAL AND LABORATORY FINDINGS OF DIABETIC KETOACIDOSIS IN A PICU OF ALBANIA

¹I Bakalli, ¹E Celaj, ¹E Kola, ¹R Lluka, ¹D Sala, ¹I Kito, ¹I Gjeto, ¹S Sallabanda, ²E Klironomi. ¹PICU, UHC Mother Teresa, Tirana, Albania; ²Statistics, Albanian University, Tirana, Albania

10.1136/archdischild-2014-307384.729

Introduction Delay diagnosis is the major cause of Diabetic ketoacidosis (DKA). Children with profound acidosis are at great risk for symptomatic cerebral oedema.

Objective To identify the epidemiological profile, clinical feature, factors related to delayed diagnosis in children with DKA and to analyse the factors associated with prolonged acidosis.

Methods We analysed the records of all children with DKA, admitted to our PICU during January 2004–December 2013. We evaluated clinical features, biochemical profile at admission, 6, 12 and 24 hrs, presence of sepsis, shock, complications and outcome. The severity of DKA was defined by the degree of acidosis: mild ($pH = 7.2–7.3$), moderate ($7.1–7.2$) and severe ($pH < 7.1$). Anion gap (AG), delta gap (DG) and delta ratio were calculated. Prolonged acidosis was analysed against various independent factors.

Results Mean age of the patients was 7.06 ± 4.24 years, with misdiagnosis in 32% of cases. By the degree of acidosis, DKA was mild in 16%, moderate in 56% and severe in 28% of cases, with prolonged acidosis (>24 hrs) in 36% of cases. Factors associated with prolonged acidosis were: $Na > 133$ mEq/L ($p = 0.01$), $HCO_3^- < 4.8$ mEq/L ($p = 0.03$), $pH < 7.01$ ($p = 0.01$), $Cl^- > 100$ mEq/L ($p = 0.02$) and $AG > 25.1$ ($p = 0.03$). HbA_{1c}, azotemia, DG and misdiagnosis didn't result significant for prolonged acidosis. Three cases are complicated with cerebral oedema. Initial blood glucose or decline in glucose had no association with cerebral oedema. Mortality rate was 8%.

Conclusion Misdiagnosis of diabetes with DKA as consequence, is still high in children in Albania. Clinical and laboratory findings help identifying the patients who require a higher level of intervention.

PO-0057 ASSOCIATION OF DIETARY PATTERN WITH BIOCHEMICAL BLOOD PROFILES AND BODY WEIGHT AS RISK FACTORS OF CARDIOVASCULAR DISEASE AMONG ADULTS WITH TYPE 2 DIABETES MELLITUS

¹N Darani Zad, ¹H Esmali, ¹S Khalatbari, ²M Vaezi, ³M Hamedani. ¹Nutrition and Dietetics, Faculty of Medicine and Health Sciences Universiti Putra Malaysia, Serdang, Malaysia; ²Faculty of Health, Islamic Azad University-Tehran Medical Branch, Tehran, Iran; ³Faculty of Engineering, Islamic Azad University-Tehran, Tehran, Iran

10.1136/archdischild-2014-307384.730

Background and aims Nutrients are established as dietary risk factors for cardiovascular disease (CVD), but dietary patterns may be a better predictor of CVD risk. This study was conducted to identify dietary patterns and evaluated their association with biochemical blood profiles and body weight among 400 adults with type 2 diabetes mellitus aged between 40–60 years.

Methods Biochemical blood profiles, anthropometric measurements, and dietary data were obtained. Food frequency questionnaire were used to derive dietary patterns. Factor analysis was conducted to ascertain the dietary patterns, and analysis of covariance was fitted to assess the relation between blood profiles, body weight and adherence to dietary patterns.

Results Three dietary patterns by factor analysis were identified, Vegetable and Poultry, Western and Mixed. After control for potential confounders, waist circumference ($b = -0.12$, $p < 0.01$) and body mass index ($b = -0.15$, $p < 0.02$) were negatively associated with vegetable and poultry dietary pattern. Conversely, total cholesterol ($\beta = 0.14$, $p < 0.008$) and fasting blood glucose ($b = 0.12$, $p < 0.01$) were positively associated with western dietary pattern. A dietary pattern labelled as mixed pattern was found to be positively related to HDL-cholesterol ($b = 0.16$, $p < 0.002$) and body mass index ($b = -0.18$, $p < 0.01$). Associations between mixed pattern, LDL-cholesterol ($b = -$