

PO-0192 WITHDRAWN

PO-0193 WITHDRAWN

PO-0194 THE CLINICAL IMPACT OF POSITIVE BLOOD CULTURE RESULTS IN A DISTRICT GENERAL PAEDIATRIC UNIT

SE Lawson, PD Donnelly, C Mark, NP Corrigan. *Paediatrics, Altnagelvin Hospital, Londonderry, UK*

10.1136/archdischild-2014-307384.851

Background and aims Blood cultures are taken to detect bacteraemia and direct antimicrobial therapy. Uncertainty regarding contamination versus bacteraemia leads to financial and human cost through increased investigations and prolonged antimicrobial therapy¹. Contamination may be due to inadequate aseptic technique, with rates up to 10%².

This audit aims to establish bacteraemia rates, contamination rates and review if positive blood cultures change clinical management.

Methods A retrospective list of all paediatric blood cultures collected between 7/8/13–7/12/13 was compiled and case-notes of positive results reviewed using a designated proforma.

Contamination was suspected with multiple organisms, particularly if skin flora, exclusively aerobic/anaerobic isolates and reassuring inflammatory markers or clinical condition.

Results

339 blood cultures were taken, 19 were positive.

47% of positive results occurred in patients <1 year old.

The main indication was pyrexia (63%)

42% of cultures were collected on the ward

Documented aseptic non-touch technique in 1 case (5%)

Antibiotics were administered before collection in 10%

68% of positive results were felt to be contaminants

Our bacteraemia rate is calculated as 1.76%.

37% of all positive cultures were repeated including three of six confirmed bacteraemia cases.

A positive culture result altered clinical management in 32%.

Conclusions Bacteraemia rates are low at 1.76%. However, there is still a significant contamination rate of 68%, especially in infants where aseptic procedures are difficult. Our results confirm that a blood culture result in isolation is of limited value and clinical correlation is paramount.

REFERENCES

- 1 Bates DW, Goldman L, Lee TH. Contaminant blood cultures and resource utilisation. The true consequences of false positive results. *JAMA* 1991 265:365–369
- 2 Taking Blood Cultures – a summary of best practice. Saving Lives: reducing infection, delivering clean and safe care. Department of Health, London 2007. Accessed (14th January 2012) via Department of Health Website at: http://hcai.dh.gov.uk/files/2011/03/Document_Blood_culture_FINAL_100826.pdf

PO-0195 A PROSPECTIVE STUDY OF DENGUE FEVER PRESENTATION IN A CHILD COHORT

¹J Liao, ²L Foo, ³E Hennessy. ¹*Paediatrics, Aberdeen Royal Infirmary, Aberdeen, UK;* ²*Sungai Buloh Hospital, Chidren's Ward, Kuala Lumpur, Malaysia;* ³*Paediatrics, Aberdeen Royal Infirmary, Aberdeen, UK*

10.1136/archdischild-2014-307384.852

Background Dengue fever (DF) infects between 50 and 100 million people each year. Local DF transmission was first reported

in Europe in 2010 and WHO now warns of a possible DF outbreak in Europe.

DF is usually self-limiting in the population (overall mortality rate of <1%.) However, 90% of patients with dengue haemorrhagic fever (severe dengue) are under 15 years old. The initial presentation of both is similar in children. Mortality rate in severe dengue is 2.5%.

Aim To describe DF presentation in a child cohort

Method During an 8 week period, data were collected for 19 children diagnosed with DF and included demography, previous dengue infection, clinical presentation and time interval between symptom onset and admission.

Results All children with DF were under 15 years old. None had previous episodes, complications or progression to dengue haemorrhagic fever. 74% of children were living in red flagged areas when symptoms occurred. Close contacts with a recent DF diagnosis were identified in 37%. Pyrexia was a first symptom in 84%. Other symptoms included rashes (42%), cough (37%), loss of appetite (37%), vomiting (37%), dizziness (37%) and headache (37%), as well as 14 other symptoms of lower frequency. An average of 5 days separated initial symptoms and admission.

Conclusion In considering differential diagnosis in children presenting in Europe with non-specific symptoms such as those found in this study, the possibility of DF should not be omitted. A travel and contact history is even more important if this potentially serious infection is to be recognised.

PO-0196 EVOLUTION OF INFECTIONS IN FOUR CACHECTIC CHILDREN. CASE REPORTS

L Marinu, S Raciula. *Pediatric: Mother and Child, University of Medicine and Pharmacy of Craiova, Craiova, Romania*


10.1136/archdischild-2014-307384.853

We can diagnose cachexy in children whose BMI is <5 percentiles and under 3-rd negative deviation. The aetiology of poor state of nutrition can be: incorrect feeding, frequent infections, congenital defects, improper care or a combination of the previous situations.


Authors present four cachectic patients: two children suffering of trisomy 21, who had associated each one heart malformation: first, a boy, N. S., aged 7 and weighting 12 Kg had Fallot tetralogy and bacterial endocarditis; second, a girl, P. S, 43 months aged, weighting 10 Kg, had ventricular septal defect and acute interstitial pneumonia. The third cachectic patient was a girl aged 8, weighting 13 kg, D. C. with Seckel syndrome, admitted for staphylococcal pneumonia; the fourth is a 11 years girl weighting 12 Kg, P. D. with spastic tetraparesis, microcephaly, diagnosed with severe sepsis.

Only the boy suffering of endocarditis, evaluated to MODS and death; the three girls were discharged healed. Trisomy 21 caused immunodeficiency and cachexy also permitted severe infections. The boy had severe hypoxia, but girls had a normal oxygen saturation. The aetiology of poor state of nutrition was combined: congenital defects, improper diet, frequent infections, inadequate care (three patients lived in orphanage, a girl had only mother). Pneumonia is frequent in thin children, with birth malformations. Congenital heart defects could be complicated to endocarditis.

Cachexy in malformed children is an important cause of immunodeficiency which leads to severe, sometimes letal infections. Infections produce denutrition and cachexy promotes severe sepsis, especially in parentless children.



Blood culture (n=4)



NPA culture (n=30)

Abstract PO-0197 Figure 1 Bacteria detected by culture (Blood and NPA)

PO-0197 BACTERIAL CULTURE VERSUS PCR FOR ETIOLOGIC DIAGNOSIS OF COMMUNITY ACQUIRED PNEUMONIA- RESULTS FROM CAPES (COMMUNITY ACQUIRED PNEUMONIA ETIOLOGY STUDY)

¹JL Mathew, ¹S Singhi, ²P Ray, ³BV Ravi Kumar, ¹A Bansal, ⁴A Nilsson. ¹Advanced Pediatrics Centre, Post Graduate Institute of Medical Education and Research, Chandigarh, India; ²Medical Microbiology, Post Graduate Institute of Medical Education and Research, Chandigarh, India; ³Managing Director, Xcyton Diagnostics Pvt Ltd, Bangalore, India; ⁴Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm, Sweden

10.1136/archdischild-2014-307384.854

Background PCR technology holds promise to overcome the limitations of bacterial culture for confirming bacterial aetiology in childhood pneumonia.

Aims To compare the diagnostic yield using bacterial culture versus PCR in a cohort of children with community acquired pneumonia (CAP).

Methods CAPES is single-centre cohort study of 2000 consecutively enrolled children (1 month-12 years) with CAP (World Health Organisation definition). All underwent blood and

nasopharyngeal aspirate (NPA) culture for bacteria. In addition, a randomly selected sub-group (10%, n = 200) underwent multiplex PCR on NPA samples to identify eight pathogenic bacteria. **Results** In the sub-cohort of 200 children, pathogenic organisms were isolated from blood and NPA in 4(2%) and 30(15%) respectively (Figure 1). The main cohort showed similar proportions (1.7% and 14.0%) and distribution of organisms, suggesting absence of selection bias. Multiplex PCR (Figure 2) yielded traces of bacteria in 193(96.5%), with one organism detected in 44(22%), two in 76(38%), three in 53(26.5%), and four in 20 (10%). Etiologic diagnosis could be confirmed by bacterial culture in 4.5% cases, whereas PCR confirmed in 22%.

Conclusion Bacterial culture techniques appear to have limited sensitivity for etiologic diagnosis, whereas PCR has much higher sensitivity, although detection of multiple pathogenic bacteria precluded confirmation of aetiology in the majority.


PO-0198 FERRITIN AND SERUM IRON IN PAEDIATRIC FEBRILE ILLNESS

¹A Nasir, ²S James, ³D Perry, ⁴J Meza. ¹Pediatrics, University of Nebraska Medical Center, Omaha, USA; ²Emergency, Children's Hospital and Medical Center, Omaha, USA; ³Pathology, Children's Hospital and Medical Center, Omaha, USA; ⁴Statistics, University of Nebraska Medical Center, Omaha, USA

10.1136/archdischild-2014-307384.855

Introduction Ferritin is the major intracellular iron binding protein in the body. Serum ferritin is an inflammatory marker. Iron sequestration is one of the innate immune responses to infection. The goal of this research was to investigate the role of serum ferritin and serum iron as clinically useful markers of infection in the paediatric emergency department.

Methods Multiple inflammatory markers, including C-reactive protein, procalcitonin, and serum ferritin, and other iron studies were measured in 37 children, from 3 months through 8 years of age, presenting to the emergency department with temperature of ≥ 39 degrees Celsius, and 38 patients in the same age group with non-febrile illness (controls). Patients with chronic inflammatory or rheumatologic conditions and those with renal or hepatic failure were excluded.



Abstract PO-0197 Figure 2 Bacteria detected by multiplex PCR