O-060 ASSESSING AUTISM IN TODDLERS BORN

¹M Pritchard, ²T de Dassel, ³D Cartwright, ³F Bogossian, ⁴J Paynter, ⁵E Beller, ⁶S Russo, ⁷C Crothers, ⁸L Johnston, ⁹J Scott. ¹Perinatal Centre for Clinical Research, The University of Queensland Royal Brisbane Women's Hospital, Brisbane, Australia; ²Psychology, Royal Brisbane Women's Hospital, Brisbane, Australia; ³Nursing, The University of Queensland, Brisbane, Australia; ⁴Psychology, Autism Early Intervention Outcomes Unit, Brisbane, Australia; ⁵Centre for Research in Evidence-Based Practice, Bond University, Robina, Australia; ⁶Psychology, Better Life Psychology, Brisbane, Australia; ⁷Perinatal, The University of Queensland, Brisbane, Australia; ⁸Nursing, Queens University, Belfast, Ireland; ⁹Centre for Clinical Research, The University of Queensland, Brisbane, Australia

10.1136/archdischild-2014-307384.128

The Autism Diagnostic Observation Schedule (ADOS) is the only available semi-structured observational assessment of autism and has not been used extensively in preterm toddlers. We used a screening and diagnostic method to determine the diagnostic prevalence of autism and the characteristics of screen positive cases.

Methods A 12-month birth-cohort of toddlers (2-and-4-years-old) born with a gestational age <29-weeks were administered the Modified Checklist of Autism in Toddlers (M-CHAT)-and follow-up interview (FI™) screen, the ADOS and neurodevelopmental assessments (Bayley Scales of Infant and Toddler Development III-2-year-olds, Wechsler Preschool and Primary Scale of Intelligence-III and Adaptive Behaviour Assessment System-Second Edition-4-year-olds). The ADOS was conducted on toddlers with M-CHAT-FI™ positive screens.

Results Complete data were available on 88% (169/192) of children. Thirteen-percent screened M-CHAT-FI™ positive. ADOS (DSM-IV-criteria) classified 1.8% with autistic disorder and none with ASD. Multivariate analysis showed that social emotional delay ($p = <0.001$) was the only neurodevelopmental factor independently associated with M-CHAT-FI™ positive screens even after adjusting for psychosocial risk and child gender. All but one of the M-CHAT-FI™ positive screens scored some level of abnormality at the domain and item level on the ADOS testing, one third of these reached the Communication Total category cut off for ASD for the combined Communication and Social Interaction Total Score. Four children displayed comorbid atypical development (DQ/IQ <2SD).

Conclusion The ADOS reports a lower incidence of autistic disorder in children born very preterm compared to studies using diagnostic interview based tools. The distinguishing features of the M-CHAT-FI™ positive cases suggest a sub-threshold communication dysfunction profile.

O-061 CELL-BASED THERAPY FOR HYPOXIC-ISCHAEMIC INJURY IN THE PRETERM BRAIN

¹RK Jellema, ²TGAM Wolfs, ³V Lima Passos, ⁴A Zwanenburg, ¹DRMG Ophelders, ¹E Kuypers, ⁵AHN Hopman, ⁶J Dudink, ⁷HW Steinbusch, ⁸P Andriessen, ⁹WTV Germeraad, ¹⁰J Vanderlocht, ¹¹BW Kramer. ¹Department of Pediatrics. School for Mental Health and Neuroscience, Maastricht University, Maastricht, Netherlands; ²Department of Pediatrics. School of Oncology and Developmental Biology, Maastricht University, Maastricht, Netherlands; ³Department of Methodology & Statistics, Maastricht University, Maastricht, Netherlands; ⁴Department of Pediatrics. Department of Biomedical Engineering, Maastricht University, Maastricht, Netherlands; ⁵Department of Molecular Cell Biology. School of Oncology and Developmental Biology, Maastricht University, Maastricht, Netherlands; ⁶Department of Neonatology and Neuroscience, Sophia Children's Hospital, Rotterdam, Netherlands; ⁷School for Mental Health and Neuroscience, Maastricht University, Maastricht, Netherlands; ⁸Department of Pediatrics, Máxima Medical Centre, Veldhoven, Netherlands; ⁹Department of Internal Medicine Division of Haematology.

School of Oncology and Developmental Biology, Maastricht University Medical Center, Maastricht, Netherlands; ¹⁰Department of Transplantation Immunology Tissue Typing Laboratory. School of Oncology and Developmental Biology, Maastricht University Medical Center, Maastricht, Netherlands; ¹¹Department of Pediatrics. School for Mental Health and Neuroscience. School of Oncology and Developmental Biology, Maastricht University, Maastricht, Netherlands

10.1136/archdischild-2014-307384.129

Background and aims Preterm infants are prone to hypoxic-ischaemic encephalopathy. No therapy exists to treat this brain injury. The objective of this study was to assess the neuroprotective effect of exogenous administration of stem cells and the mobilisation of endogenous stem cells in the ovine preterm brain after global hypoxia-ischemia.

Methods Instrumented preterm sheep were subjected to global hypoxia-ischemia by 25 min of umbilical cord occlusion at a gestational age of 104 (term is 150) days. During a 7 day reperfusion period all vital parameters, including (amplitude-integrated) electroencephalogram, were recorded. At the end of the experiment, the preterm brain was assessed by histology and diffusion tensor imaging (DTI).

Results Systemic administration of exogenous mesenchymal stem cells (MSCs) reduced cerebral inflammation (i.e. microglia proliferation) and white matter injury. MSCs induced T-cell tolerance, which was paralleled with diminished mobilisation and invasion of these cells in the preterm brain. In addition, MSCs decreased number of seizures after global hypoxia-ischemia, indicating functional improvement.

Similarly, mobilisation of endogenous stem cells using systemic granulocyte-colony stimulating factor (G-CSF) reduced cerebral inflammation and white matter injury. However, G-CSF did not reduce the number of seizures after global hypoxia-ischemia.

Conclusion We have shown for the first time in a translational animal model that cell-based therapy is effective in protecting the preterm brain against the cerebral and peripheral inflammatory responses which are involved in the aetiology of white matter injury in the preterm brain after global hypoxia-ischemia. Our studies form the basis for future clinical trials studying feasibility of cell-based therapy in preterm infants with hypoxic-ischaemic encephalopathy.

Neonatal Brain and Development – Evolving Techniques

O-062 EARLY BRAIN ACTIVITY AND CORTICAL DEVELOPMENT IN PRETERM INFANTS

¹ML Tataranno, ²P Moeskops, ³LS de Vries, ³KJ Kersbergen, ³F Groenendaal, ³LGM van Rooij, ³MC Toet, ¹G Buonocore, ²I Isgum, ³MJNL Benders. ¹Department of Molecular and Developmental Medicine, University of Siena, Siena, Italy; ²Image Sciences Institute, University Medical Center Utrecht, Utrecht, Netherlands; ³Department of Neonatology Division of Perinatology, University Medical Center Utrecht, Utrecht, Netherlands

10.1136/archdischild-2014-307384.130

Background and aim Early brain activity is crucial for neuronal growth. It is well known that the cerebral cortex develops rapidly in the last trimester of pregnancy. We investigated whether early brain activity was related to the rate of cortical