

distributed in major medical areas. Paediatrics includes a mandatory requirement of this stage presentation Session Paediatric Clinics (SPC) based in the various practice settings in order to instigate the search for knowledge providing a meaningful construction.

Methods A descriptive study, retrospective documentary of SPC presented in 2013 by the undergraduate students of the Faculty of Medicine of Petrópolis, Rio de Janeiro, Brazil.

Results There were 70 SCP, 70% from the paediatric ward, 12.86% of the NICU and 17.14% of the other scenarios. Active participation in the choice of topic, review and submission process was on average 3 students and the number of listeners was 40/SCP. The chosen themes, 52,86% are not part of the curriculum previously offered. Infectious diseases have contributed to 37,14% of realisation of the SPC.

Conclusion This study shows for the promotion of proactive methodologies as supporters of the integration of the student as the protagonist of the teaching- learning process and therefore should be encouraged. As for the themes chosen believe that awakening to the search for new knowledge has been significantly.

PO-0968 MAIN DEATH ETIOLOGIES OF CATALAN CHILDREN. EXPERIENCE OF THE ADMITTED PATIENTS IN A TERTIARY HOSPITAL


¹S Navarro, ¹MR Escobar, ¹S Ciprés, ¹M Palomares, ²G Gelabert, ³JJ García. ¹Pediatric Palliative Care Unit, Hospital Sant Joan de Déu, Barcelona, Spain; ²Clinical Archives and Documentation, Hospital Sant Joan de Déu, Barcelona, Spain; ³Pediatrics, Hospital Sant Joan de Déu, Barcelona, Spain

10.1136/archdischild-2014-307384.1587

Background and aims Paediatric palliative care is an essential aspect of medical practice for patients who need end-of-life attendance. The better understanding of the main death causes allows to anticipate the future complications in the final stage. The aim is to describe the epidemiology and characteristics of deaths at childhood in Catalonia and specifically in a tertiary paediatric hospital.

Methods Review data from the National Statistics Institute (<http://pestadistico.inteligenciadegestion.mssi.es>) on mortality of people aged 0–19 years old, during the period from 2007 to 2011. Analyse the main general causes of death by ICD (International Code of Disease) collected in death certificate and compare them with our experience.

Results During the period from 2007 to 2011, a total of 2.282 deaths were registered in Catalonia (59% males, 41% females). The mortality rate varies by age (Graph 1). In Catalonia the most frequent causes of death were conditions originated in the perinatal period (COPP) (27%), external causes (18%),



Abstract PO-0968 Graph 1 Numbers of deaths in catalonia/100.000 population (2007–2011)

congenital malformations (15%) and neoplasms (11%)(Table 1). The epidemiological study in our centre shows that 487 patients died in this period. The mean age was 3.5 years. According to sex, 47% were female and 53% male. The average length of hospitalisation was 12 days (range: 1–167 days). In our hospital, the most frequent causes of death in inpatients were COPP (45%), diseases of the respiratory system (17%) and neoplasms (15%)(Graph 2).

Conclusions In accordance with the literature, during the first year there is a peak in the mortality rate. In this period the main causes of death are COPP and congenital malformations. In adolescence the main causes of death are external causes and malignancy. In our hospital, almost a half of the total deaths occur in the first month of life due to COPP. Every centre should know his epidemiology of the main causes of death.

PO-0969 WITHDRAWN

PO-0970 A NEW GROWING PAINS DIAGNOSTIC TOOL: EVALUATION IN A MEDITERRANEAN CLINICAL SAMPLE

¹M Vassilopoulou, ²A Spathis, ²I Paspatis, ³M Tsolia. ¹PICU, Penteli's Children Hospital, Athens, Greece; ²Orthopedic Clinic, Penteli's Children Hospital, Athens, Greece; ³2nd University Pediatric Clinic, University of Athens Medical School, Athens, Greece

10.1136/archdischild-2014-307384.1588

Background "Growing pains (GP)" is the most common musculoskeletal complaint in childhood.

Aim To investigate the sensitivity and specificity of a previously validated questionnaire, for the diagnosis of GP.

Methods From 01/2013–12/ 2013, a questionnaire (Tb. 1), was administered to parents of children aged 3–8 years, who visited an orthopaedic clinic of a general children's hospital, as outpatients, complaining of lower limb pain of no apparent traumatic

Abstract PO-0968 Table 1 Causes of death in Catalonia 2007–2011

			0–1 y.o.	2–4 y.o.	5–9 y.o.	10–14 y.o.	15–19 y.o.	TOTAL
II	C00–D48	Neoplasms	15	50	59	53	68	245
IV	E00–E90	Endocrine, nutritional and metabolic diseases	40	24	6	7	12	89
VI	G00–G99	Diseases of the nervous system	59	32	10	23	35	159
X	J00–J99	Diseases of the respiratory system	20	20	10	9	15	74
XVI	P00–P96	Certain conditions originating in the perinatal period	624	1	0	1	0	626
XVII	Q00–Q99	Congenital malformations, deformations and chromosomal abnormalities	284	28	8	11	8	339
XVIII	R00–R99	Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified	93	26	6	12	18	155
XX	V01–Y98	External causes of morbidity and mortality	23	59	43	46	237	408
		Others: Infectious, diseases of the blood, digestive and circulatory system...	46	46	22	31	42	187