Causes of Mortality 2007-2011 in our center

Abstract PO-0968 Figure 1 Causes of mortality 2007-2011 in our centre

cause. An orthopaedic, blind to the parents' answers to the questionnaire, evaluated the children. A researcher estimated the score of each one of the completed questionnaires and reviewed the orthopaedics' clinical chart.

Results 35 questionnaires were completed. According to orthopaedics, 21 children had GP, while according to questionnaire, 13 children did so. When the cut-off point was reset at 7, sensitivity was 91.3% and specificity 95.4%.

Conclusions The questionnaire quantifies parental answers and may assist clinicians in GP diagnosis.

Abstract PO-0970 Table 1 Questionnaire

Select the answer that best describes your child's limb pain attack:

- Occurs during late afternoon or evening.
- Is still present next morning.
- Is bilateral.
- Occurs always in the same limb.
- Is located in muscles (thigh, calf, posterior knee, foot)
- Resolves spontaneously or with massage of the affected area.
- Is persistent and doesn't resolve.
- The child awakes at night because of pain.
- The child is otherwise well.

YES to questions 1, 3, 5, 6, 8, 9 and NO to questions 2, 4, 7 are rated 1 and are indicative of GP.

The cut-off value for the diagnosis had been set at ≥ 8 .

PO-0971 MEDICAL STUDENT, AN AUDIT AND CLINICAL GOVERNANCE

¹C Glancy, ²S Mullen. ¹Paediatrics, Craigavon Area Hospital, Belfast, UK; ²Neonatal ICU, Royal Maternity Hospital, Belfast, UK

10.1136/archdischild-2014-307384.1589

Aims To assess if paediatric undergraduate students can improve their knowledge and understanding of clinical governance when undertaking an audit as part of their clinical placement.

Methods The students were broken into 2 groups (2-3 in each group). Each group was given a topic to assess (growth charts and post-take ward round documentation) and asked to design a

prospectus. A 10 chart audit was undertaken and the results presented to the clinical team. This project will run from August 2013 until Jan 2014.

A questionnaire was filled out at the start and at the end of the placement and used as an indicator of the student's progress. The questionnaire comprised of yes and no answers as well as a rating scale from 1-5.

Results No student had undertaken an audit or quality improvement project previously (n = 17).

The students rated their undergraduate teaching in clinical governance a mean mark of 1.6 (self-rating scale marked 1-5).

There was an increased in mean score regarding their understanding of clinical governance (pre 2, post 3.2), understanding of the audit cycle (pre 1.9, post 3.9) and in the importance of clinician governance in modern medicine (pre 3.5, post 4.3).

All students (100%) felt this project helped improve their c.v. and that it would be beneficial for all undergraduates to participate in an audit.

Conclusions The results identified a self-rated improvement in knowledge of clinical governance and the audit cycle as well as support for undergraduates undertaking a quality improvement project.

PO-0972 LONGITUDINAL RELATIONSHIP BETWEEN NURSES' AND PHYSICIANS' PERCEIVED APPROPRIATENESS OF CARE AND MORAL DISTRESS IN A NEONATAL INTENSIVE CARE

¹J de Boer, ²J van Rosmalen, ¹KM Reiss, ³M van Dijk. ¹Pediatrics/NICU, Erasmus MC Sophia Children's Hospital, Rotterdam, Netherlands; ²Biostatistics, Erasmus University Medical Centre, Rotterdam, Netherlands; ³Pediatrics/PICU, Erasmus MC Sophia Children's Hospital, Rotterdam, Netherlands

10.1136/archdischild-2014-307384.1590

Background/aims Invasive treatment without sufficient benefit induces doubts among nurses and physicians about 'appropriateness of patient care'. Conflicting interpersonal moral convictions, may cause moral distress. Additional sources of moral distress are incompatible institutional requirements, workplace

deficiencies and insufficient/unsafe levels of staffing. The associated declining self-respect, frustration, and guilt cause work-dissatisfaction, burnout, turnover, and consequently, diminished patient care. In this survey study, perceived appropriateness of care and levels of moral distress were evaluated across time. We also determined if respondents' background predicted moral distress levels.

Methods After baseline assessment (background, moral distress, ethical climate), nurses and physicians of our level-III NICU evaluated day-levels of perceived appropriateness of care, the different aspects of moral distress, and ethical climate, at the end of five randomly selected shifts.

Results Response rate: nurses 87(77%)/physicians 30(91%). Moral distress (range 1–16) was low at baseline ($M = 2.21$; $SD = 1.55$), but significantly higher for nurses than for physicians ($M = 2.40/SD = 1.68$ vs $M = 1.68/SD = 0.98$; $p = 0.01$). Nurses were less likely to disagree with treatment than physicians ($OR = 2.62$, $p = 0.02$). Moral distress at day-level (range 1–4) was very low ($M = 0.08/SD = 0.21$) and significantly depended on being religious ($\beta = 0.16$; 95% CI = 0.03 to 0.28) and perceived 'overtreatment' ($\beta = 0.18$; 95% CI = 0.07 to 0.30), contrary to 'undertreatment' ($\beta = 0.17$; 95% CI -0.63 to 0.29). Highest scores were observed for the following aspects of moral distress: provider (dis)continuity, communication about patient care, and (un)safe levels of staffing.

Conclusion In earlier studies, 'expressing concerns' and 'facilitated ethics conversations' proved to diminish moral distress. Possibly in our NICU the existence of structured-multi-disciplinary-medical-ethical-decision-making explains the (very)low levels of moral distress.

PO-0973 LIFESTYLES, EATING AND ACTIVITY FOR FAMILIES (LEAF) PROGRAMME: DEVELOPING A TIER 3 INTERVENTION FOR WEIGHT MANAGEMENT IN THE EARLY YEARS

N. Sauven, R. Brandreth, K.E. Laity, H. Vickerstaff. *Paediatrics, Royal Cornwall Hospital Trust, Truro, UK*

10.1136/archdischild-2014-307384.1591

Aim To develop a Tier 3, family-focused weight management intervention for the early years.

Background In the UK roughly 1 in 4 children are overweight or obese by the time they start primary school. According to the literature there are no proven models for working with children under 6 years with extreme obesity. We were required to develop a Tier 3 weight management intervention as part of the Care Pathway for Children's Weight Management in Cornwall.

Methodology A review of national guidance and both nationally and locally run programmes for children's weight management was undertaken. The key aspects for successful interventions, appropriate for the early years, were identified. A pilot programme was then developed by our team.

Results Successful programmes were already running for children aged 7–13 years. The LEAF (Lifestyles, Eating and Activity for Families) clinic was developed for children 6 years and under. It comprises a multi-disciplinary clinic with Community Paediatrician, Specialist Children Dietician and Specialist Activity Advisor, followed by a group intervention in a community setting, before multi-disciplinary follow-up. The intervention covers a broad range of topics that aim to help empower parents to make changes that ultimately improve the body mass index of their children. After completion of the programme families are

discharged, with the specialist team providing support to primary care professionals. Co-morbidities identified including raised LFTs, deranged lipids and sleep apnoea have been improved with this intervention.

Conclusion Initial results are promising but full evaluations of the outcomes of our programme are needed.

PO-0974 CARE OF CHILDREN AND YOUNG PEOPLE WITH DELIBERATE SELF-HARM: ARE WE DOING IT RIGHT?

L. Wajid, M. Aziz, H. Bentur. *General Paediatrics, Whiston Hospital, Liverpool, UK*

10.1136/archdischild-2014-307384.1592

Background and aim Deliberate Self-harm (DSH) affects 1 in 15 young people and remains one of top 5 causes of acute hospital admissions in UK. We looked at children attending hospital with DSH to assess their pathways of care and discharge outcomes. We also looked to see if any trends emerged in their attendance to hospital.

Methods Retrospective review of patients over 6 month's period from March to August 2012.

Results 135-patients were identified, 124 with diagnosis of self-harm and 11 with previous mental health problems. Following their attendance to emergency department 90% (n122) needed inpatient assessment and 96% of these were followed-up by mental health team. Average length of stay was 3 h to 6 days. Thoughts of harm included 81% (n100), drug overdose 50% (n62) and physical injury 37% (n46). Analgesics were most common drug overdose in 26% (n29) and medical interventions was needed in 19% (n12). 61% (n83) were girls. Social factors like parental-separation was seen in 61% (n82), living in supported accommodation in 26% (n35), family history of mental illness in 23% (n31) and social-services involved in 38% (n52).

Conclusion We found that most of our patients had significant DSH needing inpatient management (5% of our admissions) and majority needed follow-up with mental health services which reinforces the need to work closely with mental health team. Social risk factors were commonly found in our cohort. Recognition and early intervention may reduce the incidence of self-harm in these children. Analgesics were the most common drug of overdose reflecting the easy availability of these to children.

PO-0975 KNOWLEDGE, ATTITUDE AND BARRIERS TOWARDS RESEARCH IN THE NATIONAL HEALTH SERVICE (NHS)

¹M. Ahmed, ²S. Boswell, ²C. Lawrie. ¹Paediatrics, Burton Hospitals NHS Foundation Trust, Burton on Trent, UK; ²Research and Development, Burton Hospitals NHS Foundation Trust, Burton on Trent, UK

10.1136/archdischild-2014-307384.1593

Background In the 21st century, healthcare professionals know a great deal about health, disease and medicine. However, much remains uncertain and continued investment in research aims to find better ways of managing patients, improving public health, reducing health inequalities and keeping people in a state of good health. Clinical Research Networks are well established across the NHS to ensure that all patients and clinicians are able to share the benefits of participating in clinical research. Barriers to research can deter healthcare professionals from contributing towards useful research projects.