Abstract PO-0975 Table 1

	Strongly agree	Agree	Unsure	Disagree	Strongly disagree
Research is imperative to the future of the NHS	35	60	2	2	1
I am aware of the research studies carried out locally in my working area	9	37	23	25	6
I am aware of the requirement of Good Clinical Practice course to participate in research	18	32	16	30	4
Time commitment puts me off research.	14	40	14	28	4
Bureaucracy (paper work) puts me off research.	10	44	13	30	3
Protected time would enable me to carry out more research.	18	60	13	9	0
I am aware of the local research team and how they can support me undertaking research	7	46	13	27	7
Research is driven for publication and self-interest	3	22	20	49	6
Research is driven for the benefit of future patient management	29	56	13	0	2

Aim To ascertain knowledge and attitude towards research and identify barriers amongst healthcare professionals in a district general hospital setting.

Methods All the healthcare professionals were emailed a questionnaire via 'survey monkey' in March 2014. An email reminder to non-respondents was sent 2 weeks later. Questions focussed on participant's knowledge and attitude towards research and identifying perceived barriers to undertaking research.

Results 87 responses were received (26 doctors and 61 allied health professionals). Following table outlines key study findings in %.

Conclusions Majority of participants perceived time commitment and bureaucracy as barrier to research. Dedicated time allocated in the job plan and administrative support would be imperative in improving research outcomes leading to innovative treatments that can significantly improve patient health in future.

PO-0976 PRESCRIBING IN THE ELECTRONIC AGE: FASTER, SAFER, BETTER?

¹P Dale, ²P Munyard. ¹Pharmacy, Royal Cornwall Hospital, Truro, UK; ²Child Health, Royal Cornwall Hospital, Truro, UK

10.1136/archdischild-2014-307384.1594

Background and aims In December 2012 electronic prescribing (EP) and administration of medicines was introduced to the Trust (800 beds, 40 paediatric beds, 3 Paediatric HDU beds, 20 admissions/day) as a pilot – specifically to the Child Health Department – before general adoption across the Trust.

Methods Pre-implementation, training was thoroughly and carefully organised for 40 paediatricians, 60 nurses and 5 pharmacists. Training was face to face and then on-line with face to face support. Mobile computing devices were distributed to the ward areas. All in-patient paper prescriptions were transcribed to electronic systems and paper prescriptions were removed. Ward based 24 h pharmacist/nurse support was available for the first week.

Results All prescriptions were legible.

Antibiotic stewardship easily audited – prescriptions with stop/review date 50%/indication 33% vs. 86%/80% respectively after an educational programme in child health.

No paediatric patients have been given a medication to which they were known to be allergic after implementation of EP.

Dispensary error rates from electronic prescription orders reduced from 5.38/month to 1.5/month post-implementation.

Only 25% of paediatric staff would go back to paper prescriptions.

The system is now adopted across most of the Trust.

Conclusions Departmental involvement in planning was most important as was “buy-in” from the junior doctors and nursing staff. Well organised training for all staff, intensive and face-to-face, is felt to be essential. EP has been accepted and is a safer system which enables audit of practice down to individual levels. Outpatient use is planned in the future.

PO-0977 WITHDRAWN

PO-0978 POPULATION-BASED STUDY OF DOWN SYNDROME PREVALENCE IN SOUTHERN THAILAND

¹S Jaruratanasirikul, ¹O Kor-anantakul, ²W Limpitkul, ³P Dissaneevate, ³S Sutthibenjakul, ⁴N Khunnarakpong, ⁴N Intharasangkanawin, ⁵A Sattapanyo, ⁵S Pathompanitrat. ¹Pediatrics, Prince of Songkla University, Songkhla, Thailand; ²Pediatrics, Medical Education Center Songkhla Hospital, Songkhla, Thailand; ³Pediatrics, Medical Education Center Hatyai Hospital, Songkhla, Thailand; ⁴Pediatrics, Medical Education Center Trang Hospital, Trang, Thailand; ⁵Pediatrics, Phatthalung Hospital, Phatthalung, Thailand

10.1136/archdischild-2014-307384.1595

Introduction Down syndrome (DS) is the most common chromosomal disorder that causes mental retardation. In Thailand, studies on DS from 1969–1978 and 1988–1999, both hospital-based, found prevalences of DS of 0.89 and 1.07 per 1,000 live births.

Objective To determine the prevalence of DS in the population-based birth defect registries in 3 provinces in southern Thailand (Songkhla, Trang, Phatthalung), and the proportion of termination of pregnancy (ToP) of DS fetuses detected by prenatal screening.

Method Data were obtained from a population-based surveillance study undertaken during 2009–2012. Entries in the birth defects registry included all live births, all stillbirths, and all ToPs following a prenatal diagnosis. Infants with clinical characteristics of DS had a chromosomal study to make a definite diagnosis.

Results Of the total 148,795 births recorded during the study period, 180 DS cases were listed, giving an average prevalence of 1.21 per 1,000 births. The median maternal age was 36.4 years with percentage of maternal age ≥ 35 years of 52.2% (94/180). Fifty-three cases (29.4%) diagnosed prenatally resulted in termination of pregnancy.