and removal of placental suppressors when clamping the cord. The locus coeruleus is activated causing the arousal.

A spontaneous resting state activity has been found in newborn infants with fMRI.² This activity may correspond to the idea of William James that there is a "stream of consciousness". It involves five hubs including the somatosensory system and the auditory and visual cortex in the infants. This is in contrast to adults where ten hubs were defined including the insula, precuneus and ventromedial prefrontal cortex. Thus the infant is probably only aware of what it feels, sees and hears in present time, while the adult relates the sensory input to memories, itself and also plans for the future.

REFERENCES

- 1 <http://www.nuffieldbioethics.org>.
- 2 Fransson P, Skold B, Horsch S, et al. Resting-state networks in the infant brain. *PNAS* 2007;104(39):15531–36

IS-020 FUNCTIONAL CONNECTIVITY IN THE INFANT BRAIN

CD Smyser. *Neurology and Pediatrics, Washington University, St. Louis, USA*

10.1136/archdischild-2014-307384.20

Advanced MRI techniques have been increasingly applied in infants to explore the structural and functional architecture of the developing brain. Functional connectivity magnetic resonance imaging (fcMRI) utilises spontaneous, low-frequency, coherent fluctuations in blood oxygen level dependent signal to identify networks of functional cerebral connections. Application of fcMRI in infants provides unique technical challenges. To obtain high-quality fcMRI data, investigators have applied new technology and modified acquisition practices. Advanced analysis techniques have also been developed to improve anatomic registration, eliminate artifactual variance and improve signal-to-noise ratios. These measures have enabled successful, robust fcMRI investigations in neonates. Importantly, these methods are transferrable across institutions and clinical populations of interest.

Neonatal fcMRI investigations have included healthy, term-born infants and prematurely-born infants with and without cerebral injury. In these populations, fcMRI data has been used to identify immature networks as early as 26 weeks postmenstrual age. These networks gradually mature. Prematurity (Figure 1) and white matter injury (Figure 2) significantly affect connectivity, altering network configuration and strength. These results demonstrate the promise of fcMRI as an investigational tool of neurodevelopment, providing insight into the earliest forms of functional cerebral development. While key groundwork has been laid, additional efforts are necessary to apply continued

Abstract IS-020 Figure 1 EEG recording of a preterm baby at 31 weeks of gestational age using clinically approved EEG cap. Support vector machine multivariate pattern analysis results illustrating connections important for differentiating healthy, term-born infants and very preterm infants scanned at term equivalent postmenstrual age. Caliber of connections weighted by difference magnitude. Vectors colored green are those stronger in term infants, while orange vectors are stronger in very preterm infants

Abstract IS-020 Figure 2 Individual fcMRI correlation maps illustrating Fisher z-transformed correlation coefficients ($z(r)$; threshold = 0.3) overlaid on subject-specific, atlas-registered T2-weighted images. Results for very preterm infants scanned at term equivalent postmenstrual age with moderate and severe white matter injury included. Results for healthy, term-born subject provided for comparison. Maps were generated using an ROI located in the hemisphere of greater injury

advances in technology and methodology. Expanded investigations will provide greater understanding of the processes underlying typical and atypical cerebral development and the role of these networks in neurodevelopmental outcomes.

IS-021 FUNCTIONAL MRI AND LANGUAGE DEVELOPMENT IN INFANTS AND CHILDREN

I Krägeloh-Mann, K Lidzba. *Pediatric Neurology, University Children's Hospital, Tübingen, Germany*

10.1136/archdischild-2014-307384.21

Language is specific to the human and can, thus, not be studied in the animal. In the adult, there is a strong dominance of the left hemisphere for most aspects of language and stroke in language regions often leads to permanent aphasia. In contrast, lesions in similar locations acquired very early do not lead to an impaired language function in the affected child.

With the advent of fMRI, the cerebral representation of language organisation can now be studied non-invasively even in smaller children.

Questions

1. Language representation during development?
2. Early left hemispheric lesions and language representation?
3. If there is language reorganisation, does it affect right hemispheric functions?
4. How good is right hemispheric language?
5. What is the time frame for reorganisation?

Answers

1. Language representation is initially bilateral and increasingly left dominated during development.

2. Early left hemispheric lesions may induce reorganisation to the right – in homotopic areas.

3. This happens on the expense of right hemispheric functions.

4. Language quality in reorganised language is not impaired for everyday language, but there are differences in complex linguistic aspects, which supports the idea of a genetic predisposition.

5. The time frame for reorganisation is not very clear, but there is no evidence for successful reorganisation after preschool age.

This work is supported by the German Research Council (DFG) and the University of Tübingen.

The Child

IS-022 CRITICAL CARE FOR THE CHILDREN OF THE WORLD

N Kissoon. *Global Child Health, BC Children's Hospital and University of British Columbia, Vancouver, Canada*

10.1136/archdischild-2014-307384.22

The burden of sepsis in the developing world Early childhood and neonatal sepsis has become a global public health concern. Nearly seven million children die each year before reaching their fifth birthday and around 50% of these deaths are attributable to sepsis and possible serious infections.

Studies have demonstrated the effectiveness of community interventions in reducing neonatal deaths due to sepsis and other severe diseases. But in resource poor settings, the lack of access to health facilities, acute shortage of a trained health workforce and unsatisfactory diagnosis and treatment in the facilities have been identified as some of the major contributors to high sepsis related mortality. Sepsis should be recognised early and treated aggressively. Therefore the quality of care at health facilities needs to be ensured to provide comprehensive management to all sepsis cases in order to prevent or decrease the onset of fatal consequences like septic shock and vital organ failure. A systematic review of thirteen studies in developing country context reported that the post discharge mortality rate may vary between 1% and 18%. The burden of morbidity and mortality of post discharged sepsis cases needs to be further evaluated in developing country context.

As representatives of the Global Sepsis Alliance (<http://globalsepsisalliance.com/>) we have had fruitful discussions with the WHO and the Global Burden for Disease experts who are eager to highlight the burden posed by sepsis. You can make a difference. Joining the World Sepsis Day initiative is a welcome start (<http://www.world-sepsis-day.org/>).

The Gut

IS-023 CLINICAL ASPECTS OF NEONATAL CHOLESTASIS

B Fischer. *Department of Pediatrics, Karolinska University Hospital CLINTEC, Stockholm, Sweden*

10.1136/archdischild-2014-307384.23

Neonatal cholestasis most often presents as prolonged jaundice, beyond 2 weeks of age. Such babies should always be investigated for conjugated hyperbilirubinemia and if this is detected further work-up for cholestasis is mandatory. Simultaneously, prothrombin time must be checked and pathologic levels need to be treated with intravenous vitamin K, to avoid detrimental bleedings. Neonatal cholestasis occurs in approximately 1 in 2500 term infants, the most common underlying diseases being

biliary atresia (BA), viral infections, α_1 -antitrypsin deficiency and other genetic cholestatic disorders.

Patients with BA most commonly present with jaundice and pale stools. Population based screening of all infants, by the use of stool colour cards sent to the parents can speed up the detection of cases with BA and thereby improve the success rate of the surgical procedure.

Once cholestasis is established, there is a need for a relatively rapid series of investigations. The reason for this "semiacute" management is to define certain situations where early treatment is lifesaving, such as septicemia, galactosemia or hypocortisolism, as well as to ensure timely surgical management in BA patients. A stepwise approach of investigations is suggested, where percutaneous liver biopsy is of value to select patients with suspected biliary atresia for laparotomy.

All patients, regardless of aetiology, should receive fat soluble vitamins as long as they are cholestatic. Although widely used, the effect of ursodeoxycholic acid is less well proven. However, it seems to be beneficiary in subsets of patient groups and the side effects are very minor.

The Heart

IS-024 FETAL CARDIAC INTERVENTIONS

¹G Tulzer, ²W Arzt. *¹Department of Pediatric Cardiology, Children's Heart Center Linz, Linz, Austria; ²Institute of Prenatal Medicine, Children's Heart Center Linz, Linz, Austria*

10.1136/archdischild-2014-307384.24

Fetal cardiac interventions are currently performed for critical aortic stenosis (CAS), pulmonary atresia intact septum (PA/IVS) and closed atrial septum in hypoplastic left heart syndrome (HLHS). The rational is to change fetal hemodynamics, prevent secondary damage and improve long term outcome at an acceptable risk for mother and fetus.

In fetal CAS the left ventricle (LV) dilates, shortening decreases and endocardial fibroelastosis develops. The most important issue is to identify those patients, who will benefit from this procedure.

Intervention has been shown to improve fetal hemodynamics. The risk of intrauterine death is around 12%. A biventricular outcome has been reported in 1/3 to 2/3 of these fetuses.

At the Children's Heart Centre Linz, 57 procedures were performed in 50 fetuses, with a success rate of 83% and a biventricular outcome after a median follow-up period of 43 months of 56%.

Intervention in fetal PA/IVS is technically more challenging. In our centre 12 attempts have been performed in 10 fetuses all with suprasystemic RV pressures. There was no serious complication so far. Successful intervention (in 8 fetuses) resulted in better RV filling and continuous, but slower than normal growth of tricuspid valve and RV. After 1 year, 4/5 newborns are biventricular and well, 1 child received a Glenn shunt. One 6 months old patient still has a Blalock Taussig shunt, 2 fetuses are still in-utero.

Intact atrial septum in fetal HLHS has to be treated with transseptal stent placement to prevent re-occlusion. Results however are still poor (Own experience with 5 procedures).

IS-025 FUNCTIONAL ECHOCARDIOGRAPHY: CORE COMPETENCY FOR THE NEONATOLOGIST?

N Evans. *Newborn Care, RPA Hospital and University of Sydney, Sydney, Australia*

10.1136/archdischild-2014-307384.25