

SUPPLEMENTARY MATERIAL**New-onset type 1 diabetes in Finnish children during the COVID-19 pandemic**

Heli Salmi *, MD, PhD (1, 2); Santtu Heinonen *, MD, PhD (1); Johanna Hästbacka, MD, PhD (1, 2);

Mitja Lääperi, MSc (1); Paula Rautiainen, MD, PhD (2); Päivi J. Miettinen, MD, PhD (1); Olli

Vapalahti, MD, PhD (3, 4, 5); Jussi Hepojoki, PhD; (3, 6); Mikael Knip, MD, PhD (1, 7)

*equal contribution

- (1) New Children's Hospital, Pediatric Research Center, University of Helsinki, and Helsinki University Hospital; Stenbäckinkatu 9, 00029 HUS, Helsinki, Finland
- (2) Department of Anesthesia and Intensive Care, New Children's Hospital, University of Helsinki and Helsinki University Hospital; Stenbäckinkatu 9, 00029 HUS, Helsinki, Finland
- (3) Department of Virology, Medicum, Faculty of Medicine, University of Helsinki; Haartmaninkatu 3, 00014, Helsinki, Finland
- (4) HUSLAB, Helsinki University Hospital, Helsinki, Finland
- (5) Department of Veterinary Biosciences, Faculty of Veterinary Medicine, University of Helsinki, Helsinki, Finland
- (6) Institute of Veterinary Pathology, Vetsuisse Faculty, University of Zürich, Zürich, Switzerland
- (7) Research Program for Clinical and Molecular Metabolism, Faculty of Medicine, University of Helsinki, Helsinki, Finland

This supplement forms part of the original submission.

We post it as supplied by the authors.

SUPPLEMENTARY METHODS

Patient selection for pediatric intensive care unit (PICU) admission

There are no absolute admission criteria to our PICU for children who do not require invasive mechanical ventilation, vasoactive therapy or extracorporeal support; admission is based on a joint evaluation by an emergency care paediatrician and a paediatric intensivist. A child with DKA is admitted if he or she is deemed to require constant monitoring of the level of consciousness, more than 2-4 hourly blood gases or immediate central venous access. Consequently, children with severe DKA (pH <7.10), severe hyperosmolarity or altered level of consciousness will, in most cases, be admitted to PICU.

Laboratory tests and data extraction

All laboratory parameters, except SARS-CoV-2 antibodies, were analysed by the Helsinki University Hospital (HUH) accredited centralized clinical laboratory HUSLAB. In 2016, the method for HbA1C measurement allowed exact measurements of values over 130 mmol/mol, whereas thereafter, they were reported as >130 mmol/mol. Thus, values exceeding 130 mmol/mol in 2016 were converted to 130 mmol/mol. Similarly, as β -Hydroxybutyric acid was occasionally measured using point of care test reporting values over 8 mmol/l as > 8 mmol/l, all values above 8 mmol/l were converted to 8 mmol/l. Duration of symptoms at the time of the diagnosis was categorized to the nearest entire week for up to 4 weeks, and to more than 4 weeks thereafter and cumulative incidence plots were created using survival survival and ggsurvminer packages[1,2].

SARS-CoV-2 antibody tests

Available serum samples were tested for the presence of IgG antibodies against SARS-CoV-2 spike protein using enzyme-linked immunosorbent assay (ELISA). SARS-CoV-2 spike protein was produced in Expi293 cells and ELISA was performed according to previously described protocols[3,4], with the exception that ELISA coating was 100 instead of 200 ng/well. The test has FDA authorization for emergency use in patient diagnostics [5]. One sample that gave a weak positive result in the ELISA test was further tested using a microneutralization assay[6].

SUPPLEMENTARY TABLES**Supplementary table 1. Admissions to paediatric intensive care unit due to new onset T1D**

during each study period (April 1 to October 31) 2016-2020. As the study period was 7 months each year, yearly population sizes at risk were scaled to 7-month person times and incidence was reported as rates per 100 000 person years. For pre-pandemic period 2016-2019 all events and scaled person years were pooled together.

T1D, type 1 diabetes

CI, confidence interval

Year	Events	Population	Scaled person years	Incidence rate per 100 000 person years (95% CI)
2016	6	370 948	216 386	2.77 (1.25-6.17)
2017	6	371 716	216 834	2.77 (1.24-6.16)
2018	9	371 223	216 547	4.16 (2.16-7.99)
2019	4	369 807	215 721	1.85 (0.70-4.94)
2016 - 2019	25	1 483 694	865 488	2.89 (1.95-4.27)
2020	20	366 754	213 940	9.35 (6.03-14.49)

Supplementary table 2. Children registered to Finnish pediatric diabetes registry due to new onset T1D during each study period (April 1 to October 31) 2016-2020. As the study period was 7 months each year, yearly population sizes at risk were scaled to 7-month person times and incidence was reported as rates per 100 000 person years. For pre-pandemic period 2016-2019 all events and scaled person years were pooled together.

T1D, type 1 diabetes

CI, confidence interval

Year	Events	Population	Scaled person years	Incidence rate per 100 000 person years (95% CI)
2016	58	253 210	147 706	39.3 (30.4-50.8)
2017	53	255 533	149 061	35.6 (27.2-46.5)
2018	58	257 141	149 999	38.7 (29.9-50.0)
2019	62	257 958	150 476	41.2 (32.1-52.8)
2016 - 2019	231	1 023 842	597 241	38.7 (34.0-44.0)
2020	84	257 158	150 009	56.0 (45.2-69.3)

FIGURE LEGEND

Supplementary figure 1. Cumulative incidences of T1D diagnoses and time since symptom onset for PICU admission (left panel) and patients registered to FPDR (right panel). The differences between the pre-pandemic and pandemic periods were not statistically significant (PICU cohort $P=0.09$; and FPDR cohort $P=0.29$).

T1D, type 1 diabetes; PICU, paediatric intensive care unit; FPDR, Finnish pediatric diabetes registry

REFERENCES

- 1 Albukadel K, Marcin K, Przemyslaw B, Scheipl F. Drawing Survival Curves using “ggplot2” [R package survminer version 0.4.9]. 2020.<https://CRAN.R-project.org/package=survminer>
- 2 Therneau T. A Package for Survival Analysis in R. R package version 3.2-10. 2021.<https://CRAN.R-project.org/package=survival>
- 3 Stadlbauer D, Amanat F, Chromikova V, *et al.* SARS-CoV-2 Seroconversion in Humans: A Detailed Protocol for a Serological Assay, Antigen Production, and Test Setup. *Curr Protoc Microbiol* 2020;**57**:e100.
- 4 Amanat F, Stadlbauer D, Strohmeier S, *et al.* A serological assay to detect SARS-CoV-2 seroconversion in humans. *Nat Med* 2020;**26**:1033–6.
- 5 U.S. Food and Drug Administration (FDA). Accelerated Emergency Use Authorization (EUA) Summary Covid-19 Elisa IgG Antibody Test. <https://www.fda.gov/media/137029/download> (accessed Oct, 28 2020).
- 6 Haveri A, Smura T, Kuivanen S, *et al.* Serological and molecular findings during SARS-CoV-2 infection: the first case study in Finland, January to February 2020. *Euro Surveill* 2020;**25**. doi:10.2807/1560-7917.ES.2020.25.11.2000266