

Table (electronic). Symptom-based score (adapted from references 11, 12, 13)

Symptom	Score										
Crying (°)	0 to 6	0 1 hour/day 1 1 - 1.5 hours/day 2 1.5 - 2 hours/day 3 2 to 3 hours/day 4 3 to 4 hours/day 5 4 to 5 hours/day 6 > 5 hours/day									
Regurgitation (27)	0 to 6	0 0-2 episodes/day 1 $\geq 3 - \leq 5$ of small volume 2 > 5 episodes of > 1 coffee spoon 3. > 5 episodes of \pm half of the feedings in < half of the feedings 4 continuous regurgitations of small volumes > 30 min after each feeding 5 regurgitation of half to complete volume of a feeding in at least half of the feedings 6. regurgitation of the complete volume after each feeding									
Stools (according to Bristol scale-28)	0 to 6	4 type 1 and 2 (hard stools) 0 type 3 and 4 (normal stools) 2 type 5 (soft stool) 4 type 6 (mushy/liquid stool, if unrelated to infection) 6 type 7 (watery stools)									
Dermatological symptoms	0 to 6	Atopic eczema <table style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th></th> <th>Head-neck-trunk</th> <th>Arms-hands-legs-feet</th> </tr> </thead> <tbody> <tr> <td>Absent</td> <td>0</td> <td>0</td> </tr> <tr> <td>Mild</td> <td>1</td> <td>1</td> </tr> </tbody> </table>		Head-neck-trunk	Arms-hands-legs-feet	Absent	0	0	Mild	1	1
	Head-neck-trunk	Arms-hands-legs-feet									
Absent	0	0									
Mild	1	1									

		Moderate 2 2
		Severe 3 3
	0 to 6	Urticaria (no 0 / yes 6)
Respiratory symptoms	0 to 3	0 no respiratory symptoms 1 mild symptoms 2 moderate symptoms 3 severe symptoms

Legend

(°) Crying was only considered if the child was crying for one week or more, assessed by the parents, without any other obvious cause.